S^+ – valued graphs and its Cartesian product

M. Sundar^{1*} and M. Chandramouleeswaran²

Abstract

In this paper, we introduce a new class of S –valued graphs, called S^+ –valued graphs by assigning the weights of the vertices as usual and edge weight is assigned by using the addition operation in the semiring.

Keywords

Semiring, S –valued graphs, Cartesian product of S –valued graphs.

AMS Subject Classification

16Y60, 05C25, 05C76.

¹ Department of Mathematics, Sree Sowdambika College of Engineering, Aruppukottai-626143, India.

² PG Department of Mathematics, Sri Ramanas College of Arts and Science for Women, Chettikurichi-626143, India.

*Corresponding author: ¹ sundar.mat14@gmail.com; ² moulee59@gmail.com

Article History: Received 21 December 2018 ; Accepted 11 February 2019

©2019 MJM.

Contents

1	Introduction	73
2	Preliminaries	73
3	S^+ –Valued Graphs	74
4	Cartesian Product of two S^+ –valued graphs	75
5	Generalized Cartesian Product of S^+ –valued graphs	76
6	Conclusion	78
	References	78

1. Introduction

In 2015, Chandramouleeswaran et.al introduced the concept of Semiring valued graphs simply called S –valued graphs [4]. In all the works done so far, the two binary operations in the semirings are not utilized. In this paper, we redefine the concept of S –valued graphs by assigning the weights to the vertices as usual and edges are assigned weights using the binary operation '+' in the semiring. We call this S –valued graph as S^+ –valued graph and it is denoted by G_S^+ . In [?] , we discussed the concept of generalized cartesian product of S –valued graphs by assigning the weights of the vertices of the two graphs from two different semirings. In this paper, we also discuss the concept of cartesian product of S^+ –valued graphs and study the regularity conditions satisfied by the S^+ –valued graphs.

2. Preliminaries

In this section, we recall some basic definitions related to S –valued graphs that are needed for our sequel.

Definition 2.1. [3] A semiring $(S, +, \cdot)$ is an algebraic system with a non-empty set S together with two binary operations $+$ and $\cdot$ such that

1. $(S, +)$ is a monoid.
2. $(S, \cdot)$ is a semigroup.
3. For all $a, b, c \in S$, $a \cdot (b + c) = a \cdot b + a \cdot c$ and $(a + b) \cdot c = a \cdot c + b \cdot c$
4. $0 \cdot x = x \cdot 0 = 0 \quad \forall x \in S$.

The element $0 \in S$ is the additive identity as well as the zero of the semiring S .

Definition 2.2. [3] Let $(S, +, \cdot)$ be a semiring. $\preceq$ is said to be a canonical pre-order if for $a, b \in S$, $a \preceq b$ if and only if there exists $c \in S$ such that $a + c = b$.

Definition 2.3. [4] Let $G = (V, E \subset V \times V)$ be a given graph with $V, E \neq \emptyset$. For any semiring $(S, +, \cdot)$, a semiring valued graph (or a S -valued graph) G^S is defined to be the graph $G^S = (V, E, \sigma, \psi)$ where $\sigma : V \rightarrow S$ and $\psi : E \rightarrow S$ is defined by

$$\psi(x, y) = \begin{cases} \min\{\sigma(x), \sigma(y)\} & \text{if } \sigma(x) \preceq \sigma(y) \text{ or } \sigma(y) \preceq \sigma(x) \\ 0 & \text{otherwise} \end{cases}$$

for every unordered pair (x, y) of $E \subset V \times V$. we call σ , a S –vertex set and ψ a S –edge set of the S –valued graph G^S .

Definition 2.4. [5] The open neighbourhood of v_i in G^S is defined as

$$N_S(v_i) = \{(v_j, \sigma(v_j)) \mid (v_i, v_j) \in E, \psi(v_i, v_j) \in S\}.$$

The closed neighbourhood of v_i in G^S is defined as

$$N_S[v_i] = N_S(v_i) \cup \{(v_i, \sigma(v_i))\}.$$

Definition 2.5. [5] The degree of a vertex v_i of the S -valued graph G^S is defined as $\deg_S(v_i) = (\sum_{v_j \in N_S(v_i)} \psi(v_i, v_j), l)$ where l is the number of edges incident with v_i .

Definition 2.6. [2] A S -valued graph G^S is said to be

1. S -vertex regular if $\sigma(v) = a, \forall v \in V$, for some $a \in S$.
2. S -edge regular if $\psi(u, v) = a, \forall (u, v) \in E$, for some $a \in S$.
3. S -regular if it is both S -vertex and S -edge regular.

Definition 2.7. [5] A S -valued graph G^S is said to be degree regular S -valued graph (d_S -regular graph) if

$$\deg_S(v) = (a, n) \forall v \in V, \text{ for some } a \in S \text{ and } n \in \mathbb{N}.$$

Definition 2.8. [6] The cartesian product of two graphs G and H is a graph, denoted by $G \square H$ whose vertex set is $V(G) \times V(H)$. Two vertices (g, h) and (g', h') are adjacent if $g = g'$ and $hh' \in E(H)$ or $gg' \in E(G)$ and $h = h'$. Thus $V(G \square H) = \{(g, h) \mid g \in V(G) \text{ and } h \in V(H)\}$,

$$E(G \square H) = \{(g, h)(g', h') \mid g = g', hh' \in E(H)$$

$$\text{or } gg' \in E(G), h = h'\}.$$

3. S^+ -Valued Graphs

In this section, we introduce the notion of S^+ -valued graphs and illustrate it with some examples.

Definition 3.1. Let $G = (V, E \subseteq V \times V)$ be a given graph with $V, E \neq \emptyset$. For any semiring $(S, +, \cdot)$, a S^+ -valued graph G^{S^+} is defined to be the graph $G^{S^+} = (V, E, \sigma, \psi)$, where $\sigma : V \rightarrow S$ and $\psi : E \rightarrow S$ is defined to be $\psi(x, y) = (\sigma(x) + \sigma(y))$, for every unordered pair $(x, y) \in E \subseteq V \times V$.

Example 3.2. Consider the semiring $S = B(5, 3) = (\{0, 1, 2, 3, 4\}, \oplus, \odot)$ with the binary operations ' $\oplus$ ' and ' $\odot$ ' defined by

$$a \oplus b = \begin{cases} a + b & \text{if } a + b \leq 4 \\ c & \text{if } c \equiv a + b \pmod{2}, 3 \leq c \leq 4 \end{cases} \text{ and}$$

$$a \odot b = \begin{cases} a \cdot b & \text{if } a \cdot b \leq 4 \\ c & \text{if } c \equiv a \cdot b \pmod{2}, 3 \leq c \leq 4 \end{cases}$$

These operations can be represented by the following Cayley Tables:

$\oplus$	0	1	2	3	4
0	0	1	2	3	4
1	1	2	3	4	3
2	2	3	4	3	4
3	3	4	3	4	3
4	4	3	4	3	4

$\odot$	0	1	2	3	4
0	0	0	0	0	0
1	0	1	2	3	4
2	0	2	4	4	4
3	0	3	4	3	4
4	0	4	4	4	4

Consider the graph $G = (V, E)$, where $V = \{v_1, v_2, v_3, v_4, v_5, v_6\}$

$$\text{and } E = \{(v_1, v_2), (v_2, v_3), (v_3, v_4), (v_3, v_6),$$

$$(v_4, v_5), (v_4, v_6), (v_5, v_1), (v_5, v_6)\}.$$

Corresponding to the graph G , we define the S^+ -valued graph G^{S^+} as follows:

Define $\sigma : V \rightarrow S$ by

$$\sigma(v_1) = 2; \sigma(v_2) = 1; \sigma(v_3) = 2;$$

$$\sigma(v_4) = 3; \sigma(v_5) = 4 \sigma(v_6) = 3.$$

Therefore the S^+ -vertex set of $G^{S^+} = \{1, 2, 3, 4\}$.

Now define $\psi : E \rightarrow S$ as follows:

$$\psi((v_1, v_2)) = (\sigma(v_1) + \sigma(v_2)) = (2 + 1) = 3.$$

$$\psi((v_2, v_3)) = (\sigma(v_2) + \sigma(v_3)) = (1 + 2) = 3.$$

$$\psi((v_3, v_4)) = (\sigma(v_3) + \sigma(v_4)) = (2 + 3) = 3.$$

$$\psi((v_3, v_6)) = (\sigma(v_3) + \sigma(v_6)) = (2 + 3) = 3.$$

$$\psi((v_4, v_5)) = (\sigma(v_4) + \sigma(v_5)) = (3 + 4) = 3.$$

$$\psi((v_4, v_6)) = (\sigma(v_4) + \sigma(v_6)) = (3 + 3) = 4.$$

$$\psi((v_5, v_1)) = (\sigma(v_5) + \sigma(v_1)) = (4 + 2) = 4.$$

$$\psi((v_5, v_6)) = (\sigma(v_5) + \sigma(v_6)) = (4 + 3) = 3.$$

Then the S^+ -edge set of $G^{S^+} = \{3, 4\}$.

The graph G and its corresponding S^+ -valued graphs $G_1^{S^+}, G_2^{S^+}, G_3^{S^+}$ are given below:

G

$G_1^{S^+}$

$G_2^{S^+}$

$G_3^{S^+}$

Remark 3.3. From the above example, we observe that for a given graph G its S^+ -valued graph G^{S^+} is not unique.

Definition 3.4. A S -valued graph G^S is said to be

1. S^+ -vertex regular if $\sigma(v)$ is same $\forall v \in V$.
2. S^+ -edge regular if $\psi(u, v)$ is same $\forall (u, v) \in E$.
3. S^+ -regular if it is both S^+ -vertex and S^+ -edge regular.

4. Cartesian Product of two S^+ -valued graphs

In this section, we discuss the cartesian product of two S^+ -valued graphs $G_1^{S^+}$ and $G_2^{S^+}$. Also we discuss the regularity conditions satisfied by the cartesian product of two S^+ -valued graphs.

Definition 4.1. Let $G_1^{S^+} = (V_1, E_1, \sigma_1, \psi_1)$ and $G_2^{S^+} = (V_2, E_2, \sigma_2, \psi_2)$ be two S^+ -valued graphs. Then the Cartesian product of $G_1^{S^+}, G_2^{S^+}$ is denoted by

$$G_{\square}^{S^+} = (V = V_1 \times V_2, E = E_1 \times E_2, \sigma = \sigma_1 \times \sigma_2, \psi = \psi_1 \times \psi_2)$$

where $V = \{w_{ij} = (v_i, u_j) \mid v_i \in V_1 \text{ and } u_j \in V_2\}$ and two vertices w_{ij} and w_{kl} are adjacent if $i = k$ and $u_j u_l \in E_2$ or $j = l$ and $v_i v_k \in E_1$. That is $E = \{e_{ij}^{kl} \mid \text{if either } i = k \text{ and } u_j u_l \in E_2 \text{ or } j = l \text{ and } v_i v_k \in E_1\}$, Define $\sigma : V_1 \times V_2 \rightarrow S \times S$ by

$$\sigma(v_i, u_j) = \sigma(w_{ij}) = (\sigma_1(v_i) + \sigma_2(u_j))$$

$$\psi : E \rightarrow S \text{ by } \psi(e_{ij}^{kl}) = \psi((v_i, u_j)(v_k, u_l))$$

$$= \begin{cases} \sigma_1(v_i) + \psi_2(u_j, u_l) & \text{if } i = k \text{ and } u_j u_l \in E_2 \\ \psi_1(v_i, v_k) + \sigma_2(u_j) & \text{if } j = l \text{ and } v_i v_k \in E_1 \end{cases}$$

Example 4.2. Consider the semiring $S = (\{0, a, b, c\}, +, \cdot)$ with the binary operations '+' and '·' defined by the following Cayley tables.

+	0	a	b	c
0	0	a	b	c
a	a	b	c	c
b	b	c	c	c
c	c	c	c	c

·	0	a	b	c
0	0	0	0	0
a	0	a	b	c
b	0	b	c	c
c	0	c	c	c

Consider the two S^+ -valued graphs $G_1^{S^+}$ and $G_2^{S^+}$. Then the cartesian product $G_{\square}^{S^+} = G_1^{S^+} \square G_2^{S^+}$ is given by $G_{\square}^{S^+} = (V, E, \sigma, \psi)$ where $V = \{w_{11}, w_{12}, w_{13}, w_{21}, w_{22}, w_{23}\}$, $E = \{e_{21}^{11}, e_{12}^{11}, e_{13}^{11}, e_{22}^{12}, e_{13}^{12}, e_{23}^{13}, e_{22}^{21}, e_{23}^{21}, e_{23}^{22}\}$

Theorem 4.3. The cartesian product of two S^+ -vertex regular graph is S^+ -vertex regular.

Proof: Let $G_1^{S^+} = (V_1, E_1, \sigma_1, \psi_1)$ and $G_2^{S^+} = (V_2, E_2, \sigma_2, \psi_2)$ be two given S^+ -vertex regular graphs.

Claim: $G_{\square}^{S^+}$ is S^+ -vertex regular

Since $G_1^{S^+}$ and $G_2^{S^+}$ are S^+ -vertex regular graphs, $\sigma_1(v_i) = s_1$ and $\sigma_2(u_j) = s_2$, for some $s_1, s_2 \in S$.

Now By definition, $\sigma(w_{ij}) = \sigma_1(v_i) + \sigma_2(u_j) = s_1 + s_2 = s_3$, for some $s_3 \in S$.

This implies that $\sigma(w_{ij})$ is equal for all $w_{ij} \in V$, for all i, j .

Hence $G_{\square}^{S^+} = G_1^{S^+} \square G_2^{S^+}$ is S^+ -vertex regular.

Example 4.4. The cartesian product of two S^+ -edge regular graphs is not S^+ -edge regular. Consider the semiring as in the example 3.2 and the following two S^+ -edge regular graphs:

Clearly $G_{\square}^{S^+}$ is not S^+ -edge regular.

Remark 4.5. In cartesian product of S -valued graphs, the cartesian product of two S -vertex regular graphs is S -edge regular. But from the above example, we observe that, the cartesian product of two S^+ -vertex regular graphs is need not be S^+ -edge regular.

Theorem 4.6. The cartesian product of two S^+ -edge regular graphs is S^+ -edge regular only if the semiring under consideration is additively idempotent.

Proof: Let $G_1^{S^+} = (V_1, E_1, \sigma_1, \psi_1)$ and $G_2^{S^+} = (V_2, E_2, \sigma_2, \psi_2)$ be two given S^+ -edge regular graphs.

Claim: $G_1^{S^+} \square G_2^{S^+}$ is S^+ -edge regular
By definition, $\psi(e_{ij}^{kl}) = \psi((v_i, u_j)(v_k, u_l))$

$$= \begin{cases} \sigma_1(v_i) + \psi_2(u_j, u_l) & \text{if } i = k \text{ and } u_j u_l \in E_2 \\ \psi_1(v_i, v_k) + \sigma_2(u_j) & \text{if } j = l \text{ and } v_i v_k \in E_1 \end{cases}$$

$$= \begin{cases} \sigma_1(v_i) + (\sigma_2(u_j) + \sigma_2(u_l)) & \text{if } i = k \text{ and } u_j u_l \in E_2 \\ (\sigma_1(v_i) + \sigma_1(v_k)) + \sigma_2(u_j) & \text{if } j = l \text{ and } v_i v_k \in E_1 \end{cases}$$

$$= \begin{cases} \sigma_1(v_i) + \sigma_2(u_j) & \text{if } i = k \text{ and } u_j u_l \in E_2 \\ \sigma_1(v_i) + \sigma_2(u_j) & \text{if } j = l \text{ and } v_i v_k \in E_1 \end{cases}$$

($\because S$ is additively idempotent)

This is true for all $e_{ij}^{kl} \in E$, therefore $G_1^{S^+} \square G_2^{S^+}$ is S^+ -edge regular.

Corollary 4.7. If S is an additively idempotent semiring and $G_1^{S^+}, G_2^{S^+}$ are two S^+ -vertex regular graphs, then their cartesian product $G_{\square}^{S^+}$ is S^+ -regular.

Example 4.8. The cartesian product of two d_S -regular S^+ -valued graphs is not d_S -regular.

Consider the following two d_S -regular S^+ -valued graphs $G_1^{S^+}$ and $G_2^{S^+}$ and their Cartesian product

Here $d_S(w_{11}) = (3, 2); d_S(w_{21}) = (4, 2); d_S(w_{22}) = (3, 2); d_S(w_{12}) = (4, 2)$.

We observe that $G_{\square}^{S^+}$ is not d_S -regular.

5. Generalized Cartesian Product of S^+ -valued graphs

In this section, we discuss the generalized cartesian product of two S^+ -valued graphs $G_1^{S^+}$ and $G_2^{S^+}$ assuming S -values from two different semirings $(S_1, +, \cdot)$ and $(S_2, +, \cdot)$. Also we discuss the regularity conditions satisfied by the generalized cartesian product of two S^+ -valued graphs.

Definition 5.1. Let $G_1^{S^+} = (V_1, E_1, \sigma_1, \psi_1)$ and $G_2^{S^+} = (V_2, E_2, \sigma_2, \psi_2)$ be two S_1^+ and S_2^+ valued graphs. Then the generalized cartesian product of these two S^+ -valued graphs is denoted by $G_{\square}^{S_1^+ \times S_2^+}$, defined by

$$G_1^{S^+} \square G_2^{S^+} = (V = V_1 \times V_2, E = E_1 \times E_2, \sigma = \sigma_1 \times \sigma_2, \psi = \psi_1 \times \psi_2),$$

where $V = \{w_{ij} = (v_i, u_j) \mid v_i \in V_1 \text{ and } u_j \in V_2\}$ and two vertices w_{ij} and w_{kl} are adjacent if $i = k$ and $u_j u_l \in E_2$ or $j = l$ and $v_i v_k \in E_1$. That is

$E = \{e_{ij}^{kl} \mid \text{if either } i = k \text{ and } u_j u_l \in E_2 \text{ or } j = l \text{ and } v_i v_k \in E_1\}$,
Define $\sigma : V_1 \times V_2 \rightarrow S_1 \times S_2$ by $\sigma(w_{ij}) = (\sigma_1(v_i), \sigma_2(u_j))$ and $\psi : E \rightarrow S$ by $\psi(e_{ij}^{kl}) = \psi((v_i, u_j)(v_k, u_l)) = (\sigma_1(v_i) + \sigma_1(v_k), \sigma_2(u_j) + \sigma_2(u_l))$

Example 5.2. Consider the semiring $S_1 = (\{0, a, b, c\}, +, \cdot)$ with the binary operations '+' and ' $\cdot$ ' defined by the following Cayley tables.

+	0	a	b	c
0	0	a	b	c
a	a	b	c	c
b	b	c	c	c
c	c	c	c	c

$\cdot$	0	a	b	c
0	0	0	0	0
a	0	a	b	c
b	0	b	c	c
c	0	c	c	c

and $S_2 = (\{0, 1, 2, 3\}, +, \cdot)$ with the binary operations '+' and ' $\cdot$ ' defined by the following Cayley tables.

+	0	1	2	3
0	0	1	2	3
1	1	1	2	3
2	2	2	2	2
3	3	3	2	2

$\cdot$	0	1	2	3
0	0	0	0	0
1	0	0	1	0
2	0	1	2	3
3	0	0	3	3

For the two S^+ -valued graphs $G_1^{S_1^+}$ and $G_2^{S_2^+}$ in figure:1 and figure:2, their generalized cartesian product $G_{\square}^{S_1^+ \times S_2^+} = G_1^{S_1^+} \square G_2^{S_2^+}$ is given in figure:3.

Figure 1

Figure 2

Figure 3

Theorem 5.3. The generalized Cartesian product of two S^+ -vertex regular graphs is S^+ -edge regular.

Proof: Given S_1 and S_2 are two semirings.

Let $G_1^{S_1^+} = (V_1, E_1, \sigma_1, \psi_1)$ and $G_2^{S_2^+} = (V_2, E_2, \sigma_2, \psi_2)$ be two S^+ -vertex regular graphs.

Claim: $G_{\square}^{S_1^+ \times S_2^+} = G_1^{S_1^+} \square G_2^{S_2^+}$ is S^+ -edge regular

Now by definition, $\sigma(w_{ij}) = (\sigma_1(v_i), \sigma_2(u_j))$.

Since $G_1^{S_1^+}$ is S_1 regular, then $\sigma_1(v_i) = s_1$, for all $v_i \in V_1$ and for some $s_1 \in S_1$ and $G_2^{S_2^+}$ is S_2 regular, then $\sigma_2(u_j) = s_2$, for all $u_j \in V_2$ and for some $s_2 \in S_2$

Now $\psi(e_{ij}^{kl}) = \psi((v_i, u_j)(v_k, u_l))$

$$\begin{aligned} &= (\sigma_1(v_i) + \sigma_1(v_k), \sigma_2(u_j) + \sigma_2(u_l)) \\ &= (s_1 + s_1, s_2 + s_2) \\ &= (s_3, s_4), \text{ for some } s_3 \in S_1 \text{ and } s_4 \in S_2. \end{aligned}$$

This is true for all $e_{ij}^{kl} \in E$ and for some $s_1 \in S_1, s_2 \in S_2$

Hence $G_{\square}^{S_1^+ \times S_2^+}$ is S^+ -edge regular.

Corollary 5.4. If S_1 and S_2 are any two semirings and $G_1^{S_1^+}, G_2^{S_2^+}$ are two S^+ -vertex regular graphs, then their product is $G_{\square}^{S_1^+ \times S_2^+}$ is S^+ -regular.

Remark 5.5. If the semiring under consideration are additively idempotent, then both S^+ -regular and S -regular are one and the same.

Example 5.6. The generalized cartesian product of two S^+ -edge regular S -valued graphs need not be S^+ -edge regular. This is illustrated in the following example. Consider the semirings S_1 and S_2 as in the example 5.2.

Consider the two S^+ -valued graphs $G_1^{S_1^+}$ and $G_2^{S_2^+}$ their generalized cartesian product $G_{\square}^{S_1^+ \times S_2^+} = G_1^{S_1^+} \square G_2^{S_2^+}$ is given below:

Figure 1

Figure 2

Figure 3

Example 5.7. The generalized cartesian product of two d_S -regular S^+ -valued graphs is not d_S -regular.

Consider the semiring $S_1 = (\{0, a, b, c\}, +, \cdot)$ with the binary operations '+' and '·' defined by the following Cayley

tables.

+	0	a	b	c
0	0	a	b	c
a	a	a	b	c
b	b	b	b	b
c	c	c	b	b

·	0	a	b	c
0	0	0	0	0
a	0	0	a	0
b	0	a	b	c
c	0	0	c	c

$S_2 = B(5, 3) = (\{0, 1, 2, 3, 4\}, \oplus, \odot)$ with the Cayley Tables:

$\oplus$	0	1	2	3	4
0	0	1	2	3	4
1	1	2	3	4	3
2	2	3	4	3	4
3	3	4	3	4	3
4	4	3	4	3	4

$\odot$	0	1	2	3	4
0	0	0	0	0	0
1	0	1	2	3	4
2	0	2	4	4	4
3	0	3	4	3	4
4	0	4	4	4	4

Consider the following two d_S -regular S^+ -valued graphs $G_1^{S^+}$ and $G_2^{S^+}$ and their cartesian product $G_{\square}^{S^+ \times S^+}$

We observe that $G_{\square}^{S^+ \times S^+}$ is not d_S -regular.

Theorem 5.8. *The generalized cartesian product of two d_S -regular S^+ -valued graphs is d_S -regular only if the corresponding factors are S^+ -vertex regular and the semirings under consideration are additively idempotent.*

6. Conclusion

In this paper, we present a new class of S -valued graphs called S^+ -valued graphs, analogous to the concept of S -valued graphs and discussed the regularity conditions on their cartesian product. Further this theory can be developed to discuss the concepts irregularity, vertex domination, edge domination and colouring, analogous to the S -valued graphs.

References

- [1] Chris Godsfil and Gordon Royle, *Algebraic Graph Theory*, Springer, 2001.
- [2] Jeyalakshmi. S, Rajkumar. M, and Chandramouleeswaran. M, Regularity on S -valued graphs, *Global J. of Pure and Applied Maths.*, 2(5)(2015), 2971–2978.
- [3] Jonathan S Golan *Semirings and Their Applications*, Kluwer Academic Publishers, London, 1997.
- [4] Rajkumar. M, Jeyalakshmi. S. and Chandramouleeswaran. M, Semiring Valued Graphs, *International Journal of Math. Sci. and Engg. Appls.*, 9(3)(2015), 141–152.
- [5] Rajkumar and Chandramouleeswaran. M, Degree Regular S -Valued Graphs, *Mathematical Sciences International Research Journal*, 4(2)(2012), 326–328.
- [6] Richard Hammack, Wilfried Imrich and Sandy Klavzar, *Handbook of Product Graphs*, University of Ljubljana and University of Maribor, Slovenia, 1990.
- [7] Sundar.M and Chandramouleeswaran. M, Cartesian product of two S -valued graphs, *Global Journal of Mathematical Sciences: Theory and Practical.*, 9(3)(2017), 347–355.

 ISSN(P):2319 – 3786
 Malaya Journal of Matematik
 ISSN(O):2321 – 5666