Dominator coloring on anti fuzzy graph

R. Muthuraj¹ and A. Sasireka^{2*}

Abstract

In this paper, the concept of dominator coloring on anti fuzzy graph is introduced and the fuzzy dominator chromatic number of anti fuzzy graph is characterized. This concept is applied on anti cartesian product of anti fuzzy graphs and obtained the results on them.

Keywords

Anti fuzzy graph, Domination number, Dominator coloring.

AMS Subject Classification

05C72, 05C15, 05C69, 05C76, 05A18.

¹ Assistant Professor, PG & Research Department of Mathematics, H.H. The Rajah's College, Pudukkottai - 622001, Tamil Nadu, India.

² Assistant Professor, Department of Mathematics, PSNA College of Engineering and Technology, Dindigul-624 622, Tamil Nadu, India.

*Corresponding author: ¹ rmr1973@yahoo.co.in; ² sasireka.pсна@gmail.com

Article History: Received 22 November 2018; Accepted 09 May 2019

©2019 MJM.

Contents

1	Introduction	217
2	Preliminaries	217
3	Dominator Coloring on Strong Anti Fuzzy Graph	218
4	Dominator coloring on Anti Fuzzy Graph with weak edges	221
5	Dominator Coloring on Anti Cartesian Product of Anti Fuzzy Graphs	221
6	Conclusion	222
	References	222

1. Introduction

Dominator coloring plays a vital role in biomedical applications such as predicting the cancer cells via buckyball and telecommunication networks, distributed computing via circulant graph. In 2005, Munoz et.al., delivered the idea of coloring on fuzzy graph[9] and described the chromatic number of fuzzy graph. In 2006, Raluca Michelle Gera established the fuzzy dominator coloring[6]. In the same time, Eslachi and Onagh pioneered the concept of fuzzy chromatic number as the number of partition of the color classes[3]. In 2015, R.Jahir Hussian and K.S.Kanzuli Fathima conferred fuzzy dominator coloring of fuzzy graph[7]. In 2012, Muhammad Akram introduced the concept of an anti fuzzy structures on graphs and characterized some basic concept of con-

nected anti fuzzy graph[8]. In 2016, R.Seethalakshmi and R.B.Gnanajothi introduced the definition of anti podal on anti fuzzy graph[20]. In 2017, this concept further developed by R.Muthuraj and A. Sasireka[10–12]. They introduced some types of anti fuzzy graphs and applied some operations on them and also derived some domination parameter on anti fuzzy graph[14, 15]. In 2018, Rifayathali et.al.,[17, 19] introduced the concept of coloring on anti fuzzy graph. In this paper, anti fuzzy graphs are considered without loops and multiple edges. V is a finite and non empty set. m and n are the number of vertices and edges of an anti fuzzy graph. The dominator coloring is defined on anti fuzzy graph. These concepts are applied on anti cartesian product of anti fuzzy graphs.

2. Preliminaries

In this section, basic concepts of anti fuzzy graph are discussed. Notations and more formal definitions are followed as in [3, 8, 10–12, 14, 16, 20].

Definition 2.1. [20] An anti fuzzy graph $G_A = (\sigma, \mu)$ is a pair of functions $\sigma : V \rightarrow [0, 1]$ and $\mu : V \times V \rightarrow [0, 1]$, with $\mu(u, v) \geq \sigma(u) \vee \sigma(v)$ for all $u, v \in V$.

Note. μ is considered as reflexive and symmetric. In all examples σ is chosen suitably. i.e., undirected anti fuzzy graphs are only considered. Without loss of generality, let us simply use the letter G_A to denote an anti fuzzy graph.

Definition 2.2. [8] Let $G_A=(V,\sigma,\mu)$ be an anti fuzzy graph. The order p of G_A is defined as $p = \sum_{x \in V} \sigma(x)$. The size q is, $q = \sum_{xy \in E} \mu(x,y)$. It is denoted by $O(G)$ and $S(G)$.

Definition 2.3. [12] Two vertices u and v in G_A are called adjacent if $(1/2) [\sigma(u) \vee \sigma(v)] \leq \mu(u,v)$.

Definition 2.4. [10] The anti complement of anti fuzzy graph $G_A=(\sigma,\mu)$ is an anti fuzzy graph $\bar{G}_A = (\bar{\sigma}, \bar{\mu})$ where $\bar{\sigma} = \sigma$ and $\bar{\mu}(u,v) = \mu(u,v) - (\sigma(u) \vee \sigma(v))$ for all u,v in V .

Definition 2.5. [20] An anti fuzzy graph $G_A = (\sigma, \mu)$ is a strong anti fuzzy graph of $\mu(u,v) = \sigma(u) \vee \sigma(v)$ for all $(u,v) \in \mu^*$ and G_A is a complete anti fuzzy graph if $\mu(u,v) = \sigma(u) \vee \sigma(v)$ for all $(u,v) \in \mu^*$ and $u,v \in \sigma^*$. Two vertices u and v are said to be neighbors if $\mu(u,v) > 0$.

Definition 2.6. [12] An edge $e = \{u, v\}$ of an anti fuzzy graph G_A is called an effective edge if $\mu(u,v) = \sigma(u) \vee \sigma(v)$.

Definition 2.7. [11] u is a vertex in an anti fuzzy graph G_A then $N(u) = \{v: (u,v) \text{ is an effective edge}\}$ is called the open neighborhood of u and $N[u] = N(u) \cup \{u\}$ is called closed neighborhood of u .

Definition 2.8. [8, 11] A path P_A in an anti fuzzy graph is a sequence of distinct vertices $u_0, u_1, u_2, \dots, u_n$ such that $\mu(u_{i-1}, u_i) > 0, 1 \leq i \leq n$. Here $n \geq 0$ is called the length of the path P_A . The consecutive pairs (u_{i-1}, u_i) are called the edges of the path.

Definition 2.9. [8] A cycle in G_A is said to be an anti fuzzy cycle if it contains more than one weakest vertex. It is denoted by C_A .

Definition 2.10. [11] Let $G_A^* = G_{A_1}^* \times G_{A_2}^* = (V, E')$ be the anti cartesian product of anti fuzzy graphs where $V = V_1 \times V_2$ and $E' = \{(u_1, u_2), (u_1, v_2) \mid u_1 \in V_1, (u_2, v_2) \in E_2\} \cup \{(u_1, w_2), (v_1, w_2) \mid w_2 \in V_2, (u_1, v_1) \in E_1\}$. Then the anti cartesian product of two anti fuzzy graphs, $G_A = G_{A_1} \times G_{A_2}$: $(\sigma_1 \times \sigma_2, \mu_1 \times \mu_2)$ is an anti fuzzy graph and is defined by

$(\sigma_1 \times \sigma_2)(u_1, u_2) = \max \{\sigma_1(u_1), \sigma_2(u_2)\}$ for all $(u_1, u_2) \in V$,

$(\mu_1 \times \mu_2)((u_1, u_2), (u_1, v_2)) = \max \{\sigma_1(u_1), \mu_2(u_2, v_2)\}$ for all $u_1 \in V_1$ and $(u_2, v_2) \in E_2$,

$(\mu_1 \times \mu_2)((u_1, w_2), (v_1, w_2)) = \max \{\sigma_2(w_2), \mu_1(u_1, v_1)\}$ for all $w_2 \in V_2$ and $(u_1, v_1) \in E_1$.

Then the anti fuzzy graph $G_A = (\sigma_1 \times \sigma_2, \mu_1 \times \mu_2)$ is said to be the anti cartesian product of anti fuzzy graphs $G_{A_1} = (\sigma_1, \mu_1)$ and $G_{A_2} = (\sigma_2, \mu_2)$.

Definition 2.11. [12] Every vertex in an anti fuzzy graph G_A has unique fuzzy values then G_A is said to be v -nodal anti fuzzy graph. i.e. $\sigma(u) = c$ for all $u \in V(G_A)$.

Definition 2.12. [12] Every edge in an anti fuzzy graph G_A has unique fuzzy values then G_A is said to be e -nodal anti fuzzy graph. i.e. $\mu(u,v) = c$ for all $(u,v) \in E(G_A)$.

Definition 2.13. [12] Every vertices and edges in an anti fuzzy graph G_A have the unique fuzzy values then G_A is called as uninodal anti fuzzy graph. If $\sigma(u) = c_1$ and $\mu(u,v) = c_2$ in an anti fuzzy graph G_A then G_A is called as binodal anti fuzzy graph.

Definition 2.14. [11, 14] The strong neighbourhood of an edge e_i in an anti fuzzy graph G_A is $N_s(e_i) = \{e_j \in E(G) \mid e_j \text{ is an effective edge with } \forall N(e_i) \text{ in } G_A \text{ and adjacent to } e_i\}$

Definition 2.15. [14] An edge $e = \{u,v\}$ of an anti fuzzy graph G_A is called an weak edge if $\mu(u,v) \neq \sigma(u) \vee \sigma(v)$.

Definition 2.16. [16] G_A is an anti fuzzy graph and $u, v \in V(G_A)$. If v is said to be a support vertex in G_A then v is adjacent to at least one end vertex u .

Definition 2.17. [17] Let G_A be an anti fuzzy graph. A family $c = \{c_1, c_2, \dots, c_k\}$ of fuzzy sets as a set V is called a k -vertex coloring of $G_A = (V, \sigma, \mu)$ if

1. $\vee c_i(x) = \sigma(x)$
2. $c_i \wedge c_j = 0$
3. For every strong edge xy of G_A , $\min\{c_i \sigma((x)), c_j \sigma((y))\} = 0, (1 \leq i \leq k)$. The least value of k for which G has k -vertex coloring denoted by $\chi(G_A)$, is called the chromatic number of the anti fuzzy graph G_A .

3. Dominator Coloring on Strong Anti Fuzzy Graph

In this section, dominator coloring is applied on strong anti fuzzy graph and the fuzzy dominator chromatic number on connected anti fuzzy graph which has strong edges only is determined. The membership value of the color class is 1. So, assigning a color to the vertices of an anti fuzzy graph as 1,2,3,... instead of Red(R), Green(G), Blue(B), Yellow(Y), etc. That is, every number(1,2,3,...) represent a one color class of G_A and the fuzzy value of color class is assigned the value 1. Depending upon the nature of dominator coloring, a vertex is classified as χ_{fd} -good and C-bad vertex. By using this characterization the anti fuzzy graph is also classified as χ_{fd} -good and χ_{fd} -excellent. The bounds are obtained for them and derived some theorems of dominator chromatic number on anti fuzzy graph.

Definition 3.1. A colored anti fuzzy graph G_A is an anti fuzzy graph in which each vertex of G_A is assigned a color from color class c . A properly colored anti fuzzy graph is a colored anti fuzzy graph whose color assignments conform to the coloring rules applied to the anti fuzzy graph. Fuzzy chromatic number of an anti fuzzy graph G_A is the least number of distinct colors and is denoted by $\chi(G_A)$.

Definition 3.2. A fuzzy dominator coloring of an anti fuzzy graph G_A is a proper fuzzy coloring in which each vertex of G_A dominates every vertex of atleast some color class.

Fuzzy dominator chromatic number of an anti fuzzy graph is the minimum number of color classes in a dominator fuzzy coloring of G_A . It is denoted by $\chi_{fd}(G_A)$.

Definition 3.3. C is the partition of independent sets of vertices of an anti fuzzy graph G_A in which $u \in D$ and $v \in V \setminus D$ is in distinct partition of color class C and each vertex of G_A dominates every vertex of atleast one color class. Such C is called the anti fuzzy dominator partition of G_A .

Example 3.4.

Figure 1. (a) Anti Fuzzy Graph G_A with dominator coloring; (b) Anti fuzzy graph G_A with coloring

In Figure 1(a), G_A is an anti fuzzy graph with proper coloring. The color classes of Figure 1(a) are c_1, c_2, c_3 and c_4 . Where $c_1 = \{v_1\}$, $c_2 = \{v_4\}$, $c_3 = \{v_2, v_5\}$ and $c_4 = \{v_3, v_6\}$. The vertices v_1 and v_4 dominates the remaining vertices and the vertices in the color class of c_3 and c_4 dominate all the vertices of atleast any one of color classes c_1 and c_2 . Therefore, Figure 1(a) - G_A is an anti fuzzy graph with proper dominator coloring.

In Figure 1(b), G_A is an anti fuzzy graph with proper coloring and the color classes are c_1, c_2 and c_3 . Where $c_1 = \{v_1, v_4\}$, $c_2 = \{v_3, v_6\}$ and $c_3 = \{v_2, v_5\}$. No vertex of any color classes dominate atleast one color class of G_A . Therefore, Figure 1(b) - G_A is not a proper dominator coloring.

Definition 3.5. A vertex $v \in V(G_A)$ is called χ_{fd} -good if a vertex v belongs to some C -partition (χ -partition) otherwise v is said to be C -bad vertex.

An anti fuzzy graph G_A is called χ_{fd} -excellent anti fuzzy graph if every vertex of G_A is χ_{fd} -good.

Example 3.6. Now, consider the following Figure 2, to determine the fuzzy dominator chromatic number of an anti fuzzy graph.

In Figure 2, the dominating set of G_A is $D = \{v_1, v_7\}$.

The color class is $c = \{c_1, c_2, c_3, c_4\} = \{\{v_1\}^1, \{v_7\}^1, \{v_2, v_4, v_5\}^1, \{v_3, v_6, v_8\}^1\}$

Therefore, $\chi_{fd}(G_A) = 1 + 1 + 1 + 1 = 4$.

$$c_1(v_i) = \begin{cases} 0.8, & \text{if } i = 1 \\ 0, & \text{otherwise} \end{cases}$$

$$c_2(v_i) = \begin{cases} 0.7, & \text{if } i = 7 \\ 0, & \text{otherwise} \end{cases}$$

Figure 2. An Anti Fuzzy Graph G_A

$$c_3(v_i) = \begin{cases} 0.5, & \text{if } i = 2, 5 \\ 0.3, & \text{if } i = 4 \\ 0, & \text{otherwise} \end{cases}$$

$$c_4(v_i) = \begin{cases} 0.1, & \text{if } i = 6 \\ 0.4, & \text{if } i = 3 \\ 0.2, & \text{if } i = 8 \\ 0, & \text{otherwise} \end{cases}$$

Here the vertices $v_1, v_2, \dots, v_8$ dominate atleast one of the color classes are c_1, c_2 of G_A . Therefore, $\chi_{fd}(G_A) = 4$.

The anti fuzzy dominator partition of Figure 2 is,

$C = \{C_1, C_2, C_3, C_4\}$.

Where $C_1 = (\{v_1\}, \{v_7\}, \{v_2, v_4, v_5\}, \{v_3, v_6, v_8\})$

$C_2 = (\{v_2\}, \{v_8\}, \{v_4, v_5, v_7\}, \{v_1, v_3, v_6\})$

$C_3 = (\{v_3\}, \{v_5\}, \{v_2, v_4, v_7\}, \{v_1, v_6, v_8\})$

$C_4 = (\{v_4\}, \{v_6\}, \{v_2, v_7, v_5\}, \{v_3, v_1, v_8\})$

Every vertex of G_A appears in a singleton in χ -partition.

Hence G_A is χ -excellent.

Every vertex of G_A dominates atleast one of the color classes of c_i and also appears in singleton of χ -partition. Hence G_A is χ_{fd} -excellent.

Proposition 3.7.

1. If an anti fuzzy graph G_A has no edges with m vertices then fuzzy dominator chromatic number is m .
2. For any bipartite anti fuzzy graph with atleast one edge, the fuzzy dominator chromatic number is at least $|D| + 1$. Where $|D|$ is the number of vertices in a minimal dominating set of an anti fuzzy graph G_A .

Example 3.8. Consider the following Figure 3, for characterizing chromatic number of an anti fuzzy graph.

Figure 3. An Anti Fuzzy Graph G_A

In Figure 3, $C_1 = (\{v_2\}, \{v_1, v_3, v_5\}, \{v_4\})$. The vertices v_1, v_3, v_5 do not appear in singleton of χ -partition. Hence G_A is not χ_{fd} -excellent.

Theorem 3.9. For any connected anti fuzzy graph, $\chi_{fd}(G_A) \geq |D| + 1$.

Proof. Suppose $N(V(D))=1$ and any one of the color in color class ' c_i ' is assigned except the color assigned by the dominating vertices in D . All the vertices in D is assigned with individual color from color class c_i . Therefore, the color class should contain $|D|$ colors. Let $v_n \in V \setminus D$ where v_n is adjacent with atleast one vertex in D . So, the vertex v_n is assigned with $c_{|D|+1}^{th}$ color and this vertex dominate any one of color class of G_A .

Therefore, $\chi_{fd}(G_A) = |D| + 1$. — — — $\rightarrow$ (1)

If $N(V(D)) \geq 1$ then all the vertices in D is assigned with an individual color from color class c_i . Therefore, the color class should contain $|D|$ colors. Assume that the vertices $v_n, v_{n+1} \in V \setminus D$. To construct a proper dominator coloring, these vertices are colored with $c_{|D|+1}^{th}$ and $c_{|D|+2}^{th}$ color. Since v_n adjacent to v_{n+1} .

Therefore, $\chi_{fd}(G_A) = |D| + 2$. — — — $\rightarrow$ (2)

Similarly repeat this process until all the vertices in $V \setminus D$ is assigned with some colors.

Hence, from (1) and (2), $\chi_{fd}(G_A) \geq |D| + 1$. $\square$

Theorem 3.10. If G_A is an anti fuzzy path with $m \geq 5$ vertices then $\chi_{fd}(G_A) = |D| + 2$.

Proof. Let v_i be the internal vertex in an anti fuzzy path G_A . It dominates v_{i-1}, v_{i+1} . Therefore, v_i may be assigned with color c_1 in c . The vertices v_{i-1}, v_{i+1} may be assigned with color $c_{|D|+1}^{th}$ in c . If $v_{i+2} \notin D$ and v_{i+1} adjacent with v_{i+2} then v_{i+2} is assigned with color $c_{|D|+2}$.

Suppose $v_{i+2} \in D$ then v_{i+2} is assigned with color $c_2 \leq c_{|D|}$. Therefore, the vertices in D are assigned with individual color and dominate any color classes of c . The vertices in $V \setminus D$ is assigned with alternate color of $c_{|D|+1}, c_{|D|+2}$. Since any pair of vertices in $V \setminus D$ which are adjacent to each other.

Hence $\chi_{fd}(G_A) = |D| + 2$. $\square$

Proposition 3.11.

1. If G_A is an anti fuzzy path with $m < 5$ vertices then $\chi_{fd}(G_A) = |D| + \lceil \frac{m}{3} \rceil$.
2. For a uninodeal anti fuzzy graph G_A , $\chi_{fd}(G_A) = \lceil \frac{2m}{3} \rceil$.
3. If G_A is e -nodal anti fuzzy graph with effective edges only then $\chi_{fd}(G_A) \geq |D| + 1$.

Remark 3.12. Consider that the $|D|$ is the number of vertices in a minimal dominating set of an anti fuzzy graph G_A .

1. If an anti fuzzy path contains two vertices then $|D| = 1$, $\lceil \frac{2}{3} \rceil = 1$. Therefore $\chi_{fd}(G_A) = 1 + 1 = 2$.
2. If an anti fuzzy path contains 3 vertices (say u_1, u_2, u_3) then $|D| = 1$ and $|V \setminus D| = 2$. That is, the internal vertex $u_2 \in D$ and assigned with color c_1 . The vertices in $V \setminus D$ are non adjacent to each other and a strong neighbor to a vertex u_2 in D . So the vertices u_1 and u_3 are assigned with same color c_2 . $|D| = 1$, $\lceil \frac{3}{3} \rceil = 1$. Therefore, $\chi_{fd}(G_A) = 1 + 1 = 2$.

Theorem 3.13. If G_A is a anti fuzzy tree with $m \geq 3$ vertices then $\chi_{fd}(G_A) = |S| + 1$, where S is the set of all support vertices in G_A .

Proof. Consider that G_A is an anti fuzzy tree with atleast 3 vertices. Then G_A has atleast one support vertex (say u). The vertex u is colored with color 1 from $|S|$ color classes which also dominate other vertices in G_A . The vertices which are adjacent to u but not a leaf of G_A is assigned with color $(|S|-1)^{th}$. Similarly, repeat this procedure until all the support vertices are assigned with some color from $|S|$ color classes. Finally, all the pendent vertices are assigned with $(|S|+1)^{th}$ color. Hence $\chi_{fd}(G_A) = |S| + 1$. $\square$

Theorem 3.14. For any connected irregular anti fuzzy graph G_A ,

1. $\chi(G_A) \geq \Delta_{I_c}$.
2. $\chi_{fd}(G_A) \geq \Delta_{I_c} + 1$.

where I_c is the set of all incident vertices of a vertex v in G_A and Δ_{I_c} is the maximum number of incident vertices of a vertex in G_A .

Proof. Consider that G_A is a connected irregular anti fuzzy graph. That is, any one of the vertices in G_A has number of incident vertices as Δ_{I_c} . For coloring an anti fuzzy graph G_A , a vertex is arbitrarily chosen (say u) which have Δ_{I_c} neighbors and assign with color 1. All the vertices in G_A , except u have atmost Δ_{I_c} neighbors. Consider a vertex $v \in V(G_A) \setminus \{u\}$ which have Δ_{I_c} neighbors. This vertex is colored with $(\Delta_{I_c} - 1)^{th}$ color. Since u is adjacent with v . Repeat this process until all the vertices are assigned with a color. Finally, the anti fuzzy graph G_A is colored using Δ_{I_c} colors. Hence $\chi(G_A) \geq \Delta_{I_c}$.

Similarly, choose another vertex $w \in V(G_A) \setminus \{u, v\}$ which have ΔI_c neighbors and colored the vertex w by $(\Delta I_c - 1)^{th}$ color. Since w is adjacent with u but not with v . Repeat this procedure until all vertices (having ΔI_c neighbors) are colored. Finally, get some vertices which does not have the ΔI_c neighbors. Such vertex can colored by $(\Delta I_c + 1)^{th}$ color. Hence the vertices which are having ΔI_c neighbors should be a member in dominating set of G_A . The color class which contains these vertices should form a dominator coloring of G_A . Therefore, $\chi_{fd}(G_A) \geq \Delta I_c + 1$. $\square$

Theorem 3.15. For any connected k -regular anti fuzzy graph G_A , $\chi_{fd}(G_A) > \Delta I_c + 1$.

Proposition 3.16.

1. For any connected anti fuzzy graph G_A , $\chi(G_A) \leq \chi_{fd}(G_A)$
2. Complete anti fuzzy graph is χ_{fd} -excellent.
3. Anti complement of complete anti fuzzy graph is also χ_{fd} -excellent.
4. Cyclic anti fuzzy graph G_A is χ_{fd} -excellent.
5. If anti fuzzy graph G_A is star then G_A is χ_{fd} -excellent.
6. Any connected k -regular anti fuzzy graph G_A is χ_{fd} -excellent.

4. Dominator coloring on Anti Fuzzy Graph with weak edges

In this section, the dominator chromatic number is derived for an anti fuzzy graph which has atleast one weakest edges. The color class which contains only the strong edges is assigned with membership fuzzy value as 1. If the color class has atleast one weakest edges then the membership value of color class is defined as follows.

Definition 4.1. Let u, v be two vertices in G_A and u, v are adjacent with one weakest edge (e) then the fuzzy color values of u is defined as $1 - [\mu(u, v) - \vee(\sigma(u), \sigma(v))]$ and the color class value is minimum of fuzzy color values of color class c_i .

Example 4.2. In Figure 4, G_A is an anti fuzzy graph with one weak edge (v_1, v_5).

The vertices v_5 and v_6 dominate all other vertices in G_A and maximum fuzzy cardinality of a dominating set with minimum number of vertices is obtained with these vertices. Therefore, it forms a minimal dominating set. So, these vertices are assigned with individual colors such as 1 and 2. The corresponding color class is c_1, c_2 and the membership value is 1.

Next, colors are assigned to the remaining vertices with color class 3, 4, ... Randomly choose a vertex v_1 , which is adjacent with v_5 but the edge (v_1, v_5) is weak edge in G_A .

Figure 4. An Anti Fuzzy Graph G_A

So the vertex v_1 is assigned with color 3 whose fuzzy color membership value is $1 - [0.6 - (0.1 \vee 0.4)] = 1 - (0.6 - 0.4) = 1 - (0.6 - 0.4) = 1 - 0.2 = 0.8$.

Then the vertices v_2 and v_4 are adjacent with v_1 as strong neighbors. Such vertices are assigned with color 4. The vertex v_3 is adjacent with v_2 and v_4 as a strong neighbor. Color 3 is assigned with fuzzy color membership value 1. $c = \{c_1, c_2, c_3, c_4\} = \{\{v_5\}^1, \{v_6\}^1, \{v_1, v_3\}^{0.8}, \{v_2, v_4\}^1\}$

Therefore, $\chi_{fd}(G_A) = |c| = 1 + 1 + 0.8 + 1 = 3.8$

Proposition 4.3.

1. If G_A is e -nodal anti fuzzy graph with weak edges then $\chi_{fd}(G_A) \geq |D| + \frac{m}{2}$.
2. In e -nodal (v -nodal) anti fuzzy graph with weak edges, the vertex incident with weak edges is assigned the fuzzy color value is greater than 1.

5. Dominator Coloring on Anti Cartesian Product of Anti Fuzzy Graphs

In this section, the dominator coloring is applied on anti cartesian product of anti fuzzy graphs and obtained the bounds of fuzzy dominator chromatic number on anti cartesian product of anti fuzzy graphs. Anti cartesian product of anti fuzzy graphs are considered as strong anti fuzzy graph and the fuzzy value of the color class is 1. If $G_{A_1} \times G_{A_2}$ has weak edges then the fuzzy value is calculated by $1 - [\mu(u, v) - \vee(\sigma(u), \sigma(v))]$. In this section, all G_{A_1} and G_{A_2} are considered as strong anti fuzzy graph.

Definition 5.1. [11] Let G_A be an anti cartesian product of anti fuzzy graphs G_{A_1} and G_{A_2} where $G_{A_1} = (\sigma_1, \mu_1)$ and $G_{A_2} = (\sigma_2, \mu_2)$ then $G_A = (\sigma_1 \times \sigma_2, \mu_1 \times \mu_2)$ is an anti fuzzy graph.

Theorem 5.2. Let G_{A_1} and G_{A_2} be the anti fuzzy paths on m_1 and m_2 vertices. If the anti Cartesian product $G_{A_1} \times G_{A_2}$ of G_{A_1} and G_{A_2} is an anti fuzzy graphs on $m_1 m_2$ vertices then

$\chi_{fd}(G_{A_1} \times G_{A_2}) \geq |D| + 2$, where $|D|$ is the number of vertices in a minimal dominating set of G_A .

Theorem 5.3. Let $G_A = G_{A_1} \times G_{A_2}$ be an anti cartesian product of anti fuzzy graphs G_{A_1} and G_{A_2} then $\chi_{fd}(G_{A_1} \times G_{A_2}) \geq \max \{ \chi_{fd}(G_{A_1}), \chi_{fd}(G_{A_2}) \}$.

Proof. By the definition of anti cartesian product of anti fuzzy graphs, any two vertices are adjacent in G_{A_1} or in G_{A_2} if and only if the corresponding pair of vertices are adjacent in $G_{A_1} \times G_{A_2}$. That is, by the definition of anti cartesian product of anti fuzzy graphs, more adjacency exists in $G_{A_1} \times G_{A_2}$. Assigning the color to the vertices depend on the adjacency in $G_{A_1} \times G_{A_2}$. Therefore, the number of colors required to the vertices of $G_{A_1} \times G_{A_2} \geq \max \{ \chi_{fd}(G_{A_1}), \chi_{fd}(G_{A_2}) \}$. $\square$

Theorem 5.4. Let G_{A_1} and G_{A_2} be the anti fuzzy cycles on m_1 and m_2 vertices. If the anti cartesian product of anti fuzzy cycle $G_A = (G_{A_1} \times G_{A_2})$ is an anti fuzzy graph on $m_1 m_2$ vertices then $\chi_{fd}(G_{A_1} \times G_{A_2}) \geq |D| + 2$, where $|D|$ is the number of vertices in a minimal dominating set of G_A .

Proof. Similar proof as in [13] for fuzzy graph $\square$

Theorem 5.5. If $G_{A_1} \times G_{A_2}$ is an anti cartesian product of anti fuzzy cycles with $m_1 m_2$ vertices then $\chi(G_{A_1} \times G_{A_2}) \leq \chi_{fd}(G_{A_1} \times G_{A_2})$. Where $|V(G_{A_1})| = m_1$, $|V(G_{A_2})| = m_2$

Proof. Every vertex in $G_{A_1} \times G_{A_2}$ are assigned with some colors in color class $\{c_1, c_2, \dots, c_m\}$ where $m = m_1 \times m_2$. By the definition of proper coloring, adjacent pair of vertices are assigned with distinct colors. Therefore, by the definition of anti cartesian product of anti fuzzy cycles the vertices have maximum 4 strong neighbors and minimum 3 neighbors. Hence $\chi(G_{A_1} \times G_{A_2}) = 3$. $\longrightarrow (1)$

In $G_{A_1} \times G_{A_2}$, there exists a member in a minimal dominating set is $|D| \geq 3$. Neighbors of the vertex in $V(D)$, each member may be assigned with same color in some color classes (say $c_{|D|+1}^{th}$ color) but a vertex (u_i, v_j) is a strong neighbor to atleast one dominating vertex and a neighbor to a vertex in $V \setminus D$. In such case, the vertex (u_i, v_j) and (u_i, v_{j+1}) is assigned with distinct colors.

Hence $\chi_{fd}(G_{A_1} \times G_{A_2}) \geq |D| + 2 \longrightarrow (2)$

From (1) and (2), $\chi(G_{A_1} \times G_{A_2}) \leq \chi_{fd}(G_{A_1} \times G_{A_2})$. $\square$

Theorem 5.6. Let $G_{A_1} \times G_{A_2}$ be an anti cartesian product of complete anti fuzzy graphs on $m = m_1 \times m_2$ vertices.

1. For $m_1 = 2$ and $m_2 \geq m_1$ then $\chi_{fd}(G_{A_1} \times G_{A_2}) \geq (m_1 \vee m_2) + 1$.

2. For $m_1, m_2 > 2$, then $\chi_{fd}(G_{A_1} \times G_{A_2}) \geq (m_1 \vee m_2) + 2$.

Example 5.7. consider a complete anti fuzzy graph K_2 . Apply the anti cartesian product on anti fuzzy graphs K_2 and itself. The resulting anti fuzzy graph is shown in Figure 5.

Here in Figure 5, $D = \{(u_1, v_2), (u_2, v_2)\}$, $|D| = 2$,
 $\gamma = .3 + .3 = 0.6$

Let the family of color classes be $c = \{c_1, c_2, c_3, c_4\}$

Figure 5. An Anti Fuzzy Graph $G_A = K_2 \times K_2$

$$\begin{aligned} c_1\{(u_i, v_j)\} &= \begin{cases} 0.3, & \text{if } i = 1, j = 2 \\ 0, & \text{otherwise} \end{cases} \\ c_2\{(u_i, v_j)\} &= \begin{cases} 0.3, & \text{if } i = 2, j = 2 \\ 0, & \text{otherwise} \end{cases} \\ c_3\{(u_i, v_j)\} &= \begin{cases} 0.2, & \text{if } i = 2, j = 1 \\ 0, & \text{otherwise} \end{cases} \\ c_4\{(u_i, v_j)\} &= \begin{cases} 0.2, & \text{if } i = 1, j = 1 \\ 0, & \text{otherwise} \end{cases} \end{aligned}$$

Hence all the vertices in $V(G_{A_1} \times G_{A_2})$ belong to some distinct color classes. It satisfies the property of fuzzy dominator coloring. The χ_{fd} -partition is $C_1 = (\{v_1\}, \{v_2\}, \{v_3\}, \{v_4\})$. Hence G_A is χ_{fd} -excellent.

Results

1. If G_{A_1} and G_{A_2} are χ_{fd} -excellent then $G_{A_1} \times G_{A_2}$ need not be χ_{fd} -excellent.

6. Conclusion

The dominator coloring is applied on some types of anti fuzzy graph and obtained the bounds of fuzzy dominator chromatic number on anti fuzzy graph. Fuzzy chromatic value is derived for anti fuzzy graph with effective and weak edges. Also, the fuzzy dominator chromatic number is applied on the anti cartesian product of anti fuzzy graph and derived some theorems on them.

References

- [1] S. Arumugam, Jay bagga and K.M.Raja chandrasekar, On dominator colorings in graphs, *Proc. Indian Acad. Sci., (Math. Sci.)* 122(4)(November 2012), 560–571.
- [2] C. Berge, *Theorey of graphs*, dover Publications, Inc.Mineola, New York, 2001.

- [3] C. Eslachi, B. N. Onagh, Vertex strength of fuzzy graphs, *International journal of Mathematics and mathematical sciences* (2006).
- [4] R. Gera, On Dominator coloring in Graphs, *Graph Theory Notes of New York LII*, (2007), 25–30.
- [5] R. Gera, On dominator colorings in bipartite graphs, *(ITNG'07)IEEE*, (2007), 1–6.
- [6] R. Gera, S. Horton and C. Ramussen, "Dominator colorings and safe clique partitions", *Congressus Numerantium*, 81(2006), 19–20.
- [7] R. jahir Hussian and K.S. Kanzol Fathima, On Fuzzy Dominator coloring in Fuzzy Graphs, *Applied Mathematical Sciences*, 9(23) (2015), 1131–1137.
- [8] Muhammad Akram, Anti Fuzzy Structures on Graphs, *Middle East journal of Scientific Research*, 11(12) (2012), 1641–1648.
- [9] S. Munoz, T. Ortuno, J. Ramirez, J. Yanez, Coloring fuzzy graphs, *Omega* 33(3) (2005), 211–221.
- [10] R. Muthuraj and A. Sasireka, On Anti Fuzzy Graph, *Advances in Fuzzy Mathematics*, 12(5) (2017), 1123–1135.
- [11] R. Muthuraj and A. Sasireka, Some Characterization on Operations of Anti Fuzzy Graphs", *International Conference on Mathematical Impacts in Science And Technology, (MIST -17), IJRASET*, (November 2017), 109–117.
- [12] R. Muthuraj and A. Sasireka, Some Types of Nodal and Edge Regular Anti Fuzzy Graph, *International Journal of Fuzzy Mathematical Archive*, 14(2) (2017), 365–378.
- [13] R. Muthuraj and A. Sasireka, Fuzzy dominator coloring and fuzzy chromatic number on Cartesian product of simple fuzzy graph, *Advances in theoretical and applied Mathematics*, 11(3) (2016), 245–260.
- [14] R. Muthuraj and A. Sasireka, Domination on Anti Fuzzy Graph, *International Journal of Mathematical Archive*, 9(5) (2018), 82–92.
- [15] R. Muthuraj and A. Sasireka, Connected Domination on Anti Fuzzy Graph, *Journal of Applied Science and Computations*, 5(8) (2018), 18–28.
- [16] R. Muthuraj and A. Sasireka, Total Domination on Anti Fuzzy Graph, *New Trends in Mathematical Sciences*, 6(4) (2018), 28–39.
- [17] A. Prasanna, M. A. Rifayathali and S. Ismail Mohideen, Anti-fuzzy graph coloring, *international journal of science and advance Research in Technology*, 4(4) (2018), 2598–2603.
- [18] T. Ramachandran, A. Naseer Ahmed, Dominator Coloring of Middle Graph, *IOSR Journal of Mathematics (IOSR-JM)*, 10(6) (IV) (2014), 38–40.
- [19] M.A. Rifayathali, A. Prasanna and S. Ismail Mohideen, Chromatic Excellence in Anti-fuzzy graphs, *journal of applied science and computations*, 5(7) (july/2018), 305–316.
- [20] R. Seethalakshmi, and R.B. Gnanajothi, Operations on Antifuzzy graphs, *Mathematical Sciences International Research Journal*, 5(2) (2016), 210–214.

*****
 ISSN(P):2319 – 3786
 Malaya Journal of Matematik
 ISSN(O):2321 – 5666
 *****