



Fixed point theorems for orthogonal F -Suzuki contraction mappings on O -complete metric space with an applications

Gunaseelan Mani¹, Arul Joseph Gnanaprakasam²,
Lakshmi Narayan Mishra^{3*} and Vishnu Narayan Mishra⁴

Abstract

In this paper, we introduce the concepts of generalized orthogonal F -contraction and orthogonal F -Suzuki contraction mappings and prove some fixed point theorems for a self mapping in orthogonal metric space. The proved results generalize and extend some of the well known results in the literature. An example to support our result is presented. As applications of the main results, we apply our main results to show the existence of a unique solution of the first-order ordinary differential equation.

Keywords

Orthogonal set, orthogonal metric space, orthogonal continuous, orthogonal preserving, generalized orthogonal F -contraction, orthogonal F -Suzuki contraction, fixed point.

AMS Subject Classification

47H10, 54H25.

¹Department of Mathematics, Sri Sankara Arts and Science College(Autonomous), Affiliated to Madras University, Enathur, Kanchipuram, Tamil Nadu, India 631 561.

²Department of Mathematics, College of Engineering and Technology, Faculty of Engineering and Technology, SRM Institute of Science and Technology, SRM Nagar, Kattankulathur 603203, Kanchipuram, Chennai, Tamil Nadu, India.

³Department of Mathematics, School of Advanced Sciences, Vellore Institute of Technology (VIT) University, Vellore 632 014, Tamil Nadu, India.

⁴Department of Mathematics, Indira Gandhi National Tribal University, Laipur, Amarkantak, Anuppur, Madhya Pradesh, India 484887.

*Corresponding author: ¹ mathsguna@yahoo.com; ² aruljoseph.alex@gmail.com; ³ lakshminarayanmishra04@gmail.com

⁴ vishnunarayanmishra@gmail.com

Article History: Received 11 November 2020; Accepted 09 January 2021

©2021 MJM.

Contents

1	Introduction	369
2	Preliminaries	370
3	Main Results	371
4	Example	374
5	Application to ordinary differential equations	375
6	Conclusion	376
	References	376

1. Introduction

The concept of an orthogonal set has many applications in several branches in mathematics and it has many types of the orthogonality. Gordji, Ramezani, De La Sen and Cho [1] introduced the new concept of an orthogonality in metric spaces

and proved the fixed point result for contraction mappings in metric spaces endowed with the new orthogonality. Furthermore, they gave the application of this results for claming the existence and uniqueness of solution of the first-ordinary differential equation while the Banach contraction mapping can not be applied in this problem. Eshaghi Gordji and Habibi [2] proved fixed point theory in generalized orthogonal metric space. Sawangsup, Sintunavarat and Cho [3] introduced the new concept of an orthogonal F -contraction mappings and proved the fixed point theorems on orthogonal-complete metric space. The orthogonal contractive type mappings has been studied by many authors and important results have been obtained by [4], [5], [6], [7], [8]. Some interesting concepts and application see [11–17]. In this paper, we introduced the new concepts generalized orthogonal F -contraction and orthogonal F -Suzuki contraction mappings and prove the fixed point theorems on orthogonal complete metric space.

2. Preliminaries

Throughout this paper, we denote by W , \mathbb{R}^+ , \mathbb{N} and \mathbb{N}_0 the nonempty set, the set of positive real numbers, the set of positive integers and the set of nonnegative integers, respectively. Firstly, we recall the concept of a control function which is introduced by Wardowski [9]. Let \mathfrak{F} denote the family of all functions $F : \mathbb{R}^+ \rightarrow \mathbb{R}$ satisfying the following properties:

(F_1) F_1 is strictly increasing;

(F_2) for each sequence $\{\alpha_n\}$ of positive numbers, we have

$$\lim_{n \rightarrow \infty} \alpha_n = 0 \iff \lim_{n \rightarrow \infty} F(\alpha_n) = -\infty;$$

(F_3) there exists $k \in (0, 1)$ such that $\lim_{\alpha \rightarrow 0^+} \alpha^k F(\alpha) = 0$.

Piri and Kumam [10] introduced a large class of functions by replacing the condition (F_3) in the definition of F -contractions with the following

(F'_3) F is continuous on $(0, \infty)$.

Let the set $\mathfrak{D} = \{F : \mathbb{R}_+ \rightarrow \mathbb{R} \mid F \text{ satisfies the conditions } (F_1), (F_2) \text{ and } (F'_3)\}$. He also proved the following existence and uniqueness of a fixed point for F -contraction mappings:

Theorem 2.1. [10] Let (W, d) be a complete metric space and a mapping $Y : W \rightarrow W$. Suppose that there exist $F \in \mathfrak{D}$ and $\tau > 0$ such that, for all $u, v \in W$

$$d(Yu, Yv) > 0 \Rightarrow \tau + F(d(Yu, Yv)) \leq F(d(u, v)).$$

Then Y has a unique fixed point $t \in W$ and for every $u \in W$, the sequence $\{Y^n u\}$ converges to t .

Next, he introduce notion of F -Suzuki contraction and proved existence and uniqueness of a fixed point theorem.

Definition 2.2. [10] Let (W, d) be a metric space. A mapping $Y : W \rightarrow W$ is said to be an F -Suzuki contraction if there exists $\tau > 0$ such that for all $u, v \in W$ with $Yu \neq Yv$,

$$\frac{1}{2}d(u, Yu) < d(u, v) \Rightarrow \tau + F(d(Yu, Yv)) \leq F(d(u, v)),$$

where $F \in \mathfrak{D}$.

Theorem 2.3. [10] Let (W, d) be a complete metric space and $Y : W \rightarrow W$ be an F -suzuki contraction. Then Y has a unique fixed point $u^* \in W$ and for every $u_0 \in W$, the sequence $\{Y^n u_0\}_{n=1}^\infty$ converges to u^* .

We give some examples of functions belonging to \mathfrak{D} as follows:

Example 2.4. Let functions $F_1, F_2, F_3, F_4 : \mathbb{R}^+ \rightarrow \mathbb{R}$ be defined by:

$$(1) F_1(\beta) = \frac{-1}{\beta} \text{ for } \beta > 0;$$

$$(2) F_2(\beta) = \frac{-1}{\beta} + \beta \text{ for all } \beta > 0;$$

$$(3) F_3(\beta) = \frac{1}{1-e^\beta} \text{ for all } \beta > 0;$$

$$(4) F_4(\beta) = \frac{1}{e^\beta - e^{-\beta}} \text{ for all } \beta > 0.$$

Then $F_1, F_2, F_3, F_4 \in \mathfrak{D}$.

On the other hand, Gordji, Ramezani, De La Sen and Cho [1] introduced the concept of an orthogonal set (or O -set), some examples and some properties of the orthogonal sets as follows:

Definition 2.5. [1] Let $W \neq \emptyset$ and $\perp \subseteq W \times W$ be a binary relation. If \perp satisfies the following condition:

$$\exists w_0 \in W : (\forall w \in W, w \perp w_0) \quad \text{or} \quad (\forall w \in W, w_0 \perp w),$$

then it is called an orthogonal set (briefly O -set). We denote this O -set by (W, \perp) .

Example 2.6. [1] Let W be the set all people in the world. Define the binary relation \perp on W by $u \perp w$ if u can give blood to w . According to the Table 1, if w_0 is a person such that his(her) blood type is O -, then we have $w_0 \perp w$ for all $w \in W$. This means that (W, \perp) is an O -set. In this O -set, w_0 (in Definition 2.5) is not unique. Note that, in this example, w_0 may be a person with blood type $AB+$. In this case, we have $w \perp w_0$ for all $w \in W$.

Table 1

Type	You can give blood to	You can receive blood from
A+	A+AB+	A+A-O+O-
O+	O-A+B+AB+	O+O-
B+	B+AB+	B+B-O+O-
AB+	AB+	Everyone
A-	A+A-AB+AB-	A-O-
O-	Everyone	O-
B-	B+B-AB+AB-	B-O-
AB-	AB+AB-	AB-B-O-A-

Example 2.7. [1] Let $W = \mathbb{Z}$. Define the binary relation \perp on W by $m \perp n$ if there exists $k \in \mathbb{Z}$ such that $m = kn$. It is easy to see that $0 \perp n$ for all $n \in \mathbb{Z}$. Hence, (W, \perp) is an O -set.

Example 2.8. [1] Let (W, d) be a metric space and $Y : W \rightarrow W$ be a Picard operator, that is, Y has a unique fixed point $w^* \in W$ and $\lim_{n \rightarrow \infty} Y^n(u) = w^*$ for all $u \in W$. We define the binary relation \perp on W by $u \perp w$ if

$$\lim_{n \rightarrow \infty} d(w, Y^n(u)) = 0.$$

Then, (W, \perp) is an O -set.

Example 2.9. [1] Let $W = [0, \infty)$ and define $w \perp u$ if $wu \in \{w, u\}$. Then, by setting $w_0 = 0$ or $w_0 = 1$, (W, \perp) is an O -set.



Now, we give the concepts of an O -sequence, a \perp -continuous mapping, an O -complete orthogonal metric space, a \perp -preserving mapping and a weakly \perp -preserving mapping.

Definition 2.10. [1] Let (W, \perp) be an O -set. A sequence $\{w_n\}$ is called an orthogonal sequence (briefly, O -sequence) if

$$(\forall n \in \mathbb{N}, w_n \perp w_{n+1}) \quad \text{or} \quad (\forall n \in \mathbb{N}, w_{n+1} \perp w_n).$$

Definition 2.11. [1] The triplet (W, \perp, d) is called an orthogonal metric space if (W, \perp) is an O -set and (W, d) is a metric space.

Definition 2.12. [1] Let (W, \perp, d) be an orthogonal metric space. Then, a mapping $Y : W \rightarrow W$ is said to be orthogonally continuous (or \perp -continuous) in $w \in W$ if for each O -sequence $\{w_n\}$ in W with $w_n \rightarrow w$ as $n \rightarrow \infty$, we have $Y(w_n) \rightarrow Y(w)$ as $n \rightarrow \infty$. Also, Y is said to be \perp -continuous on W if Y is \perp -continuous in each $w \in W$.

Remark 2.13. [1] Every continuous mapping is \perp -continuous and the converse is not true.

Definition 2.14. [1] Let (W, \perp, d) be an orthogonal metric space. Then, W is said to be orthogonally complete (briefly, O -complete) if every Cauchy O -sequence is convergent.

Remark 2.15. [1] Every complete metric space is O -complete and the converse is not true.

Definition 2.16. [1] Let (W, \perp) be an O -set. A mapping $Y : W \rightarrow W$ is said to be \perp -preserving if $Yw \perp Yu$ whenever $w \perp u$. Also $Y : W \rightarrow W$ is said to be weakly \perp -preserving if $Y(w) \perp Y(u)$ or $Y(u) \perp Y(w)$ whenever $w \perp u$.

In this paper, we modify the concepts of F -contraction and F -Suzuki contraction mappings to orthogonal sets and prove some fixed point theorems for F -contraction and F -Suzuki contraction mappings in orthogonality complete metric spaces. Also, we give some example to illustrate our results. Furthermore, we apply our results to show the existence and uniqueness of a solution of the first-order ordinary differential equation.

3. Main Results

In this section, inspired by the notions of an F -contraction mapping, F -Suzuki contraction and an orthogonal set, we introduce a new F -contraction mapping, new F -Suzuki contraction and prove some fixed point theorems for these contraction mappings in an orthogonal metric space.

Definition 3.1. Let (W, \perp, d) be an orthogonal metric space. A map $Y : W \rightarrow W$ is said to be a generalized orthogonal F -contraction mapping (briefly, generalized Y_\perp -contraction) on (W, \perp, d) if there are $F \in \mathcal{D}$ and $\tau > 0$ such that the following condition holds:

$$\forall w, u \in W \text{ with } w \perp u [d(Yw, Yu) > 0 \\ \Rightarrow \tau + F(d(Yw, Yu)) \leq F(d(w, u))].$$

Definition 3.2. Let (W, \perp, d) be an orthogonal metric space. A map $Y : W \rightarrow W$ is said to be a orthogonal F -Suzuki contraction mapping (briefly, Y_\perp -Suzuki contraction) on (W, \perp, d) if there are $F \in \mathcal{D}$ and $\tau > 0$ such that the following condition holds:

$$\forall w, u \in W \text{ with } w \perp u [d(Yw, Yu) > 0, \\ \frac{1}{2}d(w, Yw) < d(w, u) \\ \Rightarrow \tau + F(d(Yw, Yu)) \leq F(d(w, u))].$$

Now, we give the first fixed point theorem for generalized orthogonal F -contraction mapping in an O -complete orthogonal metric space (W, \perp, d) .

Theorem 3.3. Let (W, \perp, d) be an O -complete orthogonal metric space with an orthogonal element w_0 and a mapping $Y : W \rightarrow W$. Suppose that there exist $F \in \mathcal{D}$ and $\tau > 0$ such that the following conditions hold:

- (i) Y is \perp -preserving;
- (ii) Y is generalized Y_\perp -contraction mapping;
- (iii) Y is \perp -continuous.

Then Y has a unique fixed point $t \in W$ and for every $w \in W$, the sequence $\{Y^n w\}$ converges to t .

Proof. Since (W, \perp) is an O -set,

$$\exists w_0 \in W : (\forall w \in W, w \perp w_0) \quad \text{or} \quad (\forall w \in W, w_0 \perp w).$$

It follows that $w_0 \perp Yw_0$ or $Yw_0 \perp w_0$. Let

$$w_1 := Yw_0, w_2 := Yw_1 = Y^2w_0 \dots \\ \dots, w_{n+1} := Yw_n = Y^{n+1}w_0$$

for all $n \in \mathbb{N} \cup \{0\}$. If $w_n = w_{n+1}$ for any $n \in \mathbb{N} \cup \{0\}$, then it is clear that w_n is a fixed point of Y . Assume that $w_n \neq w_{n+1}$ for all $n \in \mathbb{N} \cup \{0\}$. Thus, we have $d(Yw_n, Yw_{n+1}) > 0$ for all $n \in \mathbb{N} \cup \{0\}$. Since Y is \perp -preserving, we have

$$w_n \perp w_{n+1} \quad \text{or} \quad w_{n+1} \perp w_n \tag{3.1}$$

for all $n \in \mathbb{N} \cup \{0\}$. This implies that $\{w_n\}$ is an O -sequence. Since Y is generalized Y_\perp -contraction mapping, we have

$$F(d(w_n, w_{n+1})) = F(d(Yw_{n-1}, Yw_n)) \\ \leq F(d(w_{n-1}, w_n)) - \tau, \quad \forall n \in \mathbb{N}.$$

Repeating this process, we get

$$F(d(Yw_{n-1}, Yw_n)) \leq F(d(w_{n-1}, w_n)) - \tau \\ = F(d(Yw_{n-2}, Yw_{n-1})) - \tau \\ \leq F(d(w_{n-2}, w_{n-1})) - 2\tau \\ = F(d(Yw_{n-3}, Yw_{n-2})) - 2\tau \\ \leq F(d(w_{n-3}, w_{n-2})) - 3\tau \\ \vdots \\ \leq F(d(w_0, w_1)) - n\tau. \tag{3.2}$$



From (3.2), we obtain $\lim_{n \rightarrow \infty} F(d(w_n, Yw_n)) = -\infty$.

By the property (F_2) , we have

$$\lim_{n \rightarrow \infty} d(w_n, Yw_n) = 0. \quad (3.3)$$

Now, we claim that $\{w_n\}_{n=1}^\infty$ is a Cauchy sequence. Arguing by contradiction, we assume that there exist $\varepsilon > 0$ and sequence $\{i(n)\}_{n=1}^\infty$ and $\{j(n)\}_{n=1}^\infty$ of natural numbers such that

$$\begin{aligned} i(n) > j(n) > n, \quad d(w_{i(n)}, w_{j(n)}) \geq \varepsilon, \\ d(w_{i(n)-1}, w_{j(n)}) < \varepsilon, \quad \forall n \in \mathbb{N}. \end{aligned} \quad (3.4)$$

So, we have

$$\begin{aligned} \varepsilon \leq d(w_{i(n)}, w_{j(n)}) &\leq d(w_{i(n)}, w_{i(n)-1}) + d(w_{i(n)-1}, w_{j(n)}) \\ &< d(w_{i(n)}, w_{i(n)-1}) + \varepsilon \\ &= d(w_{i(n)-1}, Yw_{i(n)-1}) + \varepsilon. \end{aligned}$$

It follows from (3.3) and the above inequality that

$$\lim_{n \rightarrow \infty} d(w_{i(n)}, w_{j(n)}) = \varepsilon. \quad (3.5)$$

On the other hand, from (3.3) there exists $N \in \mathbb{N}$, such that

$$d(w_{i(n)}, Yw_{i(n)}) < \frac{\varepsilon}{4} \quad \text{and} \quad d(w_{j(n)}, Yw_{j(n)}) < \frac{\varepsilon}{4}, \quad (3.6)$$

$\forall n \in \mathbb{N}$. Next, we claim that

$$d(Yw_{i(n)}, Yw_{j(n)}) = d(w_{i(n)+1}, w_{j(n)+1}) > 0, \quad \forall n \in \mathbb{N}. \quad (3.7)$$

Arguing by contradiction, there exists $m \geq \mathbb{N}$ such that

$$d(w_{i(m)+1}, w_{j(m)+1}) = 0. \quad (3.8)$$

It follows from (3.4), (3.6) and (3.8) that

$$\begin{aligned} \varepsilon \leq d(w_{i(m)}, w_{j(m)}) &\leq d(w_{i(m)}, w_{i(m)+1}) \\ &\quad + d(w_{i(m)+1}, w_{j(m)}) \\ &\leq d(w_{i(m)}, w_{i(m)+1}) \\ &\quad + d(w_{i(m)+1}, w_{j(m)+1}) \\ &\quad + d(w_{j(m)+1}, w_{j(m)}) \\ &= d(w_{i(m)}, Yw_{i(m)}) \\ &\quad + d(w_{i(m)+1}, w_{j(m)+1}) \\ &\quad + d(w_{j(m)}, Yw_{j(m)}) \\ &< \frac{\varepsilon}{4} + 0 + \frac{\varepsilon}{4} = \frac{\varepsilon}{2}. \end{aligned}$$

This contradiction establishes the relation (3.7). Since Y is \perp -preserving, we have

$$w_{i(n)} \perp w_{j(n)} \quad \text{or} \quad w_{j(n)} \perp w_{i(n)}.$$

Since Y is generalized orthogonal F -contraction mapping, we have

$$\tau + F(d(Yw_{i(n)}, Yw_{j(n)})) \leq F(d(w_{i(n)}, w_{j(n)})), \quad \forall n \geq \mathbb{N}. \quad (3.9)$$

From (F'_3) , (3.5) and (3.9), we obtain $\tau + F(\varepsilon) \leq F(\varepsilon)$. This contradiction shows that $\{w_n\}_{n=1}^\infty$ is a Cauchy sequence. Since W is complete, there exists $t \in W$ such that $\lim_{n \rightarrow \infty} w_n = t$. Since Y is \perp -continuous, we have

$$Yt = Y(\lim_{n \rightarrow \infty} w_n) = \lim_{n \rightarrow \infty} w_{n+1} = t$$

and so t is a fixed point of Y .

Next, we show that t is a unique fixed point of Y . Indeed, if $s, r \in W$ be two distinct fixed points of Y , that is $Ys = s \neq r = Yr$. Therefore,

$$d(Ys, Yr) = d(s, r) > 0. \quad (3.10)$$

Since Y is \perp -preserving, we have

$$s \perp r \quad \text{or} \quad r \perp s.$$

Since Y is generalized orthogonal F -contraction mapping, we have

$$\begin{aligned} F(d(s, r)) &= F(d(Ys, Yr)) < \tau + F(d(Ys, Yr)) \\ &\leq F(d(s, r)), \end{aligned}$$

which is a contradiction. Hence, Y has a unique fixed point. \square

Next, we give the fixed point theorem for an orthogonal F -Suzuki contraction mapping in an O -complete orthogonal metric space (W, \perp, d) .

Theorem 3.4. *Let (W, \perp, d) be an O -complete orthogonal metric space with an orthogonal element w_0 and a mapping $Y : W \rightarrow W$. Suppose that there exist $F \in \mathcal{D}$ and $\tau > 0$ such that the following conditions hold:*

- (i) Y is \perp -preserving;
- (ii) Y is an orthogonal F -Suzuki contraction mapping;
- (iii) Y is \perp -continuous.

Then Y has a unique fixed point $t \in W$ and for every $w \in W$, the sequence $\{Y^n w\}$ converges to t .

Proof. Since (W, \perp) is an O -set,

$$\exists w_0 \in W : (\forall w \in W, w \perp w_0) \quad \text{or} \quad (\forall w \in W, w_0 \perp w).$$

It follows that $w_0 \perp Yw_0$ or $Yw_0 \perp w_0$. Let

$$\begin{aligned} w_1 &:= Yw_0, w_2 := Yw_1 = Y^2w_0 \dots \\ \dots, w_{n+1} &:= Yw_n = Y^{n+1}w_0 \end{aligned}$$



for all $n \in \mathbb{N} \cup \{0\}$. If $w_n = w_{n+1}$ for any $n \in \mathbb{N} \cup \{0\}$, then it is clear that w_n is a fixed point of Y . Assume that $w_n \neq w_{n+1}$ for all $n \in \mathbb{N} \cup \{0\}$. Thus, we have $d(Yw_n, Yw_{n+1}) > 0$ for all $n \in \mathbb{N} \cup \{0\}$. Now,

$$\frac{1}{2}d(w_n, Yw_n) < d(w_n, Yw_n), \quad \forall n \in \mathbb{N}.$$

Since Y is \perp -preserving, we have

$$w_n \perp w_{n+1} \quad \text{or} \quad w_{n+1} \perp w_n \quad (3.11)$$

for all $n \in \mathbb{N} \cup \{0\}$. This implies that $\{w_n\}$ is an O -sequence. Since Y is an orthogonal F -Suzuki contraction mapping, we have

$$F(d(Yw_{n-1}, Yw_n)) \leq F(d(w_{n-1}, w_n)) - \tau, \quad \forall n \in \mathbb{N}.$$

Repeating this process, we get

$$\begin{aligned} F(d(Yw_{n-1}, Yw_n)) &\leq F(d(w_{n-1}, w_n)) - \tau \\ &= F(d(Yw_{n-2}, Yw_{n-1})) - \tau \\ &\leq F(d(w_{n-2}, w_{n-1})) - 2\tau \\ &= F(d(Yw_{n-3}, Yw_{n-2})) - 2\tau \\ &\leq F(d(w_{n-3}, w_{n-2})) - 3\tau \\ &\vdots \\ &\leq F(d(w_0, w_1)) - n\tau. \end{aligned} \quad (3.12)$$

From (3.12), we obtain $\lim_{n \rightarrow \infty} F(d(w_n, Yw_n)) = -\infty$.

By the property (F_2) , we have

$$\lim_{n \rightarrow \infty} d(w_n, Yw_n) = 0. \quad (3.13)$$

Now, we claim that $\{w_n\}_{n=1}^\infty$ is a Cauchy sequence. Arguing by contradiction, we assume that there exist $\varepsilon > 0$ and sequence $\{i(n)\}_{n=1}^\infty$ and $\{j(n)\}_{n=1}^\infty$ of natural numbers such that

$$\begin{aligned} i(n) > j(n) > n, \quad d(w_{i(n)}, w_{j(n)}) &\geq \varepsilon, \\ d(w_{i(n)-1}, w_{j(n)}) < \varepsilon, \quad \forall n \in \mathbb{N}. \end{aligned} \quad (3.14)$$

So, we have

$$\begin{aligned} \varepsilon &\leq d(w_{i(n)}, w_{j(n)}) \leq d(w_{i(n)}, w_{i(n)-1}) + d(w_{i(n)-1}, w_{j(n)}) \\ &< d(w_{i(n)}, w_{i(n)-1}) + \varepsilon \\ &= d(w_{i(n)-1}, Yw_{i(n)-1}) + \varepsilon. \end{aligned}$$

It follows from (3.13) and the above inequality that

$$\lim_{n \rightarrow \infty} d(w_{i(n)}, w_{j(n)}) = \varepsilon. \quad (3.15)$$

On the other hand, from (3.13) there exists $N \in \mathbb{N}$, such that

$$d(w_{i(n)}, Yw_{i(n)}) < \frac{\varepsilon}{4} \quad (3.16)$$

$$\text{and } d(w_{j(n)}, Yw_{j(n)}) < \frac{\varepsilon}{4}, \quad \forall n \in \mathbb{N}. \quad (3.17)$$

Next, we claim that

$$d(Yw_{i(n)}, Yw_{j(n)}) = d(w_{i(n)+1}, w_{j(n)+1}) > 0, \quad \forall n \in \mathbb{N}. \quad (3.18)$$

Arguing by contradiction, there exists $m \geq \mathbb{N}$ such that

$$d(w_{i(m)+1}, w_{j(m)+1}) = 0. \quad (3.19)$$

It follows from (3.14), (3.17) and (3.19) that

$$\begin{aligned} \varepsilon &\leq d(w_{i(m)}, w_{j(m)}) \leq d(w_{i(m)}, w_{i(m)+1}) \\ &\quad + d(w_{i(m)+1}, w_{j(m)}) \\ &\leq d(w_{i(m)}, w_{i(m)+1}) \\ &\quad + d(w_{i(m)+1}, w_{j(m)+1}) \\ &\quad + d(w_{j(m)+1}, w_{j(m)}) \\ &= d(w_{i(m)}, Yw_{i(m)}) \\ &\quad + d(w_{i(m)+1}, w_{j(m)+1}) \\ &\quad + d(w_{j(m)}, Yw_{j(m)}) \\ &< \frac{\varepsilon}{4} + 0 + \frac{\varepsilon}{4} = \frac{\varepsilon}{2}. \end{aligned}$$

This contradiction establishes the relation (3.18). Since Y is \perp -preserving, we have

$$w_{i(n)} \perp w_{j(n)} \quad \text{or} \quad w_{j(n)} \perp w_{i(n)}.$$

From (3.13) and (3.15), we can choose a positive integer $N \in \mathbb{N}$ such that

$$\frac{1}{2}d(w_{i(n)}, Yw_{i(n)}) < \frac{1}{2}\varepsilon < d(w_{i(n)}, w_{j(n)}), \quad \forall n \in \mathbb{N}.$$

Since Y is F -Suzuki contraction mapping, we have

$$\tau + F(d(Yw_{i(n)}, Yw_{j(n)})) \leq F(d(w_{i(n)}, w_{j(n)})), \quad \forall n \geq \mathbb{N}. \quad (3.20)$$

From (F'_3) , (3.15) and (3.20), we obtain $\tau + F(\varepsilon) \leq F(\varepsilon)$. This contradiction shows that $\{w_n\}_{n=1}^\infty$ is a Cauchy sequence. By completeness of (W, d) , $\{w_n\}_{n=1}^\infty$ converges to some point $t \in W$. Therefore,

$$\lim_{n \rightarrow \infty} d(w_n, t) = 0. \quad (3.21)$$

Now, we claim that

$$\begin{aligned} \frac{1}{2}d(w_n, Yw_n) &< d(w_n, t) \\ \text{or } \frac{1}{2}d(Yw_n, Y^2w_n) &< d(Yw_n, t), \quad \forall n \in \mathbb{N}. \end{aligned} \quad (3.22)$$

Again, assume that there exists $m \in \mathbb{N}$ such that

$$\begin{aligned} \frac{1}{2}d(w_m, Yw_m) &\geq d(w_m, t) \\ \text{and } \frac{1}{2}d(Yw_m, Y^2w_m) &\geq d(Yw_m, t). \end{aligned} \quad (3.23)$$



Therefore,

$$2d(w_m, t) \leq d(w_m, Yw_m) \leq d(w_m, t) + d(t, Yw_m),$$

which implies that

$$d(w_m, t) \leq d(t, Yw_m). \quad (3.24)$$

It follows from (3.23) and (3.24) that

$$d(w_m, t) \leq d(t, Yw_m) \leq \frac{1}{2}d(Yw_m, Y^2w_m). \quad (3.25)$$

Since $\frac{1}{2}d(w_m, Yw_m) < d(w_m, Yw_m)$, by the assumption of the theorem, we get

$$\tau + F(d(Yw_m, Y^2w_m)) \leq F(d(w_m, Yw_m)).$$

Since $\tau > 0$, this implies that

$$F(d(Yw_m, Y^2w_m)) < F(d(w_m, Yw_m)).$$

So, from (F_1) , we get

$$d(Yw_m, Y^2w_m) < d(w_m, Yw_m). \quad (3.26)$$

It follows that from (3.23), (3.25) and (3.26) that

$$\begin{aligned} d(Yw_m, Y^2w_m) &< d(w_m, Yw_m) \\ &\leq d(w_m, t) + d(t, Yw_m) \\ &\leq \frac{1}{2}d(Yw_m, Y^2w_m) + \frac{1}{2}d(Yw_m, Y^2w_m) \\ &= d(Yw_m, Y^2w_m). \end{aligned}$$

This is a contradiction. Hence, (3.22) holds. So, from (3.22), for every $n \in \mathbb{N}$, either

$$\begin{aligned} \tau + F(d(Yw_n, Yt)) &\leq F(d(w_n, t)), \\ \text{or} \\ \tau + F(d(Y^2w_n, Yt)) &\leq F(d(Yw_n, t)) = F(d(w_{n+1}, t)) \end{aligned}$$

holds. In the first case, from (3.21), we get

$$\lim_{n \rightarrow \infty} F(d(Yw_n, Yt)) = -\infty.$$

By the property (F_2) , we have

$$\lim_{n \rightarrow \infty} d(Yw_n, Yt) = 0.$$

Therefore,

$$d(t, Yt) = \lim_{n \rightarrow \infty} d(w_{n+1}, Yt) = \lim_{n \rightarrow \infty} d(Yw_n, Yt) = 0.$$

Also, in the second case, from (3.21), we get

$$\lim_{n \rightarrow \infty} F(d(Y^2w_n, Yt)) = -\infty.$$

By the property (F_2) , we have

$$\lim_{n \rightarrow \infty} d(Y^2w_n, Yt) = 0.$$

Therefore,

$$d(t, Yt) = \lim_{n \rightarrow \infty} d(w_{n+2}, Yt) = \lim_{n \rightarrow \infty} d(Y^2w_n, Yt) = 0.$$

Hence, t is a fixed point of Y . Next, we show that t is a unique fixed point of Y . Indeed, if $s, r \in W$ be two distinct fixed points of Y , that is $Ys = s \neq r = Yr$, then $d(Ys, Yr) > 0$. Since Y is \perp -preserving, we have

$$s \perp r \text{ or } r \perp s.$$

We have $0 = \frac{1}{2}d(s, Ys) < d(s, r)$. Since Y is F -Suzuki contraction mapping, we have

$$\begin{aligned} F(d(s, r)) &= F(d(Ys, Yr)) < \tau + F(d(Ys, Yr)) \\ &\leq F(d(s, r)), \end{aligned} \quad (3.27)$$

which is a contradiction. Hence, Y has a unique fixed point. \square

4. Example

Example 4.1. Let $W = [0, \infty)$ and $d : W \times W \rightarrow [0, \infty)$ be a mapping defined by $d(u, v) = |u - v|$ for all $u, v \in W$. Consider the sequence $\{S_p\}_{p \in \mathbb{N}}$ defined as

$$S_p = \frac{p(p+1)}{2}, \quad \forall p \in \mathbb{N} \cup \{0\}.$$

Define a relation \perp on X by

$$u \perp v \iff uv \in \{u, v\} \subseteq \{S_p\}.$$

Thus (W, \perp, d) is an O -complete metric space. Now, we will define a mapping $Y : W \rightarrow W$ by

$$Yu = \begin{cases} S_1 & \text{if } S_0 \leq u \leq S_1, \\ S_{p-1} & \text{if } S_p \leq u \leq S_{p+1} \quad \forall p > 1. \end{cases} \quad (4.1)$$

It is easy to see that Y is \perp -continuous and Y is \perp -preserving. Let $u, v \in W$ with $u \perp v$ and $d(Yu, Yv) > 0$. Without loss of generality, we may assume that $u < v$. This implies that $u \in \{S_0, S_1\}$ and $v = S_i$ for some $i \in \mathbb{N} \setminus \{1\}$.

$$\begin{aligned} \lim_{p \rightarrow \infty} \frac{d(Yu, Yv)}{d(u, v)} &= \lim_{p \rightarrow \infty} \frac{S_{p-1} - 1}{S_p - 1} \\ &= \lim_{p \rightarrow \infty} \frac{\frac{p(p-1)}{2} - 1}{\frac{p(p+1)}{2} - 1} \\ &= \lim_{p \rightarrow \infty} \frac{p(p-1) - 2}{p(p+1) - 2} = 1, \end{aligned} \quad (4.2)$$

Y is not a Banach contraction and a orthogonal F -Suzuki contraction. On the other hand taking $F(\beta) = \frac{-1}{\beta} + \beta \in \mathcal{D}$, we obtain the result that Y is an orthogonal F -Suzuki contraction with $\tau = 2$. To see this, let us consider the following calculation. First observe that

$$\begin{aligned} \frac{1}{2}d(S_p, YS_p) &< d(S_p, S_q) \\ &\iff [(1 = p < q) \vee (1 \leq q < p) \\ &\vee (1 < p < q)]. \end{aligned} \quad (4.3)$$



For $1 < p < q$, we have

$$\begin{aligned} |Y(S_q) - Y(S_1)| &= |S_{q-1} - S_1| \\ &= 2 + 3 + 4 + \dots + q - 1, \\ |S_q - S_1| &= 2 + 3 + \dots + q. \end{aligned} \quad (4.4)$$

Since $q > 1$ and $\frac{-1}{2+3+\dots+q-1} < \frac{-1}{2+3+\dots+q}$, we have

$$\begin{aligned} 2 - \frac{-1}{2+3+\dots+q-1} + [2+3+\dots+q-1] \\ < 2 - \frac{1}{2+3+\dots+q} + [2+3+\dots+q-1] \\ \leq \frac{-1}{2+3+\dots+q} + [2+3+\dots+q-1] + q. \end{aligned}$$

So, from (4.4), we get

$$\begin{aligned} 2 - \frac{1}{|Y(S_q) - Y(S_1)|} + |Y(S_q) - Y(S_1)| \\ < -\frac{1}{|S_q - S_1|} + |S_q - S_1|. \end{aligned}$$

For $1 \leq p < q$, similar to $1 = p < q$, we have

$$\begin{aligned} 2 - \frac{1}{|Y(S_q) - Y(S_1)|} + |Y(S_q) - Y(S_1)| \\ < -\frac{1}{|S_q - S_1|} + |S_q - S_1|. \end{aligned}$$

For $1 < p < q$, we have

$$\begin{aligned} |Y(S_q) - Y(S_p)| &= p + (p+1) + (p+2) + \dots + q - 1, \\ |S_q - S_p| &= p + (p+1) + (p+2) + \dots + q. \end{aligned} \quad (4.5)$$

Since $q > p > 1$ and $\frac{-1}{p+(p+1)+(p+2)+\dots+q-1} < \frac{-1}{p+(p+1)+(p+2)+\dots+q}$. Therefore

$$\begin{aligned} 2 - \frac{1}{p+(p+1)+(p+2)+\dots+q-1} \\ + [p+(p+1)+(p+2)+\dots+q-1] \\ < 2 - \frac{1}{p+(p+1)+(p+2)+\dots+q} \\ + [p+(p+1)+(p+2)+\dots+q-1] \\ \leq -\frac{1}{p+(p+1)+(p+2)+\dots+q} \\ + [p+(p+1)+(p+2)+\dots+q-1] + q \end{aligned}$$

So, from (4.5), we get

$$\begin{aligned} 2 - \frac{1}{|Y(S_q) - Y(S_p)|} + |Y(S_q) - Y(S_p)| \\ < -\frac{1}{|S_q - S_p|} + |S_q - S_p|. \end{aligned}$$

Therefore $\tau + F(d(Y(S_q), Y(S_p))) \leq F(d(S_q, S_p))$ for all $p, q \in \mathbb{N}$. Hence Y is an orthogonal F -Suzuki contraction and $Y(S_1) = S_1$.

5. Application to ordinary differential equations

Recall that, for any $1 \leq p < \infty$, the space $L^p(W, F, \gamma)$ (or $L^p(W)$) consists of all complex-valued measurable ζ on the underlying space W satisfying

$$\int_W |\zeta(v)|^p d\gamma(v),$$

where F is the σ -algebra of measurable sets and γ is the measure. When $p = 1$, the space $L^1(W)$ consists of all integrable functions ζ on W and we define the L^1 -norm of ζ by

$$\|\zeta\|_1 = \int_W |\zeta(v)| d\gamma(v).$$

In this section, using Theorem 3.3, we show the existence of a solution of the following differential equation:

$$\begin{cases} w'(t) = Y(t, w(t)), \text{ a.e. } t \in I := [0, T]; \\ w(0) = c, c \geq 1, \end{cases} \quad (5.1)$$

where $Y : I \times \mathbb{R} \rightarrow \mathbb{R}$ is an integrable function satisfying the following conditions:

- (i) $Y(s, x) \geq 0$ for all $x \geq 0$ and $s \in I$;
- (ii) for each $i, j \in L^1(I)$ with $i(s)j(s) \geq i(s)$ or $i(s)j(s) \geq j(s)$ for all $s \in I$, there exist $\zeta \in L^1(I)$ and $\tau > 0$ such that

$$|Y(s, i(s)) - Y(s, j(s))| \leq \frac{\zeta(s)}{1 + \tau \zeta(s)} |i(s) - j(s)| \quad (5.2)$$

and

$$|i(s) - j(s)| \leq \zeta(s) e^{D(s)}$$

for all $s \in I$, where $D(s) := \int_0^s |\zeta(r)| dr$.

Theorem 5.1. Consider the differential Eq.(5.1). If (i) and (ii) are satisfied, then the differential Eq. (5.1) has a unique positive solution.

Proof. Let $W = \{w \in C(I, \mathbb{R}) : w(t) > 0 \text{ for all } t \in I\}$. Define the orthogonality relation \perp on V by

$$i \perp j \iff i(t)j(t) \geq i(t) \text{ or } i(t)j(t) \geq j(t), \forall t \in I.$$

Since $D(t) = \int_0^t |\zeta(s)| ds$, we have $D'(t) = |\zeta(t)|$ for almost everywhere $t \in I$. Define a mapping $d : W \times W \rightarrow [0, \infty)$ by

$$d(i, j) = \|i - j\|_D = \sup_{t \in I} e^{-D(t)} |i(t) - j(t)|$$

for all $i, j \in W$. Thus, (W, d) is a metric space and also a complete metric space (see, [1] for details). Define a mapping $G : W \rightarrow W$ by

$$(Gi)(t) = c + \int_0^t Y(s, i(s)) ds.$$



Then, we see that G is \perp -continuous.

Now, we show that G is \perp -preserving. For each $i, j \in W$ with $i \perp j$ and $t \in I$, we have

$$(Gi)(t) = c + \int_0^t Y(s, i(s)) ds \geq 1.$$

It follows that $[(Gi)(t)][(Gj)(t)] \geq (Gj)(t)$

and so $(Gi)(t) \perp (Gj)(t)$. Then, G is \perp -preserving.

Next, we claim that G is generalized orthogonal F -contraction.

Let $i, j \in W$ with $i \perp j$. Suppose that $G(i) \neq G(j)$. For each $s \in I$, we have

$$|i(s) - j(s)| \leq \zeta(s)e^{D(s)}.$$

and so

$$d(i, j) = \sup_{s \in I} e^{-D(s)} |i(s) - j(s)| \leq \zeta(s).$$

It follows that

$$1 + \tau d(i, j) \leq 1 + \tau \zeta(s).$$

From (ii), for each $t \in I$, we obtain

$$\begin{aligned} |(Gi)(t) - (Gj)(t)| &\leq \int_0^t |Y(s, i(s)) - Y(s, j(s))| ds \\ &\leq \int_0^t \frac{|\zeta(s)|}{1 + \tau \zeta(s)} |i(s) - j(s)| ds \\ &\leq \int_0^t \frac{|\zeta(s)|}{1 + \tau d(i, j)} |i(s) - j(s)| \\ &\quad e^{-D(s)} e^{D(s)} ds \\ &\leq \frac{d(i, j)}{1 + \tau d(i, j)} \int_0^t |\zeta(s)| e^{D(s)} ds \\ &\leq \frac{d(i, j)}{1 + \tau d(i, j)} \left(e^{D(t)} - 1 \right) \\ &\leq \frac{d(i, j)}{1 + \tau d(i, j)} \left(e^{D(t)} - 1 \right) \end{aligned}$$

and so

$$\begin{aligned} e^{-D(t)} |(Gi)(t) - (Gj)(t)| &\leq e^{-D(t)} \left(e^{D(t)} - 1 \right) \\ &\quad \frac{d(i, j)}{1 + \tau d(i, j)} \\ &= \left(1 - e^{-D(t)} \right) \frac{d(i, j)}{1 + \tau d(i, j)} \\ &\leq \left(1 - e^{-\|\zeta\|_1} \right) \frac{d(i, j)}{1 + \tau d(i, j)}. \end{aligned}$$

Thus, it follows that

$$\begin{aligned} d(Gi, Gj) &\leq \frac{d(i, j)}{1 + \tau d(i, j)} \\ \frac{1 + \tau d(i, j)}{d(i, j)} &\leq \frac{1}{d(Gi, Gj)} \\ \tau + \frac{1}{d(i, j)} &\leq \frac{1}{d(Gi, Gj)} \\ \tau + \frac{1}{d(i, j)} &\leq \frac{1}{d(Gi, Gj)} \\ \tau - \frac{1}{d(Gi, Gj)} &\leq -\frac{1}{d(i, j)}. \end{aligned}$$

Taking a function $F : \mathbb{R}^+ \rightarrow \mathbb{R}$ by $F(z) = -\frac{1}{z}$ for all $z > 0$, it follows that G is generalized orthogonal F -contraction. By Theorem 3.3, G has a unique fixed point and hence the differential Eq. (5.1) has a unique positive solution. \square

6. Conclusion

In this paper, we proved fixed point theorems for generalized orthogonal F -contraction and F -Suzuki contraction mappings on O -complete metric space.

References

- [1] M. E. Gordji, M. Ramezani, M. De La Sen, and Y. J. Cho, On orthogonal sets and Banach fixed point theorem, *Fixed Point Theory (FPT)*, 18(2), (2017)569-578.
- [2] M. Eshaghi Gordji and H. Habibi, Fixed point theory in generalized orthogonal metric space, *Journal of Linear and Topological Algebra (JLTA)*, 6(3), 251-260.
- [3] K. Sawangsup, W. Sintunavarat and Y. J. Cho, Fixed point theorems for orthogonal F -contraction mappings on O -complete metric spaces, *J. Fixed Point Theory Appl.*, (2020)22:10.
- [4] M. Eshaghi and H. Habibi, Fixed point theory in ε -connected orthogonal metric space, *Shand Communications in Mathematical Analysis(SCMA)* Vol. 16, No.1(2019), 35-46.
- [5] N. B. Gungor, and D. Turkoglu, Fixed point theorems on orthogonal metric spaces via altering distance functions, *AIP Conference Proceedings*, 2183, 040011(2019).
- [6] O. Yamaod and W. Sintunavarat, On new orthogonal contractions in b-metric spaces, *International Journal of Pure Mathematics*, Vol. 5, 2018.
- [7] K. Sawangsup and W. Sintunavarat, Fixed point results for orthogonal Z -contraction mappings in O -complete metric space, *International Journal of Applied Physics and Mathematics*, Vol. 10, No. 1, January 2020.
- [8] T. Senapati, L. K. Dey, B. Damjanović and A. Chanda, New fixed results in orthogonal metric spaces with an Application, *Kragujevac Journal of Mathematics*, Vol. 42(4)(2018), 505-516.



- [9] D. Wardowski, Fixed points of a new type of contractive mappings in complete metric spaces, *Fixed Point Theory Appl.* 2012, 94(2012).
- [10] H. Piri, P. Kumam, Some fixed point theorems concerning F -contraction in complete metric spaces, *Fixed Point Theory Appl.* 210(2014).
- [11] N. Sharma, L. N. Mishra, V. N. Mishra, H. Almusawa, Endpoint approximation of standard three-step multi-valued iteration algorithm for nonexpansive mappings, *Applied Mathematics and Information Sciences*, Vol. 15, No. 1, (2021), pp. 73-81. DOI: 10.18576/amis/150109
- [12] N. Sharma, L. N. Mishra, V. N. Mishra, S. Pandey, Solution of Delay Differential equation via N_1^y iteration algorithm, *European J. Pure Appl. Math.*, Vol. 13, No. 5, (2020), pp. 1110-1130. DOI: <https://doi.org/10.29020/nybg.ejpam.v13i5.3756>.
- [13] L. N. Mishra, V. Dewangan, V. N. Mishra, S. Karateke, Best proximity points of admissible almost generalized weakly contractive mappings with rational expressions on b-metric spaces, *J. Math. Computer Sci.*, Vol. 22, Issue 2, (2021), pp. 97–109. doi: 10.22436/jmcs.022.02.01.
- [14] A. G. Sanatee, M. Iranmanesh, L. N. Mishra, V. N. Mishra, Generalized 2–proximal C –contraction mappings in complete ordered 2–metric space and their best proximity points, *Scientific publications of the state university of Novi pazar ser. A: Appl. Math. Inform. and Mech.* vol. 12, 1 (2020), 1-11.
- [15] L. N. Mishra, V. N. Mishra, P. Gautam, K. Negi, Fixed point Theorems for Cyclic-Ćirić-Reich-Rus contraction mapping in Quasi-Partial b-metric spaces, *Scientific publications of the state university of Novi pazar ser. A: Appl. Math. Inform. and Mech.* vol. 12, 1 (2020), 47-56.
- [16] L. N. Mishra, S. K. Tiwari, V. N. Mishra, I. A. Khan, Unique Fixed Point Theorems for Generalized Contractive Mappings in Partial Metric Spaces, *Journal of Function Spaces*, Volume 2015 (2015), Article ID 960827, 8 pages.
- [17] L. N. Mishra, S. K. Tiwari, V. N. Mishra, Fixed point theorems for generalized weakly S-contractive mappings in partial metric spaces, *Journal of Applied Analysis and Computation*, Volume 5, Number 4, 2015, pp. 600-612. doi:10.11948/2015047

ISSN(P):2319 – 3786
Malaya Journal of Matematik
ISSN(O):2321 – 5666

