



# Solution of one dimensional wave equation and wave-like equations with variable coefficients using variational iteration method

E. Rama<sup>1\*</sup> and K. Sambaiah<sup>3</sup>

## Abstract

Present work is devoted to application of variational iteration method to obtain the solutions of wave equation and wave-like equation. Numerical examples are taken to test the efficiency of this method. We have shown that the successive approximations of each problem are converging to their exact solution. Further we have shown graphically the fourth approximation values and exact values.

## Keywords

Variational iteration method, one dimensional wave equation, successive approximation.

## AMS Subject Classification

65F10.

<sup>1</sup> Department of Mathematics, University College of Science, Saifabad, Osmania University, Hyderabad, Telangana State, India.

<sup>2</sup> Formerly Professor of Mathematics, Kakatiya University, Flat No. 101, Riddhis Pebble, Madhapur, Hyderabad, Telangana, India.

\*Corresponding author: <sup>1</sup> ramamathsou@gmail.com

Article History: Received 29 January 2021; Accepted 18 March 2021

©2021 MJM.

## Contents

1	Introduction .....	902
2	Description of the method .....	902
3	Illustrative examples .....	903
	References .....	904

## 1. Introduction

Many problems in various fields of sciences and engineering yield partial differential equation. For their physical interpretation we need their solutions. It may not possible to find the exact solution of certain partial differential equations. There are many methods to find the approximate solutions of such equation. For example, numerical methods, Adomain decomposition method, Homotopy analysis method etc. J.H.He [2, 3] developed the variational iteration method. Several researchers [1, 4, 6?] working on application of this method to find the solutions of linear and nonlinear partial differential equations. E.Rama, K. Somaiah and K. Sambaiah [5] studied variational iteration method for solving various types of problems. In this paper we found the successive approximations of one dimensional wave equation and wavelike equation using variational iteration method. Further it was shown that they are converging to their exact solution.

## 2. Description of the method

Here we describe briefly the basic idea of the VIM to solve partial differential equation. Consider the following partial differential equation

$$L_t u(x,t) + L_x u(x,t) + Nu(x,t) = g(x,t), \quad (1.1)$$

where  $L_t$  is a linear operator involving partial derivatives with respect to  $t$  only,  $L_x$  is linear operator involving partial derivatives with respect to  $x$  only,  $N$  is non-linear operator and  $g(x,t)$  is a continuous function. According to VIM the correctional functional of equation (1) in  $t$  and  $x$  directions respectively are given by

$$u_{n+1}(x,t) = u_n(x,t) + \int_0^t \lambda_1(x,s) [L_s u_n + \{L_x + N\} \tilde{u}_n(x,s) - g(x,s)] ds \quad (1.2)$$

$$u_{n+1}(x,t) = u_n(x,t) + \int_0^t \lambda_2(x,s) [L_s u_n + \{L_t + N\} \tilde{u}_n(x,s) - g(x,s)] ds, \quad (1.3)$$

where  $u_n(x,t)$  denotes  $n$ th approximation of  $u(x,t)$ ,  $\lambda_1$  and  $\lambda_2$  are Lagrangian multipliers which can be identified optimally using the variational theory and  $\tilde{u}_n$  is the restricted variation.

i.e.,  $\tilde{\delta}u_n = 0$ . After determination of Lagrange multipliers  $\lambda_1$  and  $\lambda_2$  the successive approximations  $u_{n+1}(x, t)$  of  $u(x, t)$  can be obtained by suitable choice of initial approximation  $u_0(x, t)$ .

The solution of equation (1.1) is obtained as

$$u(x, t) = \lim_{n \rightarrow \infty} u_n(x, t). \quad (1.4)$$

### 3. Illustrative examples

#### Example 3.1:

Consider one dimensional wave equation

$$\frac{\partial^2 u(x, t)}{\partial t^2} - 9 \frac{\partial^2 u(x, t)}{\partial x^2} = 0 \quad (1.5)$$

with the following initial and boundary conditions i)  $u(0, t) = 0$  ii)  $u(1, t) = 0$  iii)  $u(x, 0) = 6 \sin(\pi x)$  iv)  $\frac{\partial u(x, t)}{\partial t} = 0$  at  $t = 0$ . According to the VIM the correctional formula for the wave equation (1.5) becomes

$$u_{n+1}(x, t) = u_n(x, t) + \int_0^t \lambda(s, t) \left( \frac{\partial^2}{\partial s^2} u_n(x, s) - 9 \frac{\partial^2}{\partial x^2} \tilde{u}_n(x, s) \right) ds \quad (1.6)$$

The equation (1.6) gives the following stationary conditions noting that  $\tilde{\delta}u_n = 0$ .

$$1 - \lambda'(s, t) = 0 \text{ at } s = t.$$

$$\lambda(s, t) = 0 \text{ at } s = t.$$

$$\lambda''(s, t) = 0 \text{ at } s = t,$$

where dash denotes partial derivative with respect to  $s$ .

From the above three equations we have

$$\lambda(s, t) = s - t \quad (1.7)$$

Hence the variational iteration formula for the problem considered is

$$u_{n+1}(x, t) = u_n(x, t) + \int_0^t (s - t) \left( \frac{\partial^2}{\partial s^2} u_n(x, s) - 9 \frac{\partial^2}{\partial x^2} u_n(x, s) \right) ds \quad (1.8)$$

Choose  $u_0(x, t) = u(x, 0) = 6 \sin(\pi x)$ . Substituting  $n = 0$  and the initial approximation in equation (1.8) we get

$$u_1(x, t) = u_0(x, t) + \int_0^t (s - t) \left( \frac{\partial^2}{\partial s^2} u_0(x, s) - 9 \frac{\partial^2}{\partial x^2} u_0(x, s) \right) ds$$

$$u_1(x, t) = 6 \sin(\pi x) + \int_0^t (s - t) \left( 0 - 9 \frac{\partial^2}{\partial x^2} (6 \sin(\pi x)) \right) ds$$

$$u_1(x, t) = 6 \sin(\pi x) \left[ 1 - \frac{3^2 \pi^2 t^2}{2!} \right] \quad (1.9)$$

Continuing this process the successive approximation are

$$u_2(x, t) = 6 \sin(\pi x) \left[ 1 - \frac{3^2 \pi^2 t^2}{2!} + \frac{3^4 \pi^4 t^4}{4!} \right] \quad (1.10)$$


$$u_3(x, t) = 6 \sin(\pi x) \left[ 1 - \frac{3^2 \pi^2 t^2}{2!} + \frac{3^4 \pi^4 t^4}{4!} - \frac{3^6 \pi^6 t^6}{6!} \right] \quad (1.11)$$

$$u_4(x, t) = 6 \sin(\pi x) \left[ 1 - \frac{3^2 \pi^2 t^2}{2!} + \frac{3^4 \pi^4 t^4}{4!} - \frac{3^6 \pi^6 t^6}{6!} + \frac{3^8 \pi^8}{8!} \right]$$

.....

$$u(x, t) = \lim_{n \rightarrow \infty} u_n(x, t) = 6 \sin(\pi x) \cos(3\pi t) \quad (1.12)$$

The successive approximations obtained are converging to the exact solution of the equation (1.5).



For small values of  $t$  and  $x$ , the fourth approximation values are close to the exact values of  $u(x, t)$ . However the limit of these approximations is approaching to the exact solution.

#### Example 3.2

Now consider wave-like wave equation with variable coefficients

$$\frac{\partial^2 u(x, t)}{\partial t^2} + \frac{x^2}{2} \frac{\partial^2 u(x, t)}{\partial x^2} = 0 \quad (1.13)$$

with the following initial and boundary conditions. i)  $u(x, 0) = x^2$ , ii)  $u(1, t) = \cos(t)$ , iii)  $u(0, t) = 0$ , iv)  $\frac{\partial u(x, t)}{\partial t} = 0$  at  $t = 0$ .

Proceeding as in case of earlier problem, we get

$$u_{n+1}(x, t) = u_n(x, t) + \int_0^t (s - t) \left( \frac{\partial^2}{\partial s^2} u_n(x, s) + \frac{x^2}{2} \frac{\partial^2}{\partial x^2} u_n(x, s) \right) ds \quad (1.14)$$

Choose the initial approximation  $u_0(x, t) = u(x, 0) = x^2$ . Substituting  $n = 0$  and using the initial approximation we

get

$$u_1(x, t) = x^2 \left( 1 - \frac{t^2}{2!} \right) \quad (1.15)$$

$$u_2(x, t) = x^2 \left( 1 - \frac{t^2}{2!} + \frac{t^4}{4!} \right) \quad (1.16)$$



$$u_3(x, t) = x^2 \left( 1 - \frac{t^2}{2!} + \frac{t^4}{4!} - \frac{t^6}{6!} \right) \quad (1.17)$$

$$u_4(x, t) = x^2 \left( 1 - \frac{t^2}{2!} + \frac{t^4}{4!} - \frac{t^6}{6!} + \frac{t^8}{8!} \right) \quad (1.18)$$

⋮

$$u(x, t) = \lim_{n \rightarrow \infty} u_n(x, t) = x^2 \cos(t). \quad (1.19)$$





**Example 3.3**

Now consider the homogeneous equation

$$\frac{\partial^2 u(x,t)}{\partial t^2} - \frac{\partial^2 u(x,t)}{\partial x^2} + 8u(x,t) = 0 \tag{1.19}$$

With the initial and boundary conditions i)  $u(x,0) = 0$ , ii)  $u(\pi,t) = 0$ , iii)  $u(x,0) = \sin(x)$ , iv)  $\frac{\partial u(x,t)}{\partial t} = 0$  at  $t = 0$ .

The variational iteration formula corresponding to equation (1.19) is given by

$$u_{n+1}(x,t) = u_n(x,t) + \int_0^t (s-t) \left( \frac{\partial^2}{\partial s^2} u_n(x,s) - \frac{\partial^2}{\partial x^2} u_n(x,s) + 8u_n(x,s) \right) ds \tag{1.20}$$

Choose the initial approximation  $u_0(x,t) = u(x,0) = \sin(x)$ . Substituting  $n = 0$  and the initial approximation in equation (1.20) we get

$$\begin{aligned} u_1(x,t) &= u_0(x,t) + \int_0^t (s-t) \left( \frac{\partial^2}{\partial s^2} u_0(x,s) - \frac{\partial^2}{\partial x^2} u_0(x,s) + 8u_0(x,s) \right) ds \\ &= \sin(x) + \int_0^t (s-t) \left( \frac{\partial^2}{\partial s^2} \sin(x) - \frac{\partial^2}{\partial x^2} \sin(x) + 8\sin(x) \right) ds \\ &= \sin(x) + \int_0^t (s-t)(0 + \sin(x) + 8\sin(x)) ds \end{aligned}$$

Thus,

$$u_1(x,t) = \sin(x) \left( 1 - \frac{(3t)^2}{2!} \right). \tag{1.21}$$

Continuing this process we obtain

$$\begin{aligned} u_2(x,t) &= \sin(x) \left( 1 - \frac{(3t)^2}{2!} + \frac{(3t)^4}{4!} \right) \\ u_3(x,t) &= \sin(x) \left( 1 - \frac{(3t)^2}{2!} + \frac{(3t)^4}{4!} - \frac{(3t)^6}{6!} \right) \\ u_4(x,t) &= \sin(x) \left( 1 - \frac{(3t)^2}{2!} + \frac{(3t)^4}{4!} - \frac{(3t)^6}{6!} + \frac{(3t)^8}{8!} \right) \\ &\vdots \\ u(x,t) &= \lim_{n \rightarrow \infty} u_n(x,t) = \sin(x) \cos(3t). \end{aligned}$$

The successive approximations are converging to the exact solution of (1.19).

**Conclusion**

Thus VIM can be easily applicable to obtain the solutions of one-dimensional wave equation and wave-like equation. It can be observed that the approximations are moving close to the exact solution. The same can be observed from the graphs. This shows the efficacy of this method.

**References**

- [1] M. Akbarzade and J. Langari, Application of variational iteration method to partial differential equation systems, *Int. J. of Math. Anal.*, 5(18)(2011), 863–870.
- [2] J.H. He, Variational iteration method for delay differential equations, *Communications in Nonlinear Science and Numerical simulations*, 2(4)(1997), 235–236.
- [3] J.H. He, Variational iteration method –a kind of non-linear analytical-technique: some examples, *Int. J. Non-linear Mech.*, 34(4)(1999), 699–708.
- [4] M. Matinfar, M. Saeidy, M. Madhavi and M. Rezaei Variational iteration method for exact solution of gas dynamic equation using He’s polynomials, *Bulletin of Mathematical Analysis and Applications*, 3(3)(2011), 50–55.
- [5] E. Rama, K. Somaiah and K. Sambaiah, A study of variational iteration method for solving various types of problems, *Malaya Journal of Matematik*, 9(1)(2021), 701–708.
- [6] S.A. Zahedi, H. Goodazian and M. Okazi, An investigation on two dimensional non-linear wave equation using VIM, *Indian Journal of Science and Technology*, 3(9)(2010), 1006–1008.

\*\*\*\*\*  
ISSN(P):2319 – 3786  
Malaya Journal of Matematik  
ISSN(O):2321 – 5666  
\*\*\*\*\*

