



# Some trapezoid type inequalities for functions of two variables via generalized fractional integrals

Kubilay Özçelik<sup>1</sup> and Hüseyin Budak<sup>2\*</sup>

## Abstract

In this paper, we establish some trapezoid type integral inequalities for functions of two variables involving generalized fractional integrals. The results presented here provide extensions of those given in earlier works.

## Keywords

Trapezoid type inequality, fractional integrals, bounded functions.

## AMS Subject Classification

26D07, 26D10, 26D15, 26A33.

<sup>1,2</sup>Department of Mathematics, Faculty of Science and Arts, Düzce University, Düzce, Turkey.

\*Corresponding author: <sup>1</sup> [kubilayozcelik@windowslive.com](mailto:kubilayozcelik@windowslive.com); <sup>2</sup> [hsyn.budak@gmail.com](mailto:hsyn.budak@gmail.com)

Article History: Received 16 April 2020; Accepted 09 September 2020

©2020 MJM.

## Contents

1	Introduction .....	1719
2	Some Trapezoid Type Inequalities for Generalized Fractional Integrals .....	1720
	References .....	1726

## 1. Introduction

The study of various types of integral inequalities has been the focus of great attention for well over a century by a number of mathematicians, interested both in pure and applied mathematics. One of the many fundamental mathematical discoveries of A. M. Ostrowski [23] is the following classical integral inequality associated with the differentiable mappings:

**Theorem 1.1.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable mapping on  $(a, b)$  whose derivative  $f' : (a, b) \rightarrow \mathbb{R}$  is bounded on  $(a, b)$ , i.e.  $\|f'\|_\infty = \sup_{t \in (a, b)} |f'(t)| < +\infty$ . Then, we have the following inequality:

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[ \frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right] (b-a) \|f'\|_\infty,$$

for all  $x \in [a, b]$ . The constant  $\frac{1}{4}$  is the best possible.

Ostrowski inequality has applications in quadrature, probability and optimization theory, stochastic, statistics, information and integral operator theory. Until now, a large number of

research papers and books have been written on Ostrowski inequalities and their numerous applications, see [1]–[4], [6]–[26].

**Definition 1.2.** Let  $f \in L_1[a, b]$ . The Riemann–Liouville fractional integrals  $J_{a+}^\alpha f$  and  $J_{b-}^\alpha f$  of order  $\alpha > 0$  with  $a \geq 0$  are defined by

$$J_{a+}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f(t) dt, \quad x > a$$

and

$$J_{b-}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} f(t) dt, \quad x < b,$$

respectively. Here,  $\Gamma(\alpha)$  is the Gamma function and  $J_{a+}^0 f(x) = J_{b-}^0 f(x) = f(x)$ .

**Definition 1.3.** [17] Let  $g : [a, b] \rightarrow \mathbb{R}$  be a positive increasing function on  $(a, b)$ , having a continuous derivative  $g'(x)$  on  $(a, b)$ . The left–sides  $(I_{a+;g}^\alpha f(x))$  and right–sides  $(I_{c+;g}^\alpha f(x))$  fractional integral of  $f$  with respect to the function  $g$  on  $[a, b]$  of order  $\alpha > 0$  are defined by

$$I_{a+;g}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x \frac{g'(t)f(t)}{[g(x) - g(t)]^{1-\alpha}} dt, \quad x > a$$

and

$$I_{c+;g}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_x^b \frac{g'(t)f(t)}{[g(t) - g(x)]^{1-\alpha}} dt, \quad x < b,$$

respectively.

Hadamard fractional integrals of a function with two variables can be given as follows:

**Definition 1.4.** Let  $f \in L_1(\Delta = [a, b] \times [c, d])$ . The Hadamard fractional integrals  $J_{a+,c+}^{\alpha,\beta} f$ ,  $J_{a+,d-}^{\alpha,\beta} f$ ,  $J_{b-,c+}^{\alpha,\beta} f$  and  $J_{b-,d-}^{\alpha,\beta} f$  of order  $\alpha, \beta > 0$  with  $a, c \geq 0$  are defined by

$$J_{a+,c+}^{\alpha,\beta} f(x,y) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^x \int_c^y \left(\ln \frac{x}{t}\right)^{\alpha-1} \left(\ln \frac{y}{s}\right)^{\beta-1} \frac{f(t,s)}{ts} dsdt,$$

for  $x > a, y > c$ ,

$$J_{a+,d-}^{\alpha,\beta} f(x,y) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^x \int_y^d \left(\ln \frac{x}{t}\right)^{\alpha-1} \left(\ln \frac{s}{y}\right)^{\beta-1} \frac{f(t,s)}{ts} dsdt,$$

for  $x > a, y < d$ ,

$$J_{b-,c+}^{\alpha,\beta} f(x,y) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_x^b \int_c^y \left(\ln \frac{t}{x}\right)^{\alpha-1} \left(\ln \frac{y}{s}\right)^{\beta-1} \frac{f(t,s)}{ts} dsdt,$$

for  $x < b, y > c$  and

$$J_{b-,d-}^{\alpha,\beta} f(x,y) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_x^b \int_y^d \left(\ln \frac{t}{x}\right)^{\alpha-1} \left(\ln \frac{y}{s}\right)^{\beta-1} \frac{f(t,s)}{ts} dsdt,$$

for  $x < b, y < d$ , respectively.

Now, we give the following generalized fractional integral operators:

**Definition 1.5.** [5] Let  $g : [a, b] \rightarrow \mathbb{R}$  be a positive increasing function on  $(a, b)$ , having a continuous derivative  $g'(x)$  on  $(a, b)$  and let  $w : [c, d] \rightarrow \mathbb{R}$  be a positive increasing function on  $(c, d)$ , having a continuous derivative  $w'(y)$  on  $(c, d)$ . If  $f \in L_1(\Delta)$ , then for  $\alpha, \beta > 0$ , the generalized fractional integral operators for functions of two variables are defined by

$$\mathcal{J}_{a+,c+;g,w}^{\alpha,\beta} f(x,y) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \times \int_a^x \int_c^y \frac{g'(t)}{[g(x) - g(t)]^{1-\alpha}} \frac{w'(s)}{[w(y) - w(s)]^{1-\beta}} f(t,s) dsdt,$$

for  $x > a, y > c$ ,

$$\mathcal{J}_{a+,d-;g,w}^{\alpha,\beta} f(x,y) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \times \int_a^x \int_y^d \frac{g'(t)}{[g(x) - g(t)]^{1-\alpha}} \frac{w'(s)}{[w(s) - w(y)]^{1-\beta}} f(t,s) dsdt,$$

for  $x > a, y < d$ ,

$$\mathcal{J}_{b-,c+;g,w}^{\alpha,\beta} f(x,y) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \times \int_x^b \int_c^y \frac{g'(t)}{[g(t) - g(x)]^{1-\alpha}} \frac{w'(s)}{[w(y) - w(s)]^{1-\beta}} f(t,s) dsdt,$$

for  $x < b, y > c$  and

$$\mathcal{J}_{b-,d-;g,w}^{\alpha,\beta} f(x,y) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \times \int_x^b \int_y^d \frac{g'(t)}{[g(t) - g(x)]^{1-\alpha}} \frac{w'(s)}{[w(s) - w(y)]^{1-\beta}} f(t,s) dsdt,$$

for  $x < b, y < d$ , respectively.

The aim of this study is to establish Ostrowski type integral inequalities for functions of two variables involving generalized fractional integrals. The results presented in this paper provide extensions of those given in earlier works.

## 2. Some Trapezoid Type Inequalities for Generalized Fractional Integrals

Firstly, we define the following functions which will be used frequently:

$$M_g^\alpha(a,x) = \frac{[g(x) - g(a)]^\alpha}{\Gamma(\alpha + 1)},$$

$$N_w^\beta(c,y) = \frac{[w(y) - w(c)]^\beta}{\Gamma(\beta + 1)},$$

$$M_g^\alpha(b,x) = \frac{[g(b) - g(x)]^\alpha}{\Gamma(\alpha + 1)}$$

and

$$N_w^\beta(d,y) = \frac{[w(d) - w(y)]^\beta}{\Gamma(\beta + 1)}$$

for  $(x,y) \in \Delta$ .

In addition, by choosing  $g(t) = \ln t, t \in [a, b]$  and  $w(s) = \ln s, s \in [c, d]$ , we have the following representations

$$M_{\ln}^\alpha(a,x) = \frac{[\ln \frac{x}{a}]^\alpha}{\Gamma(\alpha + 1)} \text{ and } N_{\ln}^\beta(c,y) = \frac{[\ln \frac{y}{c}]^\beta}{\Gamma(\beta + 1)}$$

and

$$M_{\ln}^\alpha(b,x) = \frac{[\ln \frac{b}{x}]^\alpha}{\Gamma(\alpha + 1)} \text{ and } N_{\ln}^\beta(d,y) = \frac{[\ln \frac{d}{y}]^\beta}{\Gamma(\beta + 1)}.$$



We also denote

$$A = M_g^\alpha(a,x)N_w^\beta(c,y)f(a,c) + M_g^\alpha(a,x)N_w^\beta(d,y)f(a,d) \\ + M_g^\alpha(b,x)N_w^\beta(c,y)f(b,c) + M_g^\alpha(b,x)N_w^\beta(d,y)f(b,d)$$

for  $(x,y) \in \Delta$ .

Throughout this paper, we denote the second partial derivative  $\frac{\partial^2 f}{\partial t \partial s}$  by  $f_{ts}$ . Moreover, let  $g : [a,b] \rightarrow \mathbb{R}$  be a positive increasing function on  $(a,b]$ , having a continuous derivative  $g'(x)$  on  $(a,b)$  and let  $w : [c,d] \rightarrow \mathbb{R}$  be a positive increasing function on  $(c,d]$ , having a continuous derivative  $w'(y)$  on  $(c,d)$ .

Now, we are in position to prove the following identity:

**Lemma 2.1.** *If  $f : \Delta \rightarrow \mathbb{R}$  be a twice partial differentiable mapping on  $\Delta^\circ$  with  $a < b, c < d$ . If  $\frac{\partial^2 f}{\partial t \partial s} \in L(\Delta)$ , then for  $\alpha, \beta > 0$  we have the following equality:*

$$A - N_w^\beta(c,y) \left[ J_{a+;g}^\alpha f(x,c) + J_{b-;g}^\alpha f(x,c) \right] \\ - N_w^\beta(d,y) \left[ J_{a+;g}^\alpha f(x,d) + J_{b-;g}^\alpha f(x,d) \right] \\ - M_g^\alpha(a,x) \left[ J_{c+;w}^\beta f(a,y) + J_{d-;w}^\beta f(a,y) \right] \\ - M_g^\alpha(b,x) \left[ J_{c+;w}^\beta f(b,y) + J_{d-;w}^\beta f(b,y) \right] \\ + \mathcal{J}_{a+,c+;g,w}^{\alpha,\beta} f(x,y) + \mathcal{J}_{a+,d-;g,w}^{\alpha,\beta} f(x,y) \\ + \mathcal{J}_{b-,c+;g,w}^{\alpha,\beta} f(x,y) + \mathcal{J}_{b-,d-;g,w}^{\alpha,\beta} f(x,y) \\ = \frac{1}{\Gamma(\alpha+1)\Gamma(\beta+1)} [I_1 - I_2 - I_3 + I_4],$$

where

$$I_1 = \int_a^x \int_c^y (g(x) - g(t))^\alpha (w(y) - w(s))^\beta f_{ts}(t,s) ds dt,$$

$$I_2 = \int_a^x \int_y^d (g(x) - g(t))^\alpha (w(s) - w(y))^\beta f_{ts}(t,s) ds dt,$$

$$I_3 = \int_x^b \int_c^y (g(t) - g(x))^\alpha (w(y) - w(s))^\beta f_{ts}(t,s) ds dt$$

and

$$I_4 = \int_x^b \int_y^d (g(t) - g(x))^\alpha (w(s) - w(y))^\beta f_{ts}(t,s) ds dt.$$

*Proof.* By integration by parts, we have

$$I_1 \tag{2.1} \\ = \int_a^x \int_c^y (g(x) - g(t))^\alpha (w(y) - w(s))^\beta f_{ts}(t,s) ds dt \\ = \int_a^x (g(x) - g(t))^\alpha \\ \times \left[ (w(y) - w(s))^\beta f_t(t,s) \Big|_c^y \right. \\ \left. + \beta \int_c^y (w(y) - w(s))^{\beta-1} w'(s) f_{ts}(t,s) ds \right] dt \\ = - \int_a^x (g(x) - g(t))^\alpha (w(y) - w(c))^\beta f_t(t,s) dt \\ + \beta \int_a^x \int_c^y (g(x) - g(t))^\alpha \\ \times (w(y) - w(s))^{\beta-1} w'(s) f_{ts}(t,s) ds dt \\ = - (w(y) - w(c))^\beta \\ \times \left[ (g(x) - g(t))^\alpha f(t,c) \Big|_a^x \right. \\ \left. + \alpha \int_a^x (g(x) - g(t))^{\alpha-1} g'(t) f(t,c) dt \right] \\ + \beta \int_c^y (w(y) - w(s))^{\beta-1} w'(s) \\ \times \left[ (g(x) - g(t))^\alpha f(t,s) \Big|_a^x \right. \\ \left. + \alpha \int_a^x (g(x) - g(t))^{\alpha-1} g'(t) f(t,s) dt \right] ds \\ = (w(y) - w(c))^\beta (g(x) - g(a))^\alpha f(a,c) \\ - \Gamma(\alpha+1) (w(y) - w(c))^\beta J_{a+;g}^\alpha f(x,c) \\ - \Gamma(\beta+1) (g(x) - g(a))^\alpha J_{c+;w}^\beta f(a,y) \\ + \Gamma(\alpha+1)\Gamma(\beta+1) J_{a+c+;g,w}^{\alpha,\beta} f(x,y).$$

Similarly, we get

$$I_2 = - (g(x) - g(a))^\alpha (w(d) - w(y))^\beta f(a,d) \tag{2.2} \\ + \Gamma(\alpha+1) (w(d) - w(y))^\beta J_{a+;g}^\alpha f(x,d) \\ + \Gamma(\beta+1) (g(x) - g(a))^\alpha J_{d-;w}^\beta f(a,y) \\ - \Gamma(\alpha+1)\Gamma(\beta+1) J_{a+d-;g,w}^{\alpha,\beta} f(x,y),$$



$$\begin{aligned}
 I_3 = & -(g(b) - g(x))^\alpha (w(y) - w(c))^\beta f(b, c) \quad (2.3) && + \mathcal{J}_{a+,c+;g,w}^{\alpha,\beta} f(x, y) + \mathcal{J}_{a+,d-;g,w}^{\alpha,\beta} f(x, y) \\
 & + \Gamma(\alpha + 1)(w(y) - w(c))^\beta J_{b-;g}^\alpha f(x, c) && + \mathcal{J}_{b-,c+;g,w}^{\alpha,\beta} f(x, y) + \mathcal{J}_{b-,d-;g,w}^{\alpha,\beta} f(x, y) \\
 & + \Gamma(\beta + 1)(g(b) - g(x))^\alpha J_{c+;w}^\beta f(b, y) && \leq \left[ M_g^{\alpha+1}(a, x) + M_g^{\beta+1}(b, x) \right] \\
 & + \Gamma(\alpha + 1)\Gamma(\beta + 1) J_{b-c+;g,w}^{\alpha,\beta} f(x, y) && \times \left[ N_g^{\alpha+1}(c, y) + N_g^{\beta+1}(d, y) \right] \left\| \frac{f_{ts}}{g'w'} \right\|_{\Delta, \infty}
 \end{aligned}$$

and

$$\begin{aligned}
 I_4 = & (g(b) - g(x))^\alpha (w(d) - w(y))^\beta f(b, d) \quad (2.4) \\
 & - \Gamma(\alpha + 1)(w(d) - w(y))^\beta J_{c+;g}^\alpha f(x, d) \\
 & - \Gamma(\beta + 1)(g(b) - g(x))^\alpha J_{d-;w}^\beta f(b, y) \\
 & + \Gamma(\alpha + 1)\Gamma(\beta + 1) J_{b-d-;g,w}^{\alpha,\beta} f(x, y).
 \end{aligned}$$

By equalities (2.1)–(2.4), we establish

$$\begin{aligned}
 & A - N_w^\beta(c, y) \left[ J_{a+;g}^\alpha f(x, c) + J_{b-;g}^\alpha f(x, c) \right] \\
 & - N_w^\beta(d, y) \left[ J_{a+;g}^\alpha f(x, d) + J_{b-;g}^\alpha f(x, d) \right] \\
 & - M_g^\alpha(a, x) \left[ J_{c+;w}^\beta f(a, y) + J_{d-;w}^\beta f(a, y) \right] \\
 & - M_g^\alpha(b, x) \left[ J_{c+;w}^\beta f(b, y) + J_{d-;w}^\beta f(b, y) \right] \\
 & + \mathcal{J}_{a+,c+;g,w}^{\alpha,\beta} f(x, y) + \mathcal{J}_{a+,d-;g,w}^{\alpha,\beta} f(x, y) \\
 & + \mathcal{J}_{b-,c+;g,w}^{\alpha,\beta} f(x, y) + \mathcal{J}_{b-,d-;g,w}^{\alpha,\beta} f(x, y) \\
 = & \frac{1}{\Gamma(\alpha + 1)\Gamma(\beta + 1)} [I_1 - I_2 - I_3 + I_4],
 \end{aligned}$$

which gives the desired result.  $\square$

**Theorem 2.2.** Under assumptions of Lemma 2.1, if the function  $\frac{f_{ts}}{g'w'} \in L_\infty(\Delta)$ , then we have the following Ostrowski type inequality for the generalized fractional integrals:

$$\begin{aligned}
 & \left| A - N_w^\beta(c, y) \left[ J_{a+;g}^\alpha f(x, c) + J_{b-;g}^\alpha f(x, c) \right] \right. \\
 & - N_w^\beta(d, y) \left[ J_{a+;g}^\alpha f(x, d) + J_{b-;g}^\alpha f(x, d) \right] \\
 & - M_g^\alpha(a, x) \left[ J_{c+;w}^\beta f(a, y) + J_{d-;w}^\beta f(a, y) \right] \\
 & \left. - M_g^\alpha(b, x) \left[ J_{c+;w}^\beta f(b, y) + J_{d-;w}^\beta f(b, y) \right] \right|
 \end{aligned}$$

where

$$\left\| \frac{f_{ts}}{g'w'} \right\|_{\Delta, \infty} = \sup_{(t,s) \in \Delta} \left| \frac{f_{ts}(t, s)}{g'(t)w'(s)} \right|.$$

*Proof.* Utilizing the Lemma 2.1, we get

$$\begin{aligned}
 & \left| A - N_w^\beta(c, y) \left[ J_{a+;g}^\alpha f(x, c) + J_{b-;g}^\alpha f(x, c) \right] \right. \quad (2.5) \\
 & - N_w^\beta(d, y) \left[ J_{a+;g}^\alpha f(x, d) + J_{b-;g}^\alpha f(x, d) \right] \\
 & - M_g^\alpha(a, x) \left[ J_{c+;w}^\beta f(a, y) + J_{d-;w}^\beta f(a, y) \right] \\
 & - M_g^\alpha(b, x) \left[ J_{c+;w}^\beta f(b, y) + J_{d-;w}^\beta f(b, y) \right] \\
 & + \mathcal{J}_{a+,c+;g,w}^{\alpha,\beta} f(x, y) + \mathcal{J}_{a+,d-;g,w}^{\alpha,\beta} f(x, y) \\
 & \left. + \mathcal{J}_{b-,c+;g,w}^{\alpha,\beta} f(x, y) + \mathcal{J}_{b-,d-;g,w}^{\alpha,\beta} f(x, y) \right| \\
 \leq & \frac{1}{\Gamma(\alpha + 1)\Gamma(\beta + 1)} [|I_1| + |I_2| + |I_3| + |I_4|].
 \end{aligned}$$

By the assumptions of Theorem 2.2, we have

$$\begin{aligned}
 & |I_1| \quad (2.6) \\
 = & \left| \int_a^x \int_c^y (g(x) - g(t))^\alpha (w(y) - w(s))^\beta f_{ts}(t, s) ds dt \right| \\
 \leq & \int_a^x \int_c^y (g(x) - g(t))^\alpha (w(y) - w(s))^\beta |f_{ts}(t, s)| ds dt \\
 = & \int_a^x \int_c^y (g(x) - g(t))^\alpha (w(y) - w(s))^\beta \\
 & \times \left| \frac{f_{ts}(t, s)}{g'(t)w'(s)} \right| g'(t)w'(s) ds dt \\
 \leq & \left\| \frac{f_{ts}}{g'w'} \right\|_{[a,x] \times [c,y], \infty} \\
 & \times \int_a^x \int_c^y (g(x) - g(t))^\alpha (w(y) - w(s))^\beta g'(t)w'(s) ds dt
 \end{aligned}$$



$$= \left\| \frac{f_{ts}}{g'w'} \right\|_{[a,x] \times [c,y], \infty} \times \frac{(g(x) - g(a))^{\alpha+1} (w(y) - w(c))^{\beta+1}}{(\alpha + 1)(\beta + 1)}.$$

Similarly, we get

$$\begin{aligned} & |I_2| \tag{2.7} \\ &= \left| \int_a^x \int_y^d (g(x) - g(t))^\alpha (w(s) - w(y))^\beta f_{ts}(t, s) ds dt \right| \\ &\leq \left\| \frac{f_{ts}}{g'w'} \right\|_{[a,x] \times [y,d], \infty} \times \frac{(g(x) - g(a))^{\alpha+1} (w(d) - w(y))^{\beta+1}}{(\alpha + 1)(\beta + 1)}, \end{aligned}$$

$$\begin{aligned} & |I_3| \tag{2.8} \\ &= \left| \int_x^b \int_c^y (g(b) - g(t))^\alpha (w(s) - w(c))^\beta f_{ts}(t, s) ds dt \right| \\ &\leq \left\| \frac{f_{ts}}{g'w'} \right\|_{[x,b] \times [c,y], \infty} \times \frac{(g(b) - g(x))^{\alpha+1} (w(y) - w(c))^{\beta+1}}{(\alpha + 1)(\beta + 1)} \end{aligned}$$

and

$$\begin{aligned} & |I_4| \tag{2.9} \\ &= \left| \int_x^b \int_y^d (g(b) - g(t))^\alpha (w(d) - w(s))^\beta f_{ts}(t, s) ds dt \right| \\ &\leq \left\| \frac{f_{ts}}{g'w'} \right\|_{[x,b] \times [y,d], \infty} \times \frac{(g(b) - g(x))^{\alpha+1} (w(d) - w(y))^{\beta+1}}{(\alpha + 1)(\beta + 1)}. \end{aligned}$$

By using the inequalities (2.6)–(2.9) in (2.5), we get

$$\begin{aligned} & \left| A - N_w^\beta(c, y) \left[ J_{a+;g}^\alpha f(x, c) + J_{b-;g}^\alpha f(x, c) \right] \right. \\ & - N_w^\beta(d, y) \left[ J_{a+;g}^\alpha f(x, d) + J_{b-;g}^\alpha f(x, d) \right] \\ & - M_g^\alpha(a, x) \left[ J_{c+;w}^\beta f(a, y) + J_{d-;w}^\beta f(a, y) \right] \\ & - M_g^\alpha(b, x) \left[ J_{c+;w}^\beta f(b, y) + J_{d-;w}^\beta f(b, y) \right] \\ & + \mathcal{J}_{a+,c+;g,w}^{\alpha,\beta} f(x, y) + \mathcal{J}_{a+,d-;g,w}^{\alpha,\beta} f(x, y) \\ & \left. + \mathcal{J}_{b-,c+;g,w}^{\alpha,\beta} f(x, y) + \mathcal{J}_{b-,d-;g,w}^{\alpha,\beta} f(x, y) \right| \end{aligned}$$

$$\begin{aligned} & \leq \frac{1}{\Gamma(\alpha + 1)\Gamma(\beta + 1)} \\ & \times \left[ \left\| \frac{f_{ts}}{g'w'} \right\|_{[a,x] \times [c,y], \infty} \times \frac{(g(x) - g(a))^{\alpha+1} (w(y) - w(c))^{\beta+1}}{(\alpha + 1)(\beta + 1)} \right. \\ & + \left\| \frac{f_{ts}}{g'w'} \right\|_{[a,x] \times [y,d], \infty} \times \frac{(g(x) - g(a))^{\alpha+1} (w(d) - w(y))^{\beta+1}}{(\alpha + 1)(\beta + 1)} \\ & + \left\| \frac{f_{ts}}{g'w'} \right\|_{[x,b] \times [c,y], \infty} \times \frac{(g(b) - g(x))^{\alpha+1} (w(y) - w(c))^{\beta+1}}{(\alpha + 1)(\beta + 1)} \\ & \left. + \left\| \frac{f_{ts}}{g'w'} \right\|_{[x,b] \times [y,d], \infty} \times \frac{(g(b) - g(x))^{\alpha+1} (w(d) - w(y))^{\beta+1}}{(\alpha + 1)(\beta + 1)} \right] \\ & \leq \left[ M_g^{\alpha+1}(a, x) + M_g^{\beta+1}(b, x) \right] \\ & \times \left[ N_g^{\alpha+1}(c, y) + N_g^{\beta+1}(d, y) \right] \left\| \frac{f_{ts}}{g'w'} \right\|_{[a,b] \times [b,d], \infty}, \end{aligned}$$

which completes the proof.  $\square$

**Corollary 2.3.** *If we choose  $g(t) = t, t \in [a, b]$  and  $w(s) = s, s \in [c, d]$  in Theorem 2.2, then we have the following Ostrowski type inequality for Riemann–Liouville fractional integrals:*

$$\begin{aligned} & \left| A - \frac{(y - c)^\beta}{\Gamma(\beta + 1)} \left[ J_a^\alpha f(x, c) + J_{b-}^\alpha f(x, c) \right] \right. \tag{2.10} \\ & - \frac{(d - y)^\beta}{\Gamma(\beta + 1)} \left[ J_{a+}^\alpha f(x, d) + J_{b-}^\alpha f(x, d) \right] \\ & - \frac{(x - a)^\alpha}{\Gamma(\alpha + 1)} \left[ J_{c+}^\beta f(a, y) + J_{d-}^\beta f(a, y) \right] \\ & - \frac{(b - x)^\alpha}{\Gamma(\alpha + 1)} \left[ J_{c+}^\beta f(b, y) + J_{d-}^\beta f(b, y) \right] \\ & + J_{a+,c+}^{\alpha,\beta} f(x, y) + J_{a+,d-}^{\alpha,\beta} f(x, y) \\ & + J_{b-,c+}^{\alpha,\beta} f(x, y) + J_{b-,d-}^{\alpha,\beta} f(x, y) \left| \right. \\ & \leq \frac{1}{\Gamma(\alpha + 2)\Gamma(\beta + 2)} \left[ (x - a)^{\alpha+1} + (b - x)^{\alpha+1} \right] \\ & \times \left[ (y - c)^{\beta+1} + (d - y)^{\beta+1} \right] \|f_{ts}\|_{\Delta, \infty}. \end{aligned}$$

**Remark 2.4.** *If we take  $\alpha = \beta = 1$  in Corollary 2.3, then we*



have the following inequality

$$\begin{aligned} & |(x-a)(y-c)f(a,c) + (x-a)(d-y)f(a,d) \\ & + (b-x)(y-c)f(b,c) + (b-x)(d-y)f(b,d) \\ & - (y-c) \int_a^b f(t,c)dt - (d-y) \int_a^b f(t,d)dt \\ & - (x-a) \int_c^d f(a,s)ds - (b-x) \int_c^d f(b,s)ds \\ & + \int_a^b \int_c^d f(t,s) dt ds \Big| \\ \leq & \frac{[(x-a)^2 + (b-x)^2][(y-c)^2 + (d-y)^2]}{4} \|f_{ts}\|_{\Delta, \infty}. \end{aligned}$$

**Corollary 2.5.** If we choose  $g(t) = \ln t, t \in [a, b]$  and  $w(s) = \ln s, s \in [c, d]$  in Theorem 2.2, then we have the following Ostrowski type inequalities for Hadamard fractional integrals:

$$\begin{aligned} & \left| A - N_{\ln}^\beta(c, y) \left[ \mathbf{J}_{a^+}^\alpha f(x, c) + \mathbf{J}_{b^-}^\alpha f(x, c) \right] \right. \\ & - N_{\ln}^\beta(d, y) \left[ \mathbf{J}_{a^+}^\alpha f(x, d) + \mathbf{J}_{b^-}^\alpha f(x, d) \right] \\ & - M_{\ln}^\alpha(a, x) \left[ \mathbf{J}_{c^+}^\beta f(a, y) + \mathbf{J}_{d^-}^\beta f(a, y) \right] \\ & - M_{\ln}^\alpha(b, x) \left[ \mathbf{J}_{c^+}^\beta f(b, y) + \mathbf{J}_{d^-}^\beta f(b, y) \right] \\ & + \mathbf{J}_{a^+, c^+}^{\alpha, \beta} f(x, y) + \mathbf{J}_{a^+, d^-}^{\alpha, \beta} f(x, y) \\ & \left. + \mathbf{J}_{b^-, c^+}^{\alpha, \beta} f(x, y) + \mathbf{J}_{b^-, d^-}^{\alpha, \beta} f(x, y) \right| \\ \leq & \left[ M_{\ln}^{\alpha+1}(a, x) + M_{\ln}^{\beta+1}(b, x) \right] \\ & \times \left[ N_{\ln}^{\alpha+1}(c, y) + N_{\ln}^{\beta+1}(d, y) \right] \left\| \frac{f_{ts}}{g'w'} \right\|_{[a, b] \times [c, d], \infty}. \end{aligned}$$

**Theorem 2.6.** Under assumptions of Lemma 2.1, if the function  $\frac{f_{ts}}{g'w'} \in L_p(\Delta)$ , then we have the following Ostrowski type

inequality for generalized fractional integrals:

$$\begin{aligned} & \left| A - N_w^\beta(c, y) \left[ \mathbf{J}_{a^+; g}^\alpha f(x, c) + \mathbf{J}_{b^-; g}^\alpha f(x, c) \right] \right. \\ & - N_w^\beta(d, y) \left[ \mathbf{J}_{a^+; g}^\alpha f(x, d) + \mathbf{J}_{b^-; g}^\alpha f(x, d) \right] \\ & - M_g^\alpha(a, x) \left[ \mathbf{J}_{c^+; w}^\beta f(a, y) + \mathbf{J}_{d^-; w}^\beta f(a, y) \right] \\ & - M_g^\alpha(b, x) \left[ \mathbf{J}_{c^+; w}^\beta f(b, y) + \mathbf{J}_{d^-; w}^\beta f(b, y) \right] \\ & + \mathcal{J}_{a^+, c^+; g, w}^{\alpha, \beta} f(x, y) + \mathcal{J}_{a^+, d^-; g, w}^{\alpha, \beta} f(x, y) \\ & \left. + \mathcal{J}_{b^-, c^+; g, w}^{\alpha, \beta} f(x, y) + \mathcal{J}_{b^-, d^-; g, w}^{\alpha, \beta} f(x, y) \right| \\ \leq & \frac{(\alpha+1)(\beta+1)}{(\alpha q+1)^{\frac{1}{q}}(\beta q+1)^{\frac{1}{q}}} \left[ M_g^{\alpha+1}(a, x) + M_g^{\alpha+1}(b, x) \right] \\ & \times \left[ N_w^{\beta+1}(c, y) + N_w^{\beta+1}(d, y) \right] \left\| \frac{f_{ts}}{g'w'} \right\|_{[a, b] \times [c, d], p} \end{aligned}$$

where  $\frac{1}{p} + \frac{1}{q} = 1$  and

$$\left\| \frac{f_{ts}}{g'w'} \right\|_{\Delta, p} = \left( \int_a^b \int_c^d \left| \frac{f_{ts}(t, s)}{g'(t)w'(s)} \right|^p ds dt \right)^{\frac{1}{p}} < +\infty.$$

*Proof.* By the Hölder's inequality and properties of the modulus, we have

$$\begin{aligned} & |I_1| \tag{2.11} \\ \leq & \int_a^x \int_c^y (g(x) - g(t))^\alpha (w(y) - w(s))^\beta |f_{ts}(t, s)| ds dt \\ \leq & \int_a^x \int_c^y (g(x) - g(t))^\alpha (w(y) - w(s))^\beta \\ & \times \left| \frac{f_{ts}(t, s)}{g'(t)w'(s)} \right| (g'(t))^{\frac{1}{p} + \frac{1}{q}} (w'(s))^{\frac{1}{p} + \frac{1}{q}} ds dt \\ \leq & \left( \int_a^x \int_c^y \left| \frac{f_{ts}(t, s)}{g'(t)w'(s)} \right|^p g'(t)w'(s) ds dt \right)^{\frac{1}{p}} \\ & \times \left( \int_a^x \int_c^y (g(x) - g(t))^{\alpha q} (w(y) - w(s))^{\beta q} g'(t)w'(s) ds dt \right)^{\frac{1}{q}} \\ \leq & \left\| \frac{f_{ts}}{g'w'} \right\|_{[a, x] \times [c, y], p} \\ & \times \left( \int_a^x \int_c^y g'(t)w'(s) ds dt \right)^{\frac{1}{p}} \end{aligned}$$



establish

$$\begin{aligned} & \times \left( \int_a^x (g(x) - g(t))^{\alpha q} g'(t) dt \right)^{\frac{1}{q}} \\ & \times \left( \int_c^y (w(y) - w(s))^{\beta q} w'(s) ds \right)^{\frac{1}{q}} \\ = & (w(y) - w(c))^{\frac{1}{p}} (g(x) - g(a))^{\frac{1}{p}} \\ & \times \left( \frac{(g(x) - g(a))^{\alpha q + 1}}{\alpha q + 1} \right)^{\frac{1}{q}} \\ & \times \left( \frac{(w(y) - w(c))^{\beta q + 1}}{\beta q + 1} \right)^{\frac{1}{q}} \left\| \frac{f_{ts}}{g'w'} \right\|_{[a,x] \times [c,y],p} \\ = & \left\| \frac{f_{ts}}{g'w'} \right\|_{[a,x] \times [c,y],p} \\ & \times \frac{(g(x) - g(a))^{\alpha + 1} (w(y) - w(c))^{\beta + 1}}{(\alpha q + 1)^{\frac{1}{q}} (\beta q + 1)^{\frac{1}{q}}}. \end{aligned}$$

Similarly, we get

$$\begin{aligned} & |I_2| \tag{2.12} \\ \leq & \left\| \frac{f_{ts}}{g'w'} \right\|_{[a,x] \times [y,d],p} \\ & \times \frac{(g(x) - g(a))^{\alpha + 1} (w(d) - w(y))^{\beta + 1}}{(\alpha q + 1)^{\frac{1}{q}} (\beta q + 1)^{\frac{1}{q}}}, \end{aligned}$$

$$\begin{aligned} & |I_3| \tag{2.13} \\ \leq & \left\| \frac{f_{ts}}{g'w'} \right\|_{[x,b] \times [c,y],p} \\ & \times \frac{(g(b) - g(x))^{\alpha + 1} (w(y) - w(c))^{\beta + 1}}{(\alpha q + 1)^{\frac{1}{q}} (\beta q + 1)^{\frac{1}{q}}} \end{aligned}$$

and

$$\begin{aligned} & |I_4| \tag{2.14} \\ \leq & \left\| \frac{f_{ts}}{g'w'} \right\|_{[x,b] \times [y,d],p} \\ & \times \frac{(g(b) - g(x))^{\alpha + 1} (w(d) - w(y))^{\beta + 1}}{(\alpha q + 1)^{\frac{1}{q}} (\beta q + 1)^{\frac{1}{q}}}. \end{aligned}$$

By substituting the inequalities (2.11)–(2.14) in (2.5), we

$$\begin{aligned} & \left| A - N_w^\beta(c, y) \left[ J_{a+;g}^\alpha f(x, c) + J_{b-;g}^\alpha f(x, c) \right] \right. \\ & - N_w^\beta(d, y) \left[ J_{a+;g}^\alpha f(x, d) + J_{b-;g}^\alpha f(x, d) \right] \\ & - M_g^\alpha(a, x) \left[ J_{c+;w}^\beta f(a, y) + J_{d-;w}^\beta f(a, y) \right] \\ & - M_g^\alpha(b, x) \left[ J_{c+;w}^\beta f(b, y) + J_{d-;w}^\beta f(b, y) \right] \\ & + \mathcal{J}_{a+,c+;g,w}^{\alpha,\beta} f(x, y) + \mathcal{J}_{a+,d-;g,w}^{\alpha,\beta} f(x, y) \\ & + \mathcal{J}_{b-,c+;g,w}^{\alpha,\beta} f(x, y) + \mathcal{J}_{b-,d-;g,w}^{\alpha,\beta} f(x, y) \left. \right| \\ \leq & \frac{1}{\Gamma(\alpha + 1)\Gamma(\beta + 1)} \\ & \times \left[ \left\| \frac{f_{ts}}{g'w'} \right\|_{[a,x] \times [c,y],p} \right. \\ & \times \frac{(g(x) - g(a))^{\alpha + 1} (w(y) - w(c))^{\beta + 1}}{(\alpha q + 1)^{\frac{1}{q}} (\beta q + 1)^{\frac{1}{q}}} \\ & + \left\| \frac{f_{ts}}{g'w'} \right\|_{[a,x] \times [y,d],p} \\ & \times \frac{(g(x) - g(a))^{\alpha + 1} (w(d) - w(y))^{\beta + 1}}{(\alpha q + 1)^{\frac{1}{q}} (\beta q + 1)^{\frac{1}{q}}} \\ & + \left\| \frac{f_{ts}}{g'w'} \right\|_{[x,b] \times [c,y],p} \\ & \times \frac{(g(b) - g(x))^{\alpha + 1} (w(y) - w(c))^{\beta + 1}}{(\alpha q + 1)^{\frac{1}{q}} (\beta q + 1)^{\frac{1}{q}}} \\ & + \left. \left\| \frac{f_{ts}}{g'w'} \right\|_{[x,b] \times [y,d],p} \right] \\ & \times \frac{(g(b) - g(x))^{\alpha + 1} (w(d) - w(y))^{\beta + 1}}{(\alpha q + 1)^{\frac{1}{q}} (\beta q + 1)^{\frac{1}{q}}} \\ \leq & \frac{1}{(\alpha q + 1)^{\frac{1}{q}} (\beta q + 1)^{\frac{1}{q}} \Gamma(\alpha + 1)\Gamma(\beta + 1)} \\ & \times \left\| \frac{f_{ts}}{g'w'} \right\|_{[a,b] \times [c,d],p} \\ & \times \left[ (g(x) - g(a))^{\alpha + 1} + (g(b) - g(x))^{\alpha + 1} \right] \\ & \times \left[ (w(y) - w(c))^{\beta + 1} + (w(d) - w(y))^{\beta + 1} \right]. \end{aligned}$$

This completes the proof.  $\square$

**Corollary 2.7.** *If we choose  $g(t) = t$ ,  $t \in [a, b]$  and  $w(s) = s$ ,  $s \in [c, d]$  in Theorem 2.6, then we have the following Ostrowski type inequality for Riemann–Liouville fractional integrals:*



$$\begin{aligned} & \left| A - \frac{(y-c)^\beta}{\Gamma(\beta+1)} [J_{a^+}^\alpha f(x,c) + J_{b^-}^\alpha f(x,c)] \right. \\ & - \frac{(d-y)^\beta}{\Gamma(\beta+1)} [J_{a^+}^\alpha f(x,d) + J_{b^-}^\alpha f(x,d)] \\ & - \frac{(x-a)^\alpha}{\Gamma(\alpha+1)} [J_{c^+}^\beta f(a,y) + J_{d^-}^\beta f(a,y)] \\ & - \frac{(b-x)^\alpha}{\Gamma(\alpha+1)} [J_{c^+}^\beta f(b,y) + J_{d^-}^\beta f(b,y)] \\ & \left. + J_{a^+,c^+}^{\alpha,\beta} f(x,y) + J_{a^+,d^-}^{\alpha,\beta} f(x,y) \right. \\ & \left. + J_{b^-,c^+}^{\alpha,\beta} f(x,y) + J_{b^-,d^-}^{\alpha,\beta} f(x,y) \right| \\ & \leq \left\| \frac{f_{ts}}{g'w'} \right\|_{[a,b] \times [c,d],p} \\ & \times \frac{[(x-a)^{\alpha+1} + (b-x)^{\alpha+1}]}{\Gamma(\alpha+1)\Gamma(\beta+1)} \\ & \times \frac{[(y-c)^{\beta+1} + (d-y)^{\beta+1}]}{(\alpha q + 1)^{\frac{1}{q}} (\beta q + 1)^{\frac{1}{q}}} \end{aligned}$$

**Remark 2.8.** If we take  $\alpha = \beta = 1$  in Corollary 2.7, then we have

$$\begin{aligned} & \left| f(x,y) - \frac{1}{b-a} \int_a^b f(t,y) dt - \frac{1}{d-c} \int_c^d f(x,s) ds \right. \\ & \left. + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(t,s) ds dt \right| \\ & \leq \frac{4(b-a)(d-c)}{(q+1)^{\frac{2}{q}}} \\ & \times \left[ \frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right] \left[ \frac{1}{4} + \frac{(y - \frac{c+d}{2})^2}{(d-c)^2} \right] \|f_{ts}\|_{\Delta,p} \end{aligned}$$

**Corollary 2.9.** If we choose  $g(t) = \ln t$ ,  $t \in [a, b]$  and  $w(s) = \ln s$ ,  $s \in [c, d]$  in Theorem 2.6, then we have the following

Ostrowski type inequality for Hadamard fractional integrals:

$$\begin{aligned} & \left| A - N_{\ln}^\beta(c,y) [J_{a^+}^\alpha f(x,c) + J_{b^-}^\alpha f(x,c)] \right. \\ & - N_{\ln}^\beta(d,y) [J_{a^+}^\alpha f(x,d) + J_{b^-}^\alpha f(x,d)] \\ & - M_{\ln}^\alpha(a,x) [J_{c^+}^\beta f(a,y) + J_{d^-}^\beta f(a,y)] \\ & - M_{\ln}^\alpha(b,x) [J_{c^+}^\beta f(b,y) + J_{d^-}^\beta f(b,y)] \\ & + J_{a^+,c^+}^{\alpha,\beta} f(x,y) + J_{a^+,d^-}^{\alpha,\beta} f(x,y) \\ & + J_{b^-,c^+}^{\alpha,\beta} f(x,y) + J_{b^-,d^-}^{\alpha,\beta} f(x,y) \left| \right. \\ & \leq \left\| \frac{f_{ts}}{g'w'} \right\|_{[a,b] \times [c,d],p} \\ & \times \frac{[(\ln \frac{x}{a})^{\alpha+1} + (\ln \frac{b}{x})^{\alpha+1}] [(\ln \frac{y}{c})^{\beta+1} + (\ln \frac{d}{y})^{\beta+1}]}{(\alpha q + 1)^{\frac{1}{q}} (\beta q + 1)^{\frac{1}{q}} \Gamma(\alpha+1)\Gamma(\beta+1)} \end{aligned}$$

**Remark 2.10.** If we choose  $x = \frac{a+b}{2}$  and  $y = \frac{c+d}{2}$  in Theorem 2.2 and Theorem 2.6, then we obtain some new inequalities for generalized fractional integrals.

### References

- [1] A.A. Aljinović, Montgomery identity and Ostrowski type inequalities for Riemann–Liouville fractional integral, *J. Math.*, (2014), Art. ID 503195, pp. 6.
- [2] A.O. Akdemir, Inequalities of Ostrowski’s type for  $m$  and  $(\alpha, m)$ -logarithmically convex functions via Riemann–Liouville fractional integrals, *J. Comput. Anal. Appl.*, 16(2) (2014), 375-383.
- [3] N.S. Barnett and S.S. Dragomi, An Ostrowski type inequality for double integrals and applications for cubature formulae, *Soochow J. Math.*, 27(1)(2001), 109–114.
- [4] H. Budak, M.Z. Sarikaya and E. Set, Generalized Ostrowski type inequalities for functions whose local fractional derivatives are generalized  $s$ -convex in the second sense, *J. Appl. Math. Comput. Mech.*, 15(4) (2016), 11-21.
- [5] H. Budak and P. Agarwal, On Hermite-Hadamard type inequalities for co-ordinated convex mappings utilizing generalized fractional integrals, accepted as an Book Chapter, *Fractional Differentiation and its Applications*, Springer.
- [6] S. S. Dragomir, The Ostrowski’s integral inequality for Lipschitzian mappings and applications, *Comput. Math. Appl.*, 38(11-12) (1999), 33-37.
- [7] S. S. Dragomir, The Ostrowski integral inequality for mappings of bounded variation, *Bull. Austral. Math. Soc.*, 60(3) (1999), 495-508.





- [8] S.S. Dragomir, On the Ostrowski's integral inequality for mappings with bounded variation and applications, *Math. Ineq. Appl.*, 4(1) (2001), 59–66.
- [9] S. S. Dragomir, A refinement of Ostrowski's inequality for absolutely continuous functions whose derivatives belong to  $L_\infty$  and applications, *Libertas Math.*, 22 (2002), 49-63.
- [10] S. S. Dragomir, Refinements of the Ostrowski inequality in terms of the cumulative variation and applications, *Analysis* (Berlin), 34(2)(2014), 223-240.
- [11] S. S. Dragomir, Ostrowski type inequalities for Riemann–Liouville fractional integrals of bounded variation, Hölder and Lipschitzian functions, *RGMA Res. Rep. Coll.*, 20(2017), Art. 48.
- [12] S. S. Dragomir, Ostrowski and trapezoid type inequalities for Riemann–Liouville fractional integrals of absolutely continuous functions with bounded derivatives, *RGMA Res. Rep. Coll.*, 20 (2017), Art 53.
- [13] S. S. Dragomir, Ostrowski type inequalities for generalized Riemann–Liouville fractional integrals of functions with bounded variation, *RGMA Res. Rep. Coll.*, 20 (2017), Art 58.
- [14] S. S. Dragomir, On some trapezoid type inequalities for generalized Riemann–Liouville fractional integrals, *RGMA Res. Rep. Coll.*, 20 (2017), Art 68, pp. 12.
- [15] S. S. Dragomir, On some Ostrowski type inequalities for generalized Riemann–Liouville fractional integrals, *RGMA Res. Rep. Coll.*, 20 (2017), Art 67, pp. 13.
- [16] S. Erden and M. Z. Sarikaya, On the Hermite–Hadamard type and Ostrowski type inequalities for the co-ordinated convex functions, *Palestine Journal of Mathematics*, 6(1)(2017), 257-270.
- [17] A. A. Kilbas, H. M. Srivastava and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, North-Holland Mathematics Studies, 204, Elsevier Sci. B.V., Amsterdam, 2006.
- [18] A. G.-Lakoud and F. Aissaoui, New fractional inequalities of Ostrowski type, *Transylv. J. Math. Mech.*, 5(2)(2013), 103-106.
- [19] M. A. Latif, S. Hussain and S. S. Dragomir, New Ostrowski type inequalities for co-ordinated convex functions, *Transylv. J. Math. Mech.*, 4(2)(2012), 125–136.
- [20] M. A. Latif, S.S. Dragomir and A. E. Matouk, New inequalities of Ostrowski type for co-ordinated convex functions via fractional integrals, *J. Fract. Calc. Appl.*, 2(1)(2012), 1–10.
- [21] M. A. Latif and S.S. Dragomir, New Ostrowski type inequalities for co-ordinated  $s$ -convex functions in the second sense, *Le Matematiche*, 67(1)(2012), 57-72.
- [22] M. Mudassar and N. Siddiqui and M. Iqbal, On co-ordinated Ostrowski and Hadamard's type inequalities for convex functions II, *TJMM*, 9(1), (2017), 35-42.
- [23] A. Ostrowski, Über die Absolutabweichung einer differentierbaren funktion von ihrem integralmittelwert, *Comment. Math. Helv.*, 10 (1938), 226–227.
- [24] M. Z. Sarikaya and H. Filiz, Note on the Ostrowski type inequalities for fractional integrals, *Vietnam J. Math.*, 42(2) (2014), 187-190.
- [25] M. Z. Sarikaya and M. K. Yildiz, Generalization and improvement of Ostrowski type inequalities, *AIP Conference Proceedings*, 1991, 020031 (2018); doi: 10.1063/1.5047904.
- [26] M. Z. Sarikaya, H. Budak, H. Yaldiz, Some new Ostrowski type inequalities for co-ordinated convex functions, *Turkish Journal of Analysis and Number Theory*, 2(5)(2014), 176-182.

\*\*\*\*\*  
 ISSN(P):2319 – 3786  
 Malaya Journal of Matematik  
 ISSN(O):2321 – 5666  
 \*\*\*\*\*

