



# Stability of system of additive functional equations in various Banach spaces: Classical Hyers methods

M. Arunkumar<sup>1\*</sup>, E. Sathya<sup>2</sup>, S. Karthikeyan<sup>3</sup>, G. Ganapathy<sup>4</sup>, T. Namachivayam<sup>5</sup>

## Abstract

In this paper, authors proved the generalized Ulam - Hyers stability of system of additive functional equations

$$f\left(\sum_{a=1}^n a x_a\right) = \sum_{a=1}^n (a f(x_a)); \quad n \geq 1$$

$$g\left(\sum_{a=1}^n 2a y_{2a}\right) = \sum_{a=1}^n (2a g(y_{2a})); \quad n \geq 1$$

$$h\left(\sum_{a=1}^n (2a-1) z_{2a-1}\right) = \sum_{a=1}^n ((2a-1) h(z_{2a-1})); \quad n \geq 1$$

where  $n$  is a positive integer, which is originating from sum of first  $n$ , natural numbers, even natural numbers and odd natural numbers, respectively in various Banach spaces.

## Keywords

Additive functional equations, Generalized Ulam - Hyers stability, Banach spaces, 2-Banach space, Quasi 2 - Banach space, Quasi - Beta - 2- Banach space, Fuzzy Quasi - Beta - 2 - Banach space, Random Quasi - Beta - 2 - Banach space.

## AMS Subject Classification

39B52, 32B72, 32B82.

<sup>1,2,5</sup> Department of Mathematics, Government Arts College, Tiruvannamalai - 606 603, TamilNadu, India.

<sup>3</sup> Department of Mathematics, R.M.K. Engineering College, Kavaraipettai - 601 206, TamilNadu, India.

<sup>4</sup> Department of Mathematics, R.M.D. Engineering College, Kavaraipettai - 601 206, Tamil Nadu, India.

\*Corresponding author: <sup>1</sup> annarun2002@gmail.co.in; <sup>2</sup> sathya24mathematics@gmail.com; <sup>3</sup> karthik.sma204@yahoo.com; <sup>4</sup> ganagandhi@yahoo.co.in; <sup>5</sup> namachi.siva@rediffmail.com

Article History: Received 01 December 2017; Accepted 27 December 2017

©2017 MJM.

## Contents

1	Introduction .....	91
2	General Solution.....	94
3	Stability Results In Banach Space .....	94
4	Stability Results In 2- Banach Space .....	98
5	Stability Results In Quasi 2-Banach Space .....	99
6	Stability Results In Quasi-Beta-2-Banach Space .	100
7	Stability Results In Fuzzy Quasi-Beta-2-Banach Space	103
8	Stability Results In Random Quasi-Beta-2-Banach Space	108
	References .....	111

## 1. Introduction

The stability of functional equations is a burning theme that has been dealt in the last seven decades. In 1940, S.M. Ulam [40], gave a spacious collection of talk before a Mathematical Colloquium at the University of Wisconsin in which he discussed the number of significant unsolved problems. One of them is the initial spot of a new line of investigation, the Stability Problem.

**Ulam Problem :** Let  $G$  be a group and  $H$  be a metric group with metric  $d(\cdot, \cdot)$ . Given  $\epsilon > 0$  does there exist a  $\delta > 0$  such that if a function  $f : G \rightarrow H$  satisfies the inequality

$$d(f(xy), f(x)f(y)) < \delta$$

for all  $x, y \in G$ , then there exists a homomorphism  $a : G \rightarrow H$  with

$$d(f(x), a(x)) < \varepsilon$$

for all  $x \in G$ ?

For the case where the answer is affirmative, the functional equation for homomorphisms will be called stable.

The first result pertaining to the stability of functional equations was presented by D.H.Hyers [19] in 1941. He has comprehensively answered the question of Ulam by assuming  $G$  and  $H$  are Banach Spaces. He proved the following celebrated theorem.

**Theorem 1.1.** [19] Let  $X, Y$  be Banach spaces and let  $f : X \rightarrow Y$  be a mapping satisfying

$$\|f(x+y) - f(x) - f(y)\| \leq \varepsilon \tag{1.1}$$

for all  $x, y \in X$ . Then the limit

$$a(x) = \lim_{n \rightarrow \infty} \frac{f(2^n x)}{2^n} \tag{1.2}$$

exists for all  $x \in X$  and  $a : X \rightarrow Y$  is the unique additive mapping satisfying

$$\|f(x) - a(x)\| \leq \varepsilon \tag{1.3}$$

for all  $x \in X$ . Moreover, if  $f(tx)$  is continuous in  $t$  for each fixed  $x \in X$ , then the function  $a$  is linear.

This pioneer result can be expressed as “Cauchy functional equation and it is stable for any pair of Banach spaces”. The method which was provided by Hyers and which produces the additive function  $a(x)$  will be called a **direct method**. This method is the most important and most powerful tool for studying the stability of various functional equations.

In 1951, T. Aoki [3] generalized the Hyers theorem for approximately linear transformation in Banach spaces, by weakening the condition for the Cauchy difference for sum of powers of norms. Then Th.M. Rassias [32] in 1978, investigated a similar case (see L. Maligranda [29]). Both proved the following Hyers-Ulam-Aoki-Rassias theorem for the “sum”.

**Theorem 1.2.** [3, 32] Let  $X$  and  $Y$  be two Banach spaces. Let  $\theta \in [0, \infty)$  and  $p \in [0, 1)$ . If a function  $f : X \rightarrow Y$  satisfies the inequality

$$\|f(x+y) - f(x) - f(y)\| \leq \theta (\|x\|^p + \|y\|^p) \tag{1.4}$$

for all  $x, y \in X$ , then there exists a unique additive mapping  $T : X \rightarrow Y$  such that

$$\|f(x) - T(x)\| \leq \frac{2\theta}{2-2^p} \|x\|^p \tag{1.5}$$

for all  $x \in X$ . Moreover, if  $f(tx)$  is continuous in  $t$  for each fixed  $x \in X$ , then the function  $T$  is linear.

This result is known as the Modified Hyers - Ulam Stability or Generalized Hyers- Ulam Stability for the additive functional equation.

In 1982-84, J.M. Rassias [31] replaced the sum by the product of powers of norms. Infact, he proved the following theorem.

**Theorem 1.3.** [31] Let  $f : E \rightarrow E'$  be a mapping from a normed vector space  $E$  into a Banach space  $E'$  subject to the inequality

$$\|f(x+y) - f(x) - f(y)\| \leq \varepsilon \|x\|^p \|y\|^p \tag{1.6}$$

for all  $x, y \in E$ , where  $\varepsilon$  and  $p$  are constants with  $\varepsilon > 0$  and  $0 \leq p < \frac{1}{2}$ . Then the limit

$$L(x) = \lim_{n \rightarrow \infty} \frac{f(2^n x)}{2^n} \tag{1.7}$$

exists for all  $x \in E$  and  $L : E \rightarrow E'$  is the unique additive mapping which satisfies

$$\|f(x) - L(x)\| \leq \frac{\varepsilon}{2-2^{2p}} \|x\|^{2p} \tag{1.8}$$

for all  $x \in E$ . If  $p < 0$ , then the inequality (1.6) holds for  $x, y \neq 0$  and (1.8) for  $x \neq 0$ .

If  $p > \frac{1}{2}$  the inequality (1.6) holds for  $x, y \in E$  and the limit

$$A(x) = \lim_{n \rightarrow \infty} 2^n f\left(\frac{x}{2^n}\right) \tag{1.9}$$

exists for all  $x \in E$  and  $A : E \rightarrow E'$  is the unique additive mapping which satisfies

$$\|f(x) - A(x)\| \leq \frac{\varepsilon}{2^{2p}-2} \|x\|^{2p} \tag{1.10}$$

for all  $x \in E$ . If in addition  $f : E \rightarrow E'$  is a mapping such that the transformation  $t \rightarrow f(tx)$  is continuous in  $t \in \mathbb{R}$  for each fixed  $x \in E$ , then  $L$  is  $\mathbb{R}$ - linear mapping.

In 1994, P. Gavruta [18] generalized all the above mentioned results by considering the control function as a function of variables and proved the following theorem.

**Theorem 1.4.** [18] Let  $E$  be an abelian group,  $F$  be a Banach space and let  $\phi : E \times E \rightarrow [0, \infty)$  be a function satisfying

$$\Phi(x, y) = \sum_{k=0}^{+\infty} \frac{1}{2^{k+1}} \phi\left(2^k x, 2^k y\right) < +\infty \tag{1.11}$$

for all  $x, y \in E$ . If a function  $f : E \rightarrow F$  satisfies the functional inequality

$$\|f(x+y) - f(x) - f(y)\| \leq \phi(x, y) \tag{1.12}$$

for all  $x, y \in E$ . Then there exists a unique additive mapping  $T : E \rightarrow F$  which satisfies

$$\|f(x) - T(x)\| \leq \Phi(x, y) \tag{1.13}$$

for all  $x \in E$ . If moreover  $f(tx)$  is continuous in  $t$  for fixed  $x \in E$ , then  $T$  is linear.



This stability property is called Generalized Hyers - Ulam - Rassias Stability of functional equation.

In 2008, a special case of Gavruta’s theorem for the unbounded Cauchy difference was obtained by K. Ravi, M. Arunkumar and J.M. Rassias [34] by considering the summation of both the sum and the product of two  $p$ -norms in the spirit of Rassias approach is called J.M. Rassias Stability of functional equation.

**Theorem 1.5.** [34] Let  $(E, \perp)$  denote an orthogonality normed space with norm  $\|\cdot\|_E$  and  $(F, \|\cdot\|_F)$  is a Banach space and  $f : E \rightarrow F$  be a mapping which satisfying the inequality

$$\begin{aligned} & \|f(mx + y) + f(mx - y) - 2f(x + y) - 2f(x - y) \\ & - 2(m^2 - 2)f(x) + 2f(y)\|_F \\ & \leq \varepsilon \left\{ \|x\|_E^p \|y\|_E^p + \left( \|x\|_E^{2p} + \|y\|_E^{2p} \right) \right\} \end{aligned} \quad (1.14)$$

for all  $x, y \in E$  with  $x \perp y$ , where  $\varepsilon$  and  $p$  are constants with  $\varepsilon, p > 0$  and either  $m > 1; p < 1$  or  $m < 1; p > 1$  with  $m \neq 0; m \neq \pm 1; m \neq \pm\sqrt{2}$  and  $-1 \neq |m|^{p-1} < 1$ .

Then the limit

$$Q(x) = \lim_{n \rightarrow \infty} \frac{f(m^n x)}{m^{2n}} \quad (1.15)$$

exists for all  $x \in E$  and  $Q : E \rightarrow F$  is the unique orthogonally Euler-Lagrange quadratic mapping such that

$$\|f(x) - Q(x)\|_F \leq \frac{\varepsilon}{2|m^2 - m^{2p}|} \|x\|_E^{2p} \quad (1.16)$$

for all  $x \in E$ .

A number is a mathematical object used to count, measure, and label. The most familiar numbers are the natural numbers (sometimes called whole numbers or counting numbers): 1, 2, 3, and so on. An even number is an integer that is *evenly divisible* by two, that is divisible by two without remainder; an odd number is an integer that is not even. (The old-fashioned term *evenly divisible* is now almost always shortened to *divisible*.) Equivalently, another way of defining an odd number is that it is an integer of the form  $n = 2k - 1$ , where  $k$  is an integer, and an even number has the form  $n = 2k$ , where  $k$  is an integer.

The sum of first  $n$  natural numbers, sum of first  $n$  even natural numbers and sum of first  $n$  odd natural numbers, are

$$1 + 2 + 3 + 4 + \dots + N = \frac{N(N + 1)}{2}$$

$$2 + 4 + 6 + 8 + \dots + 2N = N(N + 1)$$

$$1 + 3 + 5 + 7 + \dots + (2N - 1) = N^2$$

The above sum of observation can be taken as a functional

equation of the following forms

$$\begin{aligned} & f(x_1 + 2x_2 + 3x_3 + 4x_4 + \dots + nx_n) \\ & = 1f(x_1) + 2f(x_2) + 3f(x_3) + 4f(x_4) + \dots + nf(x_n) \\ & g(2y_2 + 4y_4 + 6y_6 + 8y_8 + \dots + 2ny_{2n}) \\ & = 2g(y_2) + 4g(y_4) + 6g(y_6) + 8g(y_8) + \dots + 2ng(y_{2n}) \\ & h(1z_1 + 3z_3 + 5z_5 + 7z_7 + \dots + (2n - 1)z_{2n-1}) \\ & = 1h(z_1) + 3h(z_3) + 5h(z_5) + 7h(z_7) + \dots + (2n - 1)h(z_{2n-1}) \end{aligned}$$

One of the most famous functional equations is the additive functional equation

$$f(x + y) = f(x) + f(y) \quad (1.17)$$

In 1821, it was first solved by A.L. Cauchy in the class of continuous real-valued functions. It is often called an additive Cauchy functional equation in tribute of A.L. Cauchy. The theory of additive functional equations is commonly useful to the growth of theories of further functional equations. Moreover, the properties of additive functional equations are powerful tackle in almost every field of natural and social sciences. Every solution of the additive functional equation (1.17) is called an additive function.

The solution and stability of various additive functional equations

$$f(2x - y) + f(x - 2y) = 3f(x) - 3f(y), \quad (1.18)$$

$$f(x + y - 2z) + f(2x + 2y - z) = 3f(x) + 3f(y) - 3f(z), \quad (1.19)$$

$$\begin{aligned} & f(m(x + y) - 2mz) + f(2m(x + y) - mz) \\ & = 3m[f(x) + f(y) - f(z)] \quad m \geq 1, \end{aligned} \quad (1.20)$$

$$\begin{aligned} & f\left(a \sum_{i=1}^{n-1} x_i - 2ax_n\right) + f\left(2a \sum_{i=1}^{n-1} x_i - ax_n\right) \\ & = 3a \left( \sum_{i=1}^{n-1} f(x_i) - f(x_n) \right) \quad n \geq 3, \end{aligned} \quad (1.21)$$

$$f(2x \pm y \pm z) = f(x \pm y) + f(x \pm z) \quad (1.22)$$

$$f(x) + f(x) = f(2x) \quad (1.23)$$

$$f(y) = \frac{f(y + z) + f(y - z)}{2} \quad (1.24)$$

$$f(x) = \sum_{\ell=1}^n \left( \frac{f(x + \ell y_\ell) + f(x - \ell y_\ell)}{2\ell} \right) \quad (1.25)$$

$$f\left(nx_0 \pm \sum_{i=1}^n x_i\right) = \sum_{i=1}^n f(x_0 \pm x_i) \quad (1.26)$$

were discussed by D.O. Lee [27], K. Ravi, M. Arunkumar [33], M. Arunkumar [4, 5, 7, 8, 11].

In this paper, authors proved the generalized Ulam - Hyers



stability of system of additive functional equations

$$f\left(\sum_{a=1}^n a x_a\right) = \sum_{a=1}^n (a f(x_a)), \quad n \geq 1 \tag{1.27}$$

$$g\left(\sum_{a=1}^n 2a y_{2a}\right) = \sum_{a=1}^n (2a g(y_{2a})), \quad n \geq 1 \tag{1.28}$$

$$h\left(\sum_{a=1}^n (2a-1) z_{2a-1}\right) = \sum_{a=1}^n ((2a-1) h(z_{2a-1})), \quad n \geq 1 \tag{1.29}$$

where  $n$  is a positive integer, which is originating from, sum of first  $n$  natural numbers, sum of first  $n$  even natural numbers and sum of first  $n$  odd natural numbers in various Banach spaces and having solutions

$$f(x) = b_1 x; \tag{1.30}$$

$$g(y) = b_2 y; \tag{1.31}$$

$$h(z) = b_3 z \tag{1.32}$$

where  $b_1, b_2, b_3$  are constants.

In this paper, authors proved the generalized Ulam - Hyers stability of system of additive functional equations where  $n$  is a positive integer, which is originating from sum of first  $n$ , natural numbers, even natural numbers and odd natural numbers, respectively in various Banach spaces.

## 2. General Solution

In this section, we give the general solution of the functional equations (1.27), (1.28) and (1.29). To prove the general solution, we let us take  $A$  and  $B$  be real vector spaces.

The proof of the following theorems are proved by the additive property of functions. Hence the details of the proof are omitted.

**Theorem 2.1.** *An additive function  $f : A \rightarrow B$  satisfies the functional equation (1.17) for all  $x, y \in A$  if and only if  $f : A \rightarrow B$  satisfies the functional equation (1.27) for all  $x_1, x_2, \dots, x_n \in A$ .*

**Theorem 2.2.** *An additive function  $g : A \rightarrow B$  satisfies the functional equation*

$$g(x+y) = g(x) + g(y) \tag{2.1}$$

*for all  $x, y \in A$  if and only if  $g : A \rightarrow B$  satisfies the functional equation (1.28) for all  $y_2, y_4, \dots, y_{2n} \in A$ .*

**Theorem 2.3.** *An additive function  $h : A \rightarrow B$  satisfies the functional equation*

$$h(x+y) = h(x) + h(y) \tag{2.2}$$

*for all  $x, y \in A$  if and only if  $h : A \rightarrow B$  satisfies the functional equation (1.29) for all  $z_1, z_3, \dots, z_{2n-1} \in A$ .*

## 3. Stability Results In Banach Space

In this section, we investigate the generalized Ulam - Hyers stability of the functional equations (1.27), (1.28) and (1.29) in Banach spaces using Hyers Method.

Now, we give basic definitions and notations in Banach Space.

**Definition 3.1.** *Let  $X$  be a set. A function  $\|\cdot\| : X \times X \rightarrow [0, \infty)$  is called a **Normed Linear space** on  $X$  if  $\|\cdot\|$  satisfies the following conditions*

- (i)  $\|x\| = 0$  if and only if  $x = 0$  for all  $x \in X$ ;
- (ii)  $\|x, y\| = 0$  if and only if  $x = y$  for all  $x, y \in X$ ;
- (iii)  $\|\lambda x\| = |\lambda| \|x\|$  for all  $x \in X$  and  $\lambda \in \mathbb{R}$ ;
- (iv)  $\|x + y\| \leq \|x\| + \|y\|$  for all  $x, y \in X$ ;
- (v)  $\|x, z\| \leq \|x, y\| + \|y, z\|$  for all  $x, y, z \in X$ ;

**Definition 3.2.** *A sequence  $\{x_n\}$  in a normed linear space  $X$  is called a **convergent** sequence if there is an  $x \in X$  such that  $\lim_{n \rightarrow \infty} \|x_n - x\| = 0$*

**Definition 3.3.** *A sequence  $\{x_n\}$  in a normed linear space  $X$  is called a **Cauchy sequence** if there exists a point,  $x_m \in X$  such that  $\lim_{n, m \rightarrow \infty} \|x_n - x_m\| = 0$*

**Definition 3.4.** *A normed linear space is said to be **Complete** if every Cauchy sequence converges.*

**Definition 3.5.** *A complete normed linear space is said to be a **Banach Space***

To prove stability results, let us consider  $\mathcal{A}$  and  $\mathcal{B}$  be Banach spaces.

**Theorem 3.6.** *Let  $f, g, h : \mathcal{A} \rightarrow \mathcal{B}$  be a mapping satisfying the following inequalities*

$$\left\| f\left(\sum_{a=1}^n a x_a\right) - \sum_{a=1}^n (a f(x_a)) \right\| \leq N(x_1, x_2, \dots, x_n); \tag{3.1}$$

$$\left\| g\left(\sum_{a=1}^n 2a y_{2a}\right) - \sum_{a=1}^n (2a g(y_{2a})) \right\| \leq E(y_2, y_4, \dots, y_{2n}); \tag{3.2}$$

$$\left\| h\left(\sum_{a=1}^n (2a-1) z_{2a-1}\right) - \sum_{a=1}^n ((2a-1) h(z_{2a-1})) \right\| \leq O(z_1, z_3, \dots, z_{2n-1}); \tag{3.3}$$

*for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$ , where  $N : \mathcal{A}^n \rightarrow [0, \infty)$ ,  $E : \mathcal{A}^{2n} \rightarrow [0, \infty)$  and  $O : \mathcal{A}^{2n-1} \rightarrow [0, \infty)$  satisfying the conditions*

$$\lim_{\alpha \rightarrow \infty} \frac{N(\kappa^{\alpha\gamma} x_1, \kappa^{\alpha\gamma} x_2, \dots, \kappa^{\alpha\gamma} x_n)}{\kappa^{\alpha\gamma}} = 0; \tag{3.4}$$

$$\lim_{\alpha \rightarrow \infty} \frac{E(\rho^{\alpha\gamma} y_2, \rho^{\alpha\gamma} y_4, \dots, \rho^{\alpha\gamma} y_{2n})}{\rho^{\alpha\gamma}} = 0; \tag{3.5}$$

$$\lim_{\alpha \rightarrow \infty} \frac{O(\tau^{\alpha\gamma} z_1, \tau^{\alpha\gamma} z_3, \dots, \tau^{\alpha\gamma} z_{2n-1})}{\tau^{\alpha\gamma}} = 0; \tag{3.6}$$



for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$  with  $\gamma = \pm 1$ . Then there exists one and only additive mapping  $A_f, A_g, A_h: \mathcal{A} \rightarrow \mathcal{B}$  satisfying (1.27), (1.28), (1.29), and

$$\|A_f(x) - f(x)\| \leq \frac{1}{\kappa} \sum_{\beta=\frac{1-\gamma}{2}}^{\infty} \frac{N(\kappa^{\beta\gamma}x, \kappa^{\beta\gamma}x, \dots, \kappa^{\beta\gamma}x)}{\kappa^{\beta\gamma}}; \tag{3.7}$$

$$\|A_g(y) - g(y)\| \leq \frac{1}{\rho} \sum_{\beta=\frac{1-\gamma}{2}}^{\infty} \frac{E(\rho^{\beta\gamma}y, \rho^{\beta\gamma}y, \dots, \rho^{\beta\gamma}y)}{\rho^{\beta\gamma}}; \tag{3.8}$$

$$\|A_h(z) - h(z)\| \leq \frac{1}{\tau} \sum_{\beta=\frac{1-\gamma}{2}}^{\infty} \frac{O(\tau^{\beta\gamma}z, \tau^{\beta\gamma}z, \dots, \tau^{\beta\gamma}z)}{\tau^{\beta\gamma}}; \tag{3.9}$$

for all  $x, y, z \in \mathcal{A}$ , respectively. The mappings  $A_f, A_g, A_h$  are respectively defined as

$$A_f(x) = \lim_{\alpha \rightarrow \infty} \frac{f(\kappa^{\alpha\gamma}x)}{\kappa^{\alpha\gamma}}; \tag{3.10}$$

$$A_g(y) = \lim_{\alpha \rightarrow \infty} \frac{g(\rho^{\alpha\gamma}y)}{\rho^{\alpha\gamma}}; \tag{3.11}$$

$$A_h(z) = \lim_{\alpha \rightarrow \infty} \frac{h(\tau^{\alpha\gamma}z)}{\tau^{\alpha\gamma}}; \tag{3.12}$$

for all  $x, y, z \in \mathcal{A}$ .

*Proof.* Changing

$$(x_1, x_2, \dots, x_n) = (x, x, \dots, x) \text{ in (3.1);}$$

$$(y_2, y_4, \dots, y_{2n}) = (y, y, \dots, y) \text{ in (3.2);}$$

$$(z_1, z_3, \dots, z_{2n-1}) = (z, z, \dots, z) \text{ in (3.3);}$$

we arrive the following inequalities

$$\left\| f\left(\sum_{a=1}^n ax\right) - \left(\sum_{a=1}^n a\right) f(x) \right\| \leq N(x, x, \dots, x); \tag{3.13}$$

$$\left\| g\left(\sum_{a=1}^n 2ay\right) - \left(\sum_{a=1}^n 2a\right) g(y) \right\| \leq E(y, y, \dots, y); \tag{3.14}$$

$$\left\| h\left(\sum_{a=1}^n (2a-1)z\right) - \left(\sum_{a=1}^n (2a-1)\right) h(z) \right\| \leq O(z, z, \dots, z); \tag{3.15}$$

for all  $x, y, z \in \mathcal{A}$ . Define

$$\sum_{a=1}^n a = \frac{n(n+1)}{2} = \kappa \text{ in (3.13);}$$

$$\sum_{a=1}^n 2a = n(n+1) = \rho \text{ in (3.14);}$$

$$\sum_{a=1}^n (2a-1) = n^2 = \tau \text{ in (3.15);}$$

we obtain the succeeding inequalities

$$\|f(\kappa x) - \kappa f(x)\| \leq N(x, x, \dots, x); \tag{3.16}$$

$$\|g(\rho y) - \rho g(y)\| \leq E(y, y, \dots, y); \tag{3.17}$$

$$\|h(\tau z) - \tau h(z)\| \leq O(z, z, \dots, z); \tag{3.18}$$

for all  $x, y, z \in \mathcal{A}$ . It follows from above inequalities

$$\left\| \frac{f(\kappa x)}{\kappa} - f(x) \right\| \leq \frac{N(x, x, \dots, x)}{\kappa}; \tag{3.19}$$

$$\left\| \frac{g(\rho y)}{\rho} - g(y) \right\| \leq \frac{E(y, y, \dots, y)}{\rho}; \tag{3.20}$$

$$\left\| \frac{h(\tau z)}{\tau} - h(z) \right\| \leq \frac{O(z, z, \dots, z)}{\tau}; \tag{3.21}$$

for all  $x, y, z \in \mathcal{A}$ . Replacing

$$x = \kappa x \text{ and } \div \kappa \text{ in (3.19);}$$

$$y = \rho y \text{ and } \div \rho \text{ in (3.20);}$$

$$z = \tau z \text{ and } \div \tau \text{ in (3.21);}$$

we arrive the following inequalities

$$\left\| \frac{f(\kappa^2 x)}{\kappa^2} - \frac{f(\kappa x)}{\kappa} \right\| \leq \frac{N(\kappa x, \kappa x, \dots, \kappa x)}{\kappa^2}; \tag{3.22}$$

$$\left\| \frac{g(\rho^2 y)}{\rho^2} - \frac{g(\rho y)}{\rho} \right\| \leq \frac{E(\rho y, \rho y, \dots, \rho y)}{\rho^2}; \tag{3.23}$$

$$\left\| \frac{h(\tau^2 z)}{\tau^2} - \frac{h(\tau z)}{\tau} \right\| \leq \frac{O(\tau z, \tau z, \dots, \tau z)}{\tau^2}; \tag{3.24}$$

for all  $x, y, z \in \mathcal{A}$ . With the help of triangle inequality from

(3.19) and (3.22); (3.20) and (3.23); (3.21) and (3.24);

we achieve the subsequent inequalities

$$\begin{aligned} & \left\| \frac{f(\kappa^2 x)}{\kappa^2} - f(x) \right\| \\ & \leq \left\| \frac{f(\kappa^2 x)}{\kappa^2} - \frac{f(\kappa x)}{\kappa} \right\| + \left\| \frac{f(\kappa x)}{\kappa} - f(x) \right\| \\ & \leq \frac{1}{\kappa} \left( N(x, x, \dots, x) + \frac{N(\kappa x, \kappa x, \dots, \kappa x)}{\kappa} \right); \end{aligned} \tag{3.25}$$

$$\begin{aligned} & \left\| \frac{g(\rho^2 y)}{\rho^2} - g(y) \right\| \\ & \leq \left\| \frac{g(\rho^2 y)}{\rho^2} - \frac{g(\rho y)}{\rho} \right\| + \left\| \frac{g(\rho y)}{\rho} - g(y) \right\| \\ & \leq \frac{1}{\rho} \left( E(y, y, \dots, y) + \frac{E(\rho y, \rho y, \dots, \rho y)}{\rho} \right); \end{aligned} \tag{3.26}$$



$$\begin{aligned} & \left\| \frac{h(\tau^2 z)}{\tau^2} - h(z) \right\| \\ & \leq \left\| \frac{h(\tau^2 z)}{\tau^2} - \frac{h(\tau z)}{\tau} \right\| + \left\| \frac{h(\tau z)}{\tau} - h(z) \right\| \\ & \leq \frac{1}{\tau} \left( E(z, z, \dots, z) + \frac{E(\tau z, \tau z, \dots, \tau)}{\tau} \right); \quad (3.27) \end{aligned}$$

for all  $x, y, z \in \mathcal{A}$ . Generalizing for a positive integer  $\alpha$ , we arrive

$$\left\| \frac{f(\kappa^\alpha x)}{\kappa^\alpha} - f(x) \right\| \leq \frac{1}{\kappa} \sum_{\beta=0}^{\alpha} \frac{N(\kappa^\beta x, \kappa^\beta x, \dots, \kappa^\beta x)}{\kappa^\beta}; \quad (3.28)$$

$$\left\| \frac{g(\rho^\alpha y)}{\rho^\alpha} - g(y) \right\| \leq \frac{1}{\rho} \sum_{\beta=0}^{\alpha} \frac{E(\rho^\beta y, \rho^\beta y, \dots, \rho^\beta y)}{\rho^\beta}; \quad (3.29)$$

$$\left\| \frac{h(\tau^\alpha z)}{\tau^\alpha} - h(z) \right\| \leq \frac{1}{\tau} \sum_{\beta=0}^{\alpha} \frac{O(\tau^\beta z, \tau^\beta z, \dots, \tau^\beta z)}{\tau^\beta}; \quad (3.30)$$

for all  $x, y, z \in \mathcal{A}$ . It is easy to verify that the sequences

$$\left\{ \frac{f(\kappa^\alpha x)}{\kappa^\alpha} \right\}; \left\{ \frac{g(\rho^\alpha y)}{\rho^\alpha} \right\}; \left\{ \frac{h(\tau^\alpha z)}{\tau^\alpha} \right\}$$

are Cauchy sequences. Indeed replacing

$$\begin{aligned} x &= \kappa^\delta x \quad \text{and} \quad \div \quad \kappa^\delta \quad \text{in} \quad (3.28); \\ y &= \rho^\delta y \quad \text{and} \quad \div \quad \rho^\delta \quad \text{in} \quad (3.29); \\ z &= \tau^\delta z \quad \text{and} \quad \div \quad \tau^\delta \quad \text{in} \quad (3.30); \end{aligned}$$

we obtain

$$\begin{aligned} & \left\| \frac{f(\kappa^{\alpha+\delta} x)}{\kappa^{\alpha+\delta}} - \frac{f(\kappa^\delta x)}{\kappa^\delta} \right\| \\ & \leq \frac{1}{\kappa} \sum_{\beta=0}^{\alpha} \frac{N(\kappa^{\beta+\delta} x, \kappa^{\beta+\delta} x, \dots, \kappa^{\beta+\delta} x)}{\kappa^{\beta+\delta}} \\ & \rightarrow 0 \quad \text{as} \quad \delta \rightarrow \infty; \\ & \left\| \frac{g(\rho^{\alpha+\delta} y)}{\rho^{\alpha+\delta}} - \frac{g(\rho^\delta y)}{\rho^\delta} \right\| \\ & \leq \frac{1}{\rho} \sum_{\beta=0}^{\alpha} \frac{E(\rho^{\beta+\delta} y, \rho^{\beta+\delta} y, \dots, \rho^{\beta+\delta} y)}{\rho^{\beta+\delta}} \\ & \rightarrow 0 \quad \text{as} \quad \delta \rightarrow \infty; \\ & \left\| \frac{h(\tau^{\alpha+\delta} z)}{\tau^{\alpha+\delta}} - \frac{h(\tau^\delta z)}{\tau^\delta} \right\| \\ & \leq \frac{1}{\tau} \sum_{\beta=0}^{\alpha} \frac{O(\tau^{\beta+\delta} z, \tau^{\beta+\delta} z, \dots, \tau^{\beta+\delta} z)}{\tau^{\beta+\delta}} \\ & \rightarrow 0 \quad \text{as} \quad \delta \rightarrow \infty; \end{aligned}$$

for all  $x, y, z \in \mathcal{A}$ . This proves the existence of Cauchy sequences. Since  $\mathcal{B}$  is a Banach space, this sequences converges to a point  $A_f; A_g; A_h$  respectively and it defined by

$$\begin{aligned} A_f(x) &= \lim_{\alpha \rightarrow \infty} \frac{f(\kappa^\alpha x)}{\kappa^\alpha}; \\ A_g(y) &= \lim_{\alpha \rightarrow \infty} \frac{g(\rho^\alpha y)}{\rho^\alpha}; \\ A_h(z) &= \lim_{\alpha \rightarrow \infty} \frac{h(\tau^\alpha z)}{\tau^\alpha}; \end{aligned}$$

for all  $x, y, z \in \mathcal{A}$ . Taking limit as  $\alpha$  tends to infinity in (3.28), (3.29) and (3.30), we see that (3.7), (3.8) and (3.9) holds respectively for all  $x, y, z \in \mathcal{A}$  with  $\gamma = 1$ .

To show the mappings  $A_f(x); A_g(y); A_h(z)$  satisfies the functional equations (1.27), (1.28) and (1.29) by replacing

$$(x_1, x_2, \dots, x_n) = (\kappa^\alpha x_1, \kappa^\alpha x_2, \dots, \kappa^\alpha x_n) \quad \text{in} \quad (3.1)$$

$$(y_2, y_4, \dots, y_{2n}) = (\rho^\alpha y_2, \rho^\alpha y_4, \dots, \rho^\alpha y_{2n}) \quad \text{in} \quad (3.2)$$

$$(z_1, z_3, \dots, z_{2n-1}) = (\tau^\alpha z_1, \tau^\alpha z_3, \dots, \tau^\alpha z_{2n-1}) \quad \text{in} \quad (3.3)$$

respectively, we arrive

$$\begin{aligned} & \left\| A_f \left( \sum_{a=1}^n a x_a \right) - \sum_{a=1}^n (a A_f(x_a)) \right\| \\ & = \frac{1}{\kappa^\alpha} \left\| f \left( \sum_{a=1}^n a \kappa^\alpha x_a \right) - \sum_{a=1}^n (a f(\kappa^\alpha x_a)) \right\| \\ & \leq \frac{1}{\kappa^\alpha} N(\kappa^\alpha x_1, \kappa^\alpha x_2, \dots, \kappa^\alpha x_n) \\ & \left\| A_g \left( \sum_{a=1}^n 2a y_{2a} \right) - \sum_{a=1}^n (2a A_g(y_{2a})) \right\| \\ & = \frac{1}{\rho^\alpha} \left\| g \left( \sum_{a=1}^n 2a \rho^\alpha y_{2a} \right) - \sum_{a=1}^n (2a g(\rho^\alpha y_{2a})) \right\| \\ & \leq \frac{1}{\rho^\alpha} E(\rho^\alpha y_2, \rho^\alpha y_4, \dots, \rho^\alpha y_{2n}) \\ & \left\| A_h \left( \sum_{a=1}^n (2a-1) z_{2a-1} \right) - \sum_{a=1}^n ((2a-1) A_h(z_{2a-1})) \right\| \\ & = \frac{1}{\tau^\alpha} \left\| h \left( \sum_{a=1}^n (2a-1) \tau^\alpha z_{2a-1} \right) - \sum_{a=1}^n ((2a-1) h(\tau^\alpha z_{2a-1})) \right\| \\ & \leq \frac{1}{\tau^\alpha} O(\tau^\alpha z_1, \tau^\alpha z_3, \dots, \tau^\alpha z_{2n-1}) \end{aligned}$$

for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$ . Taking limit as  $\alpha$  tends to infinity in the above inequalities, it proves that  $A_f(x); A_g(y); A_h(z)$  satisfies the functional equations (1.27), (1.28) and (1.29) for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$ .

In order to prove the being  $A_f(x); A_g(y); A_h(z)$  are unique. Let  $A'_f(x); A'_g(y); A'_h(z)$  be another additive mappings satisfying (3.10), (3.11), (3.12) and (1.27), (1.28), (1.29) respectively.





Now

the following inequalities

$$\begin{aligned} & \|A_f(x) - A'_f(x)\| \\ & \leq \frac{1}{\kappa^\alpha} \{ \|A_f(\kappa^\alpha x) - f(\kappa^\alpha x)\| + \|A'_f(\kappa^\alpha x) - f(\kappa^\alpha x)\| \} \\ & \leq \frac{2}{\kappa} \sum_{\beta=0}^{\infty} \frac{N(\kappa^{\beta+\alpha}x, \kappa^{\beta+\alpha}x, \dots, \kappa^{\beta+\alpha}x)}{\kappa^{\beta+\alpha}} \\ & \rightarrow 0 \text{ as } \alpha \rightarrow \infty; \\ & \|A_g(y) - A'_g(y)\| \\ & \leq \frac{1}{\rho^\alpha} \{ \|A_g(\rho^\alpha y) - g(\rho^\alpha y)\| + \|A'_g(\rho^\alpha y) - g(\rho^\alpha y)\| \} \\ & \leq \frac{2}{\rho} \sum_{\beta=0}^{\infty} \frac{E(\rho^{\beta+\alpha}y, \rho^{\beta+\alpha}y, \dots, \rho^{\beta+\alpha}y)}{\rho^{\beta+\alpha}} \\ & \rightarrow 0 \text{ as } \alpha \rightarrow \infty; \\ & \|A_h(z) - A'_h(z)\| \\ & \leq \frac{1}{\tau^\alpha} \{ \|A_h(\tau^\alpha z) - h(\tau^\alpha z)\| + \|A'_h(\tau^\alpha z) - h(\tau^\alpha z)\| \} \\ & \leq \frac{2}{\tau} \sum_{\beta=0}^{\infty} \frac{O(\tau^{\beta+\alpha}z, \tau^{\beta+\alpha}z, \dots, \tau^{\beta+\alpha}z)}{\tau^{\beta+\alpha}} \\ & \rightarrow 0 \text{ as } \alpha \rightarrow \infty; \end{aligned}$$

$$\begin{aligned} & \left\| f\left(\sum_{a=1}^n a x_a\right) - \sum_{a=1}^n (a f(x_a)) \right\| \\ & \leq \begin{cases} \lambda, & \mu \neq 1; \\ \sum_{a=1}^n \lambda \|x_a\|^\mu, & n\mu \neq 1; \\ \prod_{a=1}^n \lambda \|x_a\|^\mu + \sum_{a=1}^n \lambda \|x_a\|^{n\mu}, & n\mu = 1; \end{cases} \end{aligned} \tag{3.34}$$

$$\begin{aligned} & \left\| g\left(\sum_{a=1}^n 2a y_{2a}\right) - \sum_{a=1}^n (2a g(y_{2a})) \right\| \\ & \leq \begin{cases} \lambda, & \mu \neq 1; \\ \sum_{a=1}^n \lambda \|y_{2a}\|^\mu, & n\mu \neq 1; \\ \prod_{a=1}^n \lambda \|y_{2a}\|^\mu + \sum_{a=1}^n \lambda \|y_{2a}\|^{n\mu}, & n\mu = 1; \end{cases} \end{aligned} \tag{3.35}$$

$$\begin{aligned} & \left\| h\left(\sum_{a=1}^n (2a-1) z_{2a-1}\right) - \sum_{a=1}^n ((2a-1) h(z_{2a-1})) \right\| \\ & \leq \begin{cases} \lambda, & \mu \neq 1; \\ \sum_{a=1}^n \lambda \|z_{2a-1}\|^\mu, & n\mu \neq 1; \\ \prod_{a=1}^n \lambda \|z_{2a-1}\|^\mu + \sum_{a=1}^n \lambda \|z_{2a-1}\|^{n\mu}, & n\mu = 1; \end{cases} \end{aligned} \tag{3.36}$$

for all  $x, y, z \in \mathcal{A}$ . Hence the mappings  $A_f(x); A_g(y); A_h(z)$  are unique. Thus the theorem holds for  $\gamma = 1$ .

Also, if we replace

$$x = \frac{x}{\kappa} \text{ in (3.16);}$$

$$y = \frac{y}{\rho} \text{ in (3.17);}$$

$$z = \frac{z}{\tau} \text{ in (3.18);}$$

we arrive

$$\left\| f(x) - \kappa f\left(\frac{x}{\kappa}\right) \right\| \leq N\left(\frac{x}{\kappa}, \frac{x}{\kappa}, \dots, \frac{x}{\kappa}\right); \tag{3.31}$$

$$\left\| g(y) - \rho g\left(\frac{y}{\rho}\right) \right\| \leq E\left(\frac{y}{\rho}, \frac{y}{\rho}, \dots, \frac{y}{\rho}\right); \tag{3.32}$$

$$\left\| h(z) - \tau h\left(\frac{z}{\tau}\right) \right\| \leq O\left(\frac{z}{\tau}, \frac{z}{\tau}, \dots, \frac{z}{\tau}\right); \tag{3.33}$$

for all  $x, y, z \in \mathcal{A}$ . The rest of proof is similar to that of the case  $\gamma = 1$ . Thus the theorem holds for  $\gamma = -1$  also. This completes the proof of the Theorem.  $\square$

The following corollary is an immediate consequence of Theorem 3.6 concerning the Hyers - Ulam, Hyers - Ulam - Rassias and J.M.Rassias stabilities of the functional equations (1.27), (1.28) and (1.29).

**Corollary 3.7.** Let  $f, g, h : \mathcal{A} \rightarrow \mathcal{B}$  be a mapping satisfying for all  $x, y, z \in \mathcal{A}$ , respectively.

for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$ , where  $\lambda$  and  $\mu$  are positive constants. Then there exists one and only additive mapping  $A_f, A_g, A_h : \mathcal{A} \rightarrow \mathcal{B}$  satisfying (1.27), (1.28), (1.29), and

$$\|A_f(x) - f(x)\| \leq \begin{cases} \lambda \\ \frac{|\kappa - 1|}{n\lambda \|x\|^\mu}; \\ \frac{|\kappa - \kappa^\mu|}{(n+1)\lambda \|x\|^{n\mu}}; \\ \frac{|\kappa - \kappa^{n\mu}|}{|\kappa - \kappa^{n\mu}|}; \end{cases} \tag{3.37}$$

$$\|A_g(y) - g(y)\| \leq \begin{cases} \lambda \\ \frac{|\rho - 1|}{n\lambda \|y\|^\mu}; \\ \frac{|\rho - \rho^\mu|}{(n+1)\lambda \|y\|^{n\mu}}; \\ \frac{|\rho - \rho^{n\mu}|}{|\rho - \rho^{n\mu}|}; \end{cases} \tag{3.38}$$

$$\|A_h(z) - h(z)\| \leq \begin{cases} \lambda \\ \frac{|\tau - 1|}{n\lambda \|z\|^\mu}; \\ \frac{|\tau - \tau^\mu|}{(n+1)\lambda \|z\|^{n\mu}}; \\ \frac{|\tau - \tau^{n\mu}|}{|\tau - \tau^{n\mu}|}; \end{cases} \tag{3.39}$$



*Proof.* If we replace

$$N(x_1, x_2, \dots, x_n) = \begin{cases} \lambda, \\ \sum_{a=1}^n \lambda \|x_a\|^\mu, \\ \prod_{a=1}^n \lambda \|x_a\|^\mu + \sum_{a=1}^n \lambda \|x_a\|^{n\mu}, \end{cases}$$

$$E(y_2, y_4, \dots, y_{2n}) = \begin{cases} \lambda, \\ \sum_{a=1}^n \lambda \|y_{2a}\|^\mu, \\ \prod_{a=1}^n \lambda \|y_{2a}\|^\mu + \sum_{a=1}^n \lambda \|y_{2a}\|^{n\mu}, \end{cases}$$

$$O(z_1, z_3, \dots, z_{2n-1}) = \begin{cases} \lambda, \\ \sum_{a=1}^n \lambda \|z_{2a-1}\|^\mu, \\ \prod_{a=1}^n \lambda \|z_{2a-1}\|^\mu + \sum_{a=1}^n \lambda \|z_{2a-1}\|^{n\mu}, \end{cases}$$

in Theorem 3.6, we arrive our desired result. □

### 4. Stability Results In 2- Banach Space

In this section, we establish the generalized Ulam - Hyers stability of the functional equations (1.27), (1.28) and (1.29) in 2- Banach spaces using Hyers Method.

Now, we give basic definitions and notations in 2- Banach spaces.

**Definition 4.1.** Let  $X$  be a linear space of dimension greater than 1. Suppose  $\|(\bullet, \bullet)\|$  is a real-valued function on  $X \times X$  satisfying the following conditions:

- (2N1)  $\|(x, y)\| = 0$  if and only if  $x, y$  are linearly dependent vectors,
- (2N2)  $\|(x, y)\| = \|(y, x)\|$  for all  $x, y \in X$ ,
- (2N3)  $\|(\lambda x, y)\| = |\lambda| \|(x, y)\|$  for all  $\lambda \in \mathbb{R}$  and for all  $x, y \in X$ ,
- (2N4)  $\|(x + y, z)\| \leq \|(x, z)\| + \|(y, z)\|$  for all  $x, y, z \in X$ .

Then  $\|(\bullet, \bullet)\|$  is called a 2-norm on  $X$  and the pair  $(X, \|(\bullet, \bullet)\|)$  is called 2-normed linear space.

**Definition 4.2.** A sequence  $\{x_n\}$  in a linear 2-normed space  $X$  is called a Cauchy sequence if there are two points  $y, z \in X$  such that  $y$  and  $z$  are linearly independent,

$$\lim_{l, m \rightarrow \infty} \|(x_l - x_m, y)\| = 0 \quad \text{and} \quad \lim_{l, m \rightarrow \infty} \|(x_l - x_m, z)\| = 0.$$

A sequence  $\{x_n\}$  in a linear 2-normed space  $X$  is called a convergent sequence if there is an  $x, w \in X$  such that

$$\lim_{n \rightarrow \infty} \|(x_n - x, w)\| = 0$$

for all  $y \in X$ . If  $\{x_n\}$  converges to  $x$ , write  $x_n \rightarrow x$  as  $n \rightarrow \infty$  and call  $x$  the limit of  $\{x_n\}$ . In this case, we also write  $\lim_{n \rightarrow \infty} x_n = x$ . A linear 2-normed space in which every Cauchy sequence is a convergent sequence is called a 2-Banach space.

To prove stability results, let us consider  $\mathcal{A}$  be a 2-normed space and  $\mathcal{B}$  be a 2-Banach space. The proof of the following theorem and corollary is similar lines to the of Theorem 3.6 and Corollary 3.7.

**Theorem 4.3.** Let  $f, g, h : \mathcal{A} \rightarrow \mathcal{B}$  be a mapping satisfying the following inequalities

$$\left\| \left( f \left( \sum_{a=1}^n a x_a \right) - \sum_{a=1}^n (a f(x_a)), u \right) \right\| \leq N(x_1, x_2, \dots, x_n); \quad (4.1)$$

$$\left\| \left( g \left( \sum_{a=1}^n 2a y_{2a} \right) - \sum_{a=1}^n (2a g(y_{2a})), u \right) \right\| \leq E(y_2, y_4, \dots, y_{2n}); \quad (4.2)$$

$$\left\| \left( h \left( \sum_{a=1}^n (2a - 1) z_{2a-1} \right) - \sum_{a=1}^n ((2a - 1) h(z_{2a-1})), u \right) \right\| \leq O(z_1, z_3, \dots, z_{2n-1}); \quad (4.3)$$

for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$  and all  $u \in \mathcal{B}$ , where  $N : \mathcal{A}^n \rightarrow [0, \infty)$ ,  $E : \mathcal{A}^{2n} \rightarrow [0, \infty)$  and  $O : \mathcal{A}^{2n-1} \rightarrow [0, \infty)$  satisfying the conditions

$$\lim_{\alpha \rightarrow \infty} \frac{N(\kappa^{\alpha\gamma} x_1, \kappa^{\alpha\gamma} x_2, \dots, \kappa^{\alpha\gamma} x_n)}{\kappa^{\alpha\gamma}} = 0; \quad (4.4)$$

$$\lim_{\alpha \rightarrow \infty} \frac{E(\rho^{\alpha\gamma} y_2, \rho^{\alpha\gamma} y_4, \dots, \rho^{\alpha\gamma} y_{2n})}{\rho^{\alpha\gamma}} = 0; \quad (4.5)$$

$$\lim_{\alpha \rightarrow \infty} \frac{O(\tau^{\alpha\gamma} z_1, \tau^{\alpha\gamma} z_3, \dots, \tau^{\alpha\gamma} z_{2n-1})}{\tau^{\alpha\gamma}} = 0; \quad (4.6)$$

for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$  with  $\gamma = \pm 1$ . Then there exists one and only additive mapping  $A_f, A_g, A_h : \mathcal{A} \rightarrow \mathcal{B}$  satisfying (1.27), (1.28), (1.29), and

$$\|(A_f(x) - f(x), u)\| \leq \frac{1}{\kappa} \sum_{\beta=\frac{1-\gamma}{2}}^{\infty} \frac{N(\kappa^{\beta\gamma} x, \kappa^{\beta\gamma} x, \dots, \kappa^{\beta\gamma} x)}{\kappa^{\beta\gamma}}; \quad (4.7)$$

$$\|(A_g(y) - g(y), u)\| \leq \frac{1}{\rho} \sum_{\beta=\frac{1-\gamma}{2}}^{\infty} \frac{E(\rho^{\beta\gamma} y, \rho^{\beta\gamma} y, \dots, \rho^{\beta\gamma} y)}{\rho^{\beta\gamma}}; \quad (4.8)$$

$$\|(A_h(z) - h(z), u)\| \leq \frac{1}{\tau} \sum_{\beta=\frac{1-\gamma}{2}}^{\infty} \frac{O(\tau^{\beta\gamma} z, \tau^{\beta\gamma} z, \dots, \tau^{\beta\gamma} z)}{\tau^{\beta\gamma}}; \quad (4.9)$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$ , respectively. The mappings  $A_f, A_g, A_h$  are respectively defined as

$$A_f(x) = \lim_{\alpha \rightarrow \infty} \frac{f(\kappa^{\alpha\gamma} x)}{\kappa^{\alpha\gamma}}; \quad (4.10)$$

$$A_g(y) = \lim_{\alpha \rightarrow \infty} \frac{g(\rho^{\alpha\gamma} y)}{\rho^{\alpha\gamma}}; \quad (4.11)$$

$$A_h(z) = \lim_{\alpha \rightarrow \infty} \frac{h(\tau^{\alpha\gamma} z)}{\tau^{\alpha\gamma}}; \quad (4.12)$$

for all  $x, y, z \in \mathcal{A}$ .





**Corollary 4.4.** Let  $f, g, h : \mathcal{A} \rightarrow \mathcal{B}$  be a mapping satisfying the following inequalities

$$\begin{aligned} & \left\| \left( f \left( \sum_{a=1}^n a x_a \right) - \sum_{a=1}^n (a f(x_a)), u \right) \right\| \\ & \leq \begin{cases} \lambda, & \mu \neq 1; \\ \sum_{a=1}^n \lambda \|x_a, u\|^\mu, & n\mu \neq 1; \\ \prod_{a=1}^n \lambda \|x_a, u\|^\mu + \sum_{a=1}^n \lambda \|x_a, u\|^{n\mu}, & n\mu \neq 1; \end{cases} \end{aligned} \tag{4.13}$$

$$\begin{aligned} & \left\| \left( g \left( \sum_{a=1}^n 2a y_{2a} \right) - \sum_{a=1}^n (2a g(y_{2a})), u \right) \right\| \\ & \leq \begin{cases} \lambda, & \mu \neq 1; \\ \sum_{a=1}^n \lambda \|y_{2a}, u\|^\mu, & n\mu \neq 1; \\ \prod_{a=1}^n \lambda \|y_{2a}, u\|^\mu + \sum_{a=1}^n \lambda \|y_{2a}, u\|^{n\mu}, & n\mu \neq 1; \end{cases} \end{aligned} \tag{4.14}$$

$$\begin{aligned} & \left\| \left( h \left( \sum_{a=1}^n (2a-1) z_{2a-1} \right) \right. \right. \\ & \quad \left. \left. - \sum_{a=1}^n ((2a-1) h(z_{2a-1})), u \right) \right\| \\ & \leq \begin{cases} \lambda, & \mu \neq 1; \\ \sum_{a=1}^n \lambda \|z_{2a-1}, u\|^\mu, & n\mu \neq 1; \\ \prod_{a=1}^n \lambda \|z_{2a-1}, u\|^\mu + \sum_{a=1}^n \lambda \|z_{2a-1}, u\|^{n\mu}, & n\mu \neq 1; \end{cases} \end{aligned} \tag{4.15}$$

for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$  and all  $u \in \mathcal{B}$ , where  $\lambda$  and  $\mu$  are positive constants. Then there exists one and only additive mapping  $A_f, A_g, A_h : \mathcal{A} \rightarrow \mathcal{B}$  satisfying (1.27), (1.28), (1.29), and

$$\| (A_f(x) - f(x), u) \| \leq \begin{cases} \frac{\lambda}{|\kappa - 1|}; \\ \frac{n\lambda \|x, u\|^\mu}{|\kappa - \kappa^\mu|}; \\ \frac{(n+1)\lambda \|x, u\|^{n\mu}}{|\kappa - \kappa^{n\mu}|}; \end{cases} \tag{4.16}$$

$$\| (A_g(y) - g(y), u) \| \leq \begin{cases} \frac{\lambda}{|\rho - 1|}; \\ \frac{n\lambda \|y, u\|^\mu}{|\rho - \rho^\mu|}; \\ \frac{(n+1)\lambda \|y, u\|^{n\mu}}{|\rho - \rho^{n\mu}|}; \end{cases} \tag{4.17}$$

$$\| (A_h(z) - h(z), u) \| \leq \begin{cases} \frac{\lambda}{|\tau - 1|}; \\ \frac{n\lambda \|z, u\|^\mu}{|\tau - \tau^\mu|}; \\ \frac{(n+1)\lambda \|z, u\|^{n\mu}}{|\tau - \tau^{n\mu}|}; \end{cases} \tag{4.18}$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$ , respectively.

## 5. Stability Results In Quasi 2-Banach Space

In this section, we investigate the generalized Ulam - Hyers stability of the functional equations (1.27), (1.28) and (1.29) in Quasi 2-Banach Space using Hyers Method.

Now, we give basic definitions and notations in Quasi 2-Banach Space.

**Definition 5.1.** Let  $X$  be a linear space of dimension greater than or equal to 2. Suppose  $\|(\bullet, \bullet)\|$  is a real-valued function on  $X \times X$  satisfying the following conditions:

(Q2N1)  $\|(x, y)\| = 0$  if and only if  $x, y$  are linearly dependent vectors,

(Q2N2)  $\|(x, y)\| = \|(y, x)\|$  for all  $x, y \in X$ ,

(Q2N3)  $\|(\lambda x, y)\| = |\lambda| \|(x, y)\|$  for all  $\lambda \in \mathbb{R}$  and for all  $x, y \in X$ ,

(Q2N4) It exists a constant  $K \geq 1$  such that

$$\|(x + y, z)\| \leq K(\|(x, z)\| + \|(y, z)\|)$$

for all  $x, y, z \in X$ .

Then  $\|(\bullet, \bullet)\|$  is called a quasi 2-norm on  $X$  and the pair  $(X, \|(\bullet, \bullet)\|)$  is called quasi 2-normed linear space. The smallest possible number  $K$  such that it satisfies the condition (Q2N4) is called a modulus of concavity of the quasi 2-norm  $\|(\bullet, \bullet)\|$ .

Sometimes the condition (Q2N4) called the triangle inequality. Further, M. Kir and M. Acikgoz [26] gave few examples of trivial quasi 2-normed spaces and consider the question about completing the quasi 2-normed space. A quasi 2-normed space in which every Cauchy sequence is a convergent sequence is called a quasi 2-Banach space.

To prove stability results, let us consider  $\mathcal{A}$  be a Quasi 2-normed space and  $\mathcal{B}$  be a Quasi 2-Banach space. The proof of the following theorem and corollary is similar lines to the of Theorem 3.6 and Corollary 3.7.

**Theorem 5.2.** Let  $f, g, h : \mathcal{A} \rightarrow \mathcal{B}$  be a mapping satisfying the following inequalities

$$\left\| \left( f \left( \sum_{a=1}^n a x_a \right) - \sum_{a=1}^n (a f(x_a)), u \right) \right\| \leq N(x_1, x_2, \dots, x_n); \tag{5.1}$$

$$\left\| \left( g \left( \sum_{a=1}^n 2a y_{2a} \right) - \sum_{a=1}^n (2a g(y_{2a})), u \right) \right\| \leq E(y_2, y_4, \dots, y_{2n}); \tag{5.2}$$

$$\left\| \left( h \left( \sum_{a=1}^n (2a-1) z_{2a-1} \right) - \sum_{a=1}^n ((2a-1) h(z_{2a-1})), u \right) \right\| \leq O(z_1, z_3, \dots, z_{2n-1}); \tag{5.3}$$

for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$  and all  $u \in \mathcal{B}$ , where  $N : \mathcal{A}^n \rightarrow [0, \infty)$ ,  $E : \mathcal{A}^{2n} \rightarrow [0, \infty)$  and



$O : \mathcal{A}^{2n-1} \rightarrow [0, \infty)$  satisfying the conditions

$$\lim_{\alpha \rightarrow \infty} \frac{N(\kappa^{\alpha\gamma}x_1, \kappa^{\alpha\gamma}x_2, \dots, \kappa^{\alpha\gamma}x_n)}{\kappa^{\alpha\gamma}} = 0; \tag{5.4}$$

$$\lim_{\alpha \rightarrow \infty} \frac{E(\rho^{\alpha\gamma}y_2, \rho^{\alpha\gamma}y_4, \dots, \rho^{\alpha\gamma}y_{2n})}{\rho^{\alpha\gamma}} = 0; \tag{5.5}$$

$$\lim_{\alpha \rightarrow \infty} \frac{O(\tau^{\alpha\gamma}z_1, \tau^{\alpha\gamma}z_3, \dots, \tau^{\alpha\gamma}z_{2n-1})}{\tau^{\alpha\gamma}} = 0; \tag{5.6}$$

for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$  with  $\gamma = \pm 1$ . Then there exists one and only additive mapping  $A_f, A_g, A_h : \mathcal{A} \rightarrow \mathcal{B}$  satisfying (1.27), (1.28), (1.29), and

$$\begin{aligned} & \| (A_f(x) - f(x), u) \| \\ & \leq \frac{K^{\alpha-1}}{\kappa} \sum_{\beta=\frac{1}{2}\gamma}^{\infty} \frac{N(\kappa^{\beta\gamma}x, \kappa^{\beta\gamma}x, \dots, \kappa^{\beta\gamma}x)}{\kappa^{\beta\gamma}}; \end{aligned} \tag{5.7}$$

$$\begin{aligned} & \| (A_g(y) - g(y), u) \| \\ & \leq \frac{K^{\alpha-1}}{\rho} \sum_{\beta=\frac{1}{2}\gamma}^{\infty} \frac{E(\rho^{\beta\gamma}y, \rho^{\beta\gamma}y, \dots, \rho^{\beta\gamma}y)}{\rho^{\beta\gamma}}; \end{aligned} \tag{5.8}$$

$$\begin{aligned} & \| (A_h(z) - h(z), u) \| \\ & \leq \frac{K^{\alpha-1}}{\tau} \sum_{\beta=\frac{1}{2}\gamma}^{\infty} \frac{O(\tau^{\beta\gamma}z, \tau^{\beta\gamma}z, \dots, \tau^{\beta\gamma}z)}{\tau^{\beta\gamma}}; \end{aligned} \tag{5.9}$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$ , respectively. The mappings  $A_f, A_g, A_h$  are respectively defined as

$$A_f(x) = \lim_{\alpha \rightarrow \infty} \frac{f(\kappa^{\alpha\gamma}x)}{\kappa^{\alpha\gamma}}; \tag{5.10}$$

$$A_g(y) = \lim_{\alpha \rightarrow \infty} \frac{g(\rho^{\alpha\gamma}y)}{\rho^{\alpha\gamma}}; \tag{5.11}$$

$$A_h(z) = \lim_{\alpha \rightarrow \infty} \frac{h(\tau^{\alpha\gamma}z)}{\tau^{\alpha\gamma}}; \tag{5.12}$$

for all  $x, y, z \in \mathcal{A}$ .

**Corollary 5.3.** Let  $f, g, h : \mathcal{A} \rightarrow \mathcal{B}$  be a mapping satisfying the following inequalities

$$\begin{aligned} & \left\| \left( f \left( \sum_{a=1}^n a x_a \right) - \sum_{a=1}^n (a f(x_a)), u \right) \right\| \\ & \leq \begin{cases} \lambda, & \mu \neq 1; \\ \sum_{a=1}^n \lambda \|x_a, u\|^\mu, & n\mu \neq 1; \\ \prod_{a=1}^n \lambda \|x_a, u\|^\mu + \sum_{a=1}^n \lambda \|x_a, u\|^{n\mu}, & n\mu \neq 1; \end{cases} \end{aligned} \tag{5.13}$$

$$\begin{aligned} & \left\| \left( g \left( \sum_{a=1}^n 2a y_{2a} \right) - \sum_{a=1}^n (2a g(y_{2a})), u \right) \right\| \\ & \leq \begin{cases} \lambda, & \mu \neq 1; \\ \sum_{a=1}^n \lambda \|y_{2a}, u\|^\mu, & n\mu \neq 1; \\ \prod_{a=1}^n \lambda \|y_{2a}, u\|^\mu + \sum_{a=1}^n \lambda \|y_{2a}, u\|^{n\mu}, & n\mu \neq 1; \end{cases} \end{aligned} \tag{5.14}$$

$$\begin{aligned} & \left\| \left( h \left( \sum_{a=1}^n (2a-1) z_{2a-1} \right) - \sum_{a=1}^n ((2a-1) h(z_{2a-1})), u \right) \right\| \\ & \leq \begin{cases} \lambda, & \mu \neq 1; \\ \sum_{a=1}^n \lambda \|z_{2a-1}, u\|^\mu, & n\mu \neq 1; \\ \prod_{a=1}^n \lambda \|z_{2a-1}, u\|^\mu + \sum_{a=1}^n \lambda \|z_{2a-1}, u\|^{n\mu}, & n\mu \neq 1; \end{cases} \end{aligned} \tag{5.15}$$

for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$  and all  $u \in \mathcal{B}$ , where  $\lambda$  and  $\mu$  are positive constants. Then there exists one and only additive mapping  $A_f, A_g, A_h : \mathcal{A} \rightarrow \mathcal{B}$  satisfying (1.27), (1.28), (1.29), and

$$\| (A_f(x) - f(x), u) \| \leq \begin{cases} \frac{K^{\alpha-1}\lambda}{|\kappa-1|}; \\ \frac{K^{\alpha-1}n\lambda \|x, u\|^\mu}{|\kappa-\kappa^\mu|}; \\ \frac{K^{\alpha-1}(n+1)\lambda \|x, u\|^{n\mu}}{|\kappa-\kappa^{n\mu}|}; \end{cases} \tag{5.16}$$

$$\| (A_g(y) - g(y), u) \| \leq \begin{cases} \frac{K^{\alpha-1}\lambda}{|\rho-1|}; \\ \frac{K^{\alpha-1}n\lambda \|y, u\|^\mu}{|\rho-\rho^\mu|}; \\ \frac{K^{\alpha-1}(n+1)\lambda \|y, u\|^{n\mu}}{|\rho-\rho^{n\mu}|}; \end{cases} \tag{5.17}$$

$$\| (A_h(z) - h(z), u) \| \leq \begin{cases} \frac{K^{\alpha-1}\lambda}{|\tau-1|}; \\ \frac{K^{\alpha-1}n\lambda \|z, u\|^\mu}{|\tau-\tau^\mu|}; \\ \frac{K^{\alpha-1}(n+1)\lambda \|z, u\|^{n\mu}}{|\tau-\tau^{n\mu}|}; \end{cases} \tag{5.18}$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$ , respectively.

## 6. Stability Results In Quasi-Beta-2-Banach Space

In this section, we discussed the generalized Ulam - Hyers stability of the functional equations (1.27), (1.28) and (1.29) in quasi- $\beta$ -2-Banach space using Hyers Method.

Now, we give basic definitions and notations in quasi- $\beta$ -2-Banach space.

**Definition 6.1.** Let  $X$  be a linear space of dimension greater than or equal to 2. Suppose  $\|(\bullet, \bullet)\|$  is a real-valued function on  $X \times X$  satisfying the following conditions:



(QB2N1)  $\|(x, y)\| = 0$  if and only if  $x, y$  are linearly dependent vectors,

(QB2N2)  $\|(x, y)\| = \|(y, x)\|$  for all  $x, y \in X$ ,

(QB2N3)  $\|(\lambda x, y)\| = |\lambda|^\beta \|(x, y)\|$  for all  $\lambda \in \mathbb{R}$  and for all  $x, y \in X$  where  $\beta$  is a real number with  $0 < \beta \leq 1$

(QB2N4) If exists a constant  $K \geq 1$  such that  $\|(x + y, z)\| \leq K(\|(x, z)\| + \|(y, z)\|)$  for all  $x, y, z \in X$ .

The pair  $(X, \|(\bullet, \bullet)\|)$  is called quasi- $\beta$ -normed space if  $\|(\bullet, \bullet)\|$  is a quasi- $\beta$ -2-norm on  $X$ . The smallest possible  $K$  is called the modulus of concavity of  $\|\cdot\|$ .

**Definition 6.2.** A quasi- $\beta$ -2-Banach space is a complete quasi- $\beta$ -normed space.

To prove stability results, let us consider  $\mathcal{A}$  as a quasi- $\beta$ -2-Banach space and  $\mathcal{B}$  as a quasi- $\beta$ -2-Banach space.

**Theorem 6.3.** Let  $f, g, h : \mathcal{A} \rightarrow \mathcal{B}$  be a mapping satisfying the following inequalities

$$\left\| \left( f \left( \sum_{a=1}^n a x_a \right) - \sum_{a=1}^n (a f(x_a)), u \right) \right\| \leq N(x_1, x_2, \dots, x_n); \tag{6.1}$$

$$\left\| \left( g \left( \sum_{a=1}^n 2a y_{2a} \right) - \sum_{a=1}^n (2a g(y_{2a})), u \right) \right\| \leq E(y_2, y_4, \dots, y_{2n}); \tag{6.2}$$

$$\left\| \left( h \left( \sum_{a=1}^n (2a-1) z_{2a-1} \right) - \sum_{a=1}^n ((2a-1) h(z_{2a-1})), u \right) \right\| \leq O(z_1, z_3, \dots, z_{2n-1}); \tag{6.3}$$

for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$  and all  $u \in \mathcal{B}$ , where  $N : \mathcal{A}^n \rightarrow [0, \infty)$ ,  $E : \mathcal{A}^{2n} \rightarrow [0, \infty)$  and  $O : \mathcal{A}^{2n-1} \rightarrow [0, \infty)$  satisfying the conditions

$$\lim_{\alpha \rightarrow \infty} \frac{N(\kappa^{\alpha\gamma} x_1, \kappa^{\alpha\gamma} x_2, \dots, \kappa^{\alpha\gamma} x_n)}{\kappa^{\alpha\gamma}} = 0; \tag{6.4}$$

$$\lim_{\alpha \rightarrow \infty} \frac{E(\rho^{\alpha\gamma} y_2, \rho^{\alpha\gamma} y_4, \dots, \rho^{\alpha\gamma} y_{2n})}{\rho^{\alpha\gamma}} = 0; \tag{6.5}$$

$$\lim_{\alpha \rightarrow \infty} \frac{O(\tau^{\alpha\gamma} z_1, \tau^{\alpha\gamma} z_3, \dots, \tau^{\alpha\gamma} z_{2n-1})}{\tau^{\alpha\gamma}} = 0; \tag{6.6}$$

for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$  with  $\gamma = \pm 1$ . Then there exists one and only additive mapping

$A_f, A_g, A_h : \mathcal{A} \rightarrow \mathcal{B}$  satisfying (1.27), (1.28), (1.29), and

$$\begin{aligned} & \| (A_f(x) - f(x), u) \| \\ & \leq \frac{K^{\alpha-1}}{\kappa^\beta} \sum_{\eta=\frac{1-\gamma}{2}}^{\infty} \frac{N(\kappa^{\eta\gamma} x, \kappa^{\eta\gamma} x, \dots, \kappa^{\eta\gamma} x)}{\kappa^{\eta\gamma}}; \end{aligned} \tag{6.7}$$

$$\begin{aligned} & \| (A_g(y) - g(y), u) \| \\ & \leq \frac{K^{\alpha-1}}{\rho^\beta} \sum_{\eta=\frac{1-\gamma}{2}}^{\infty} \frac{E(\rho^{\eta\gamma} y, \rho^{\eta\gamma} y, \dots, \rho^{\eta\gamma} y)}{\rho^{\eta\gamma}}; \end{aligned} \tag{6.8}$$

$$\begin{aligned} & \| (A_h(z) - h(z), u) \| \\ & \leq \frac{K^{\alpha-1}}{\tau^\beta} \sum_{\eta=\frac{1-\gamma}{2}}^{\infty} \frac{O(\tau^{\eta\gamma} z, \tau^{\eta\gamma} z, \dots, \tau^{\eta\gamma} z)}{\tau^{\eta\gamma}}; \end{aligned} \tag{6.9}$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$ , respectively. The mappings  $A_f, A_g, A_h$  are respectively defined as

$$A_f(x) = \lim_{\alpha \rightarrow \infty} \frac{f(\kappa^{\alpha\gamma} x)}{\kappa^{\alpha\gamma}}; \tag{6.10}$$

$$A_g(y) = \lim_{\alpha \rightarrow \infty} \frac{g(\rho^{\alpha\gamma} y)}{\rho^{\alpha\gamma}}; \tag{6.11}$$

$$A_h(z) = \lim_{\alpha \rightarrow \infty} \frac{h(\tau^{\alpha\gamma} z)}{\tau^{\alpha\gamma}}; \tag{6.12}$$

for all  $x, y, z \in \mathcal{A}$ .

*Proof.* Changing

$$(x_1, x_2, \dots, x_n) = (x, x, \dots, x) \text{ in (6.1);}$$

$$(y_2, y_4, \dots, y_{2n}) = (y, y, \dots, y) \text{ in (6.2);}$$

$$(z_1, z_3, \dots, z_{2n-1}) = (z, z, \dots, z) \text{ in (6.3);}$$

we arrive the following inequalities

$$\left\| f \left( \sum_{a=1}^n a x \right) - \left( \sum_{a=1}^n a \right) f(x) \right\| \leq N(x, x, \dots, x); \tag{6.13}$$

$$\left\| g \left( \sum_{a=1}^n 2a y \right) - \left( \sum_{a=1}^n 2a \right) g(y) \right\| \leq E(y, y, \dots, y); \tag{6.14}$$

$$\left\| h \left( \sum_{a=1}^n (2a-1) z \right) - \left( \sum_{a=1}^n (2a-1) \right) h(z) \right\| \leq O(z, z, \dots, z); \tag{6.15}$$

for all  $x, y, z \in \mathcal{A}$ . Define

$$\sum_{a=1}^n a = \frac{n(n+1)}{2} = \kappa \text{ in (6.13);}$$

$$\sum_{a=1}^n 2a = n(n+1) = \rho \text{ in (6.14);}$$

$$\sum_{a=1}^n (2a-1) = n^2 = \tau \text{ in (6.15);}$$



we obtain the succeeding inequalities

$$\|f(\kappa x) - \kappa f(x)\| \leq N(x, x, \dots, x); \tag{6.16}$$

$$\|g(\rho y) - \rho g(y)\| \leq E(y, y, \dots, y); \tag{6.17}$$

$$\|h(\tau z) - \tau h(z)\| \leq O(z, z, \dots, z); \tag{6.18}$$

for all  $x, y, z \in \mathcal{A}$ . It follows from above inequalities

$$\left\| \frac{f(\kappa x)}{\kappa} - f(x) \right\| \leq \frac{N(x, x, \dots, x)}{\kappa^\beta}; \tag{6.19}$$

$$\left\| \frac{g(\rho y)}{\rho} - g(y) \right\| \leq \frac{E(y, y, \dots, y)}{\rho^\beta}; \tag{6.20}$$

$$\left\| \frac{h(\tau z)}{\tau} - h(z) \right\| \leq \frac{O(z, z, \dots, z)}{\tau^\beta}; \tag{6.21}$$

for all  $x, y, z \in \mathcal{A}$ . Replacing

$$x = \kappa x \text{ and } \div \kappa \text{ in (6.19);}$$

$$y = \rho y \text{ and } \div \rho \text{ in (6.20);}$$

$$z = \tau z \text{ and } \div \tau \text{ in (6.21);}$$

we arrive the following inequalities

$$\left\| \frac{f(\kappa^2 x)}{\kappa^2} - \frac{f(\kappa x)}{\kappa} \right\| \leq \frac{N(\kappa x, \kappa x, \dots, \kappa x)}{\kappa^\beta \kappa}; \tag{6.22}$$

$$\left\| \frac{g(\rho^2 y)}{\rho^2} - \frac{g(\rho y)}{\rho} \right\| \leq \frac{E(\rho y, \rho y, \dots, \rho y)}{\rho^\beta \rho}; \tag{6.23}$$

$$\left\| \frac{h(\tau^2 z)}{\tau^2} - \frac{h(\tau z)}{\tau} \right\| \leq \frac{O(\tau z, \tau z, \dots, \tau z)}{\tau^\beta \tau}; \tag{6.24}$$

for all  $x, y, z \in \mathcal{A}$ . With the help of triangle inequality from (6.19) and (6.22); (6.20) and (6.23); (6.21) and (6.24);

we achieve the subsequent inequalities

$$\begin{aligned} & \left\| \frac{f(\kappa^2 x)}{\kappa^2} - f(x) \right\| \\ & \leq \left\| \frac{f(\kappa^2 x)}{\kappa^2} - \frac{f(\kappa x)}{\kappa} \right\| + \left\| \frac{f(\kappa x)}{\kappa} - f(x) \right\| \\ & \leq \frac{K}{\kappa^\beta} \left( N(x, x, \dots, x) + \frac{N(\kappa x, \kappa x, \dots, \kappa x)}{\kappa} \right); \end{aligned} \tag{6.25}$$

$$\begin{aligned} & \left\| \frac{g(\rho^2 y)}{\rho^2} - g(y) \right\| \\ & \leq \left\| \frac{g(\rho^2 y)}{\rho^2} - \frac{g(\rho y)}{\rho} \right\| + \left\| \frac{g(\rho y)}{\rho} - g(y) \right\| \\ & \leq \frac{K}{\rho^\beta} \left( E(y, y, \dots, y) + \frac{E(\rho y, \rho y, \dots, \rho y)}{\rho} \right); \end{aligned} \tag{6.26}$$

$$\begin{aligned} & \left\| \frac{h(\tau^2 z)}{\tau^2} - h(z) \right\| \\ & \leq \left\| \frac{h(\tau^2 z)}{\tau^2} - \frac{h(\tau z)}{\tau} \right\| + \left\| \frac{h(\tau z)}{\tau} - h(z) \right\| \\ & \leq \frac{K}{\tau^\beta} \left( E(z, z, \dots, z) + \frac{E(\tau z, \tau z, \dots, \tau z)}{\tau} \right); \end{aligned} \tag{6.27}$$

for all  $x, y, z \in \mathcal{A}$ . Generalizing for a positive integer  $\alpha$ , we arrive

$$\left\| \frac{f(\kappa^\alpha x)}{\kappa^\alpha} - f(x) \right\| \leq \frac{K^{\alpha-1}}{\kappa^\beta} \sum_{\eta=0}^{\alpha} \frac{N(\kappa^\eta x, \kappa^\eta x, \dots, \kappa^\eta x)}{\kappa^\eta}; \tag{6.28}$$

$$\left\| \frac{g(\rho^\alpha y)}{\rho^\alpha} - g(y) \right\| \leq \frac{K^{\alpha-1}}{\rho^\beta} \sum_{\eta=0}^{\alpha} \frac{E(\rho^\eta y, \rho^\eta y, \dots, \rho^\eta y)}{\rho^\eta}; \tag{6.29}$$

$$\left\| \frac{h(\tau^\alpha z)}{\tau^\alpha} - h(z) \right\| \leq \frac{K^{\alpha-1}}{\tau^\beta} \sum_{\eta=0}^{\alpha} \frac{O(\tau^\eta z, \tau^\eta z, \dots, \tau^\eta z)}{\tau^\eta}; \tag{6.30}$$

for all  $x, y, z \in \mathcal{A}$ . The rest of proof is similar lines to the of Theorem 3.6.  $\square$

**Corollary 6.4.** Let  $f, g, h : \mathcal{A} \rightarrow \mathcal{B}$  be a mapping satisfying the following inequalities

$$\begin{aligned} & \left\| \left( f \left( \sum_{a=1}^n a x_a \right) - \sum_{a=1}^n (a f(x_a)), u \right) \right\| \\ & \leq \begin{cases} \lambda, & \mu \neq 1; \\ \prod_{a=1}^n \lambda \|x_a, u\|^\mu + \sum_{a=1}^n \lambda \|x_a, u\|^{n\mu}, & n\mu \neq 1; \end{cases} \end{aligned} \tag{6.31}$$

$$\begin{aligned} & \left\| \left( g \left( \sum_{a=1}^n 2a y_{2a} \right) - \sum_{a=1}^n (2a g(y_{2a})), u \right) \right\| \\ & \leq \begin{cases} \lambda, & \mu \neq 1; \\ \prod_{a=1}^n \lambda \|y_{2a}, u\|^\mu + \sum_{a=1}^n \lambda \|y_{2a}, u\|^{n\mu}, & n\mu \neq 1; \end{cases} \end{aligned} \tag{6.32}$$

$$\begin{aligned} & \left\| \left( h \left( \sum_{a=1}^n (2a-1) z_{2a-1} \right) - \sum_{a=1}^n ((2a-1) h(z_{2a-1})), u \right) \right\| \\ & \leq \begin{cases} \lambda, & \mu \neq 1; \\ \prod_{a=1}^n \lambda \|z_{2a-1}, u\|^\mu + \sum_{a=1}^n \lambda \|z_{2a-1}, u\|^{n\mu}, & n\mu \neq 1; \end{cases} \end{aligned} \tag{6.33}$$

for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$  and all  $u \in \mathcal{B}$ , where  $\lambda$  and  $\mu$  are positive constants. Then there exists one and only additive mapping  $A_f, A_g, A_h : \mathcal{A} \rightarrow \mathcal{B}$



satisfying (1.27), (1.28), (1.29), and

$$\|(A_f(x) - f(x), u)\| \leq \begin{cases} \frac{K^{\alpha-1} \lambda \kappa}{\kappa^\beta |\kappa - 1|}; \\ \frac{K^{\alpha-1} n \lambda \kappa \|x, u\|^\mu}{\kappa^\beta |\kappa - \kappa^\mu|}; \\ \frac{K^{\alpha-1} (n+1) \lambda \kappa \|x, u\|^{n\mu}}{\kappa^\beta |\kappa - \kappa^{n\mu}|}; \end{cases} \tag{6.34}$$

$$\|(A_g(y) - g(y), u)\| \leq \begin{cases} \frac{K^{\alpha-1} \lambda \rho}{\rho^\beta |\rho - 1|}; \\ \frac{K^{\alpha-1} n \lambda \rho \|y, u\|^\mu}{\rho^\beta |\rho - \rho^\mu|}; \\ \frac{K^{\alpha-1} (n+1) \lambda \rho \|y, u\|^{n\mu}}{\rho^\beta |\rho - \rho^{n\mu}|}; \end{cases} \tag{6.35}$$

$$\|(A_h(z) - h(z), u)\| \leq \begin{cases} \frac{K^{\alpha-1} \lambda \tau}{\tau^\beta |\tau - 1|}; \\ \frac{K^{\alpha-1} n \lambda \tau \|z, u\|^\mu}{\tau^\beta |\tau - \tau^\mu|}; \\ \frac{K^{\alpha-1} (n+1) \lambda \tau \|z, u\|^{n\mu}}{\tau^\beta |\tau - \tau^{n\mu}|}; \end{cases} \tag{6.36}$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$ , respectively.

### 7. Stability Results In Fuzzy Quasi-Beta-2-Banach Space

In this section, we investigate the generalized Ulam - Hyers stability of the functional equations (1.27), (1.28) and (1.29) in fuzzy quasi- $\beta$ -2-Banach spaces using Hyers Method.

Now, we give basic definitions and notations in Fuzzy quasi- $\beta$ -2-Banach space.

**Definition 7.1.** Let  $X$  be a linear space of dimension greater than or equal to 2. A function  $N : X \times X \times \mathbb{R} \rightarrow [0, 1]$  is said to be a fuzzy quasi- $\beta$ -2-norm on  $X$  if for all  $x, y, z \in X$  and all  $s, t \in \mathbb{R}$ ,

- (2QBFN1)  $N(x, z, c) = 0$  for  $c \leq 0$ ;
- (2QBFN2)  $x = 0$  if and only if  $N(x, z, c) = 1$  for all  $c > 0$ ;
- (2QBFN3)  $N(cx, z, t) = N\left(x, z, \frac{t}{|c|^\beta}\right)$  if  $c \neq 0$  where  $\beta$  is a real number with  $0 < \beta \leq 1$
- (2QBFN4)  $N(x + y, z, s + t) \geq \min\{N(x, z, Ks), N(y, z, Kt)\}$ ; a constant  $K \geq 1$
- (2QBFN5)  $N(x, z, \cdot)$  is a non-decreasing function on  $\mathbb{R}$  and  $\lim_{t \rightarrow \infty} N(x, z, t) = 1$ ;
- (2QBFN6) for  $x \neq 0, N(x, z, \cdot)$  is (upper semi) continuous on  $\mathbb{R}$ .

The pair  $(X, N)$  is called a fuzzy quasi- $\beta$ -2-Banach space.

**Example 7.2.** Let  $X$  be a linear space. Then

$$N(x, z, t) = \begin{cases} \frac{t}{t + \|x\|}, & t > 0, \quad x, z \in X, \\ 0, & t \leq 0, \quad x, z \in X \end{cases}$$

is a fuzzy quasi- $\beta$ -2-normed space on  $X$ .

**Example 7.3.** Let  $X$  be a linear space. Then

$$N(x, z, t) = \begin{cases} 0, & t \leq 0, \\ \frac{t}{\|x\|}, & 0 < t \leq \|x\|, z \in X \\ 1, & t > \|x\|, z \in X \end{cases}$$

is a fuzzy quasi- $\beta$ -2-normed space on  $X$ .

**Definition 7.4.** Let  $X$  be a fuzzy quasi- $\beta$ -2-normed space. Let  $x_n$  be a sequence in  $X$ . Then  $x_n$  is said to be convergent if there exists  $x, z \in X$  such that  $\lim_{n \rightarrow \infty} N(x_n - x, z, t) = 1$  for all  $t > 0$ . In that case,  $x$  is called the limit of the sequence  $x_n$  and we denote it by  $N - \lim_{n \rightarrow \infty} x_n = x$ .

**Definition 7.5.** A sequence  $x_n$  in  $X$  is called Cauchy if for each  $\varepsilon > 0$  and each  $t > 0$  there exists  $n_0$  such that for all  $n \geq n_0$  and all  $p > 0$ , we have  $N(x_{n+p} - x_n, z, t) > 1 - \varepsilon$ .

**Definition 7.6.** Every convergent sequence in a fuzzy quasi- $\beta$ -2-normed space is Cauchy. If each Cauchy sequence is convergent, then the 2-norm is said to be complete and the fuzzy quasi- $\beta$ -2-normed space is called a fuzzy quasi- $\beta$ -2-Banach space.

To prove stability results, let us consider  $\mathcal{A}$  is a fuzzy quasi- $\beta$ -2-normed space and  $\mathcal{B}$  is a fuzzy quasi- $\beta$ -2-Banach space.

**Theorem 7.7.** Let  $f, g, h : \mathcal{A} \rightarrow \mathcal{B}$  be a mapping satisfying the following inequalities

$$\begin{aligned} & \mathcal{N} \left( \left( f \left( \sum_{a=1}^n a x_a \right) - \sum_{a=1}^n (a f(x_a)) \right), u \right), t \\ & \geq \mathcal{N}'(N(x_1, x_2, \dots, x_n), u, t); \end{aligned} \tag{7.1}$$

$$\begin{aligned} & \mathcal{N} \left( \left( g \left( \sum_{a=1}^n 2a y_{2a} \right) - \sum_{a=1}^n (2a g(y_{2a})) \right), u \right), t \\ & \geq \mathcal{N}'(E(y_2, y_4, \dots, y_{2n}), u, t); \end{aligned} \tag{7.2}$$

$$\begin{aligned} & \mathcal{N} \left( \left( h \left( \sum_{a=1}^n (2a - 1) z_{2a-1} \right) - \sum_{a=1}^n ((2a - 1) h(z_{2a-1})) \right), u \right), t \\ & \geq \mathcal{N}'(O(z_1, z_3, \dots, z_{2n-1}), u, t); \end{aligned} \tag{7.3}$$

for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ , where  $N : \mathcal{A}^n \rightarrow [0, \infty)$ ,  $E : \mathcal{A}^{2n} \rightarrow [0, \infty)$  and  $O : \mathcal{A}^{2n-1} \rightarrow [0, \infty)$  satisfying the conditions

$$\lim_{\alpha \rightarrow \infty} \mathcal{N}'(N(\kappa^{\alpha\gamma} x_1, \kappa^{\alpha\gamma} x_2, \dots, \kappa^{\alpha\gamma} x_n), u, \kappa^{\alpha\gamma} t) = 1; \tag{7.4}$$

$$\lim_{\alpha \rightarrow \infty} \mathcal{N}'(E(\rho^{\alpha\gamma} y_2, \rho^{\alpha\gamma} y_4, \dots, \rho^{\alpha\gamma} y_{2n}), u, \rho^{\alpha\gamma} t) = 1; \tag{7.5}$$

$$\lim_{\alpha \rightarrow \infty} \mathcal{N}'(O(\tau^{\alpha\gamma} z_1, \tau^{\alpha\gamma} z_3, \dots, \tau^{\alpha\gamma} z_{2n-1}), u, \tau^{\alpha\gamma} t) = 1; \tag{7.6}$$



with the conditions

$$\begin{aligned} & \mathcal{N}'(N(\kappa^\alpha x_1, \kappa^\alpha x_2, \dots, \kappa^\alpha x_n), u, t) \\ &= \mathcal{N}'(v^\alpha N(x_1, x_2, \dots, x_n), u, t); \end{aligned} \quad (7.7)$$

$$\begin{aligned} & \mathcal{N}'(E(\rho^\alpha y_2, \rho^\alpha y_4, \dots, \rho^\alpha y_{2n}), u, t) \\ &= \mathcal{N}'(v^\alpha E(y_2, y_4, \dots, y_{2n}), u, t); \end{aligned} \quad (7.8)$$

$$\begin{aligned} & \mathcal{N}'(O(\tau^\alpha z_1, \tau^\alpha z_3, \dots, \tau^\alpha z_{2n-1}), u, t) \\ &= \mathcal{N}'(v^\alpha O(z_1, z_3, \dots, z_{2n-1}), u, t); \end{aligned} \quad (7.9)$$

for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$  all  $u \in \mathcal{B}$  and  $t > 0$  with  $\gamma = \pm 1$ . Then there exists one and only additive mapping  $A_f, A_g, A_h : \mathcal{A} \rightarrow \mathcal{B}$  satisfying (1.27), (1.28), (1.29), and

$$\begin{aligned} & \mathcal{N}((A_f(x) - f(x), u), t) \\ & \geq \mathcal{N}'(N(x, x, \dots, x), u, K\kappa^{\beta-1}|\kappa - v|t); \end{aligned} \quad (7.10)$$

$$\begin{aligned} & \mathcal{N}((A_g(y) - g(y), u), t) \\ & \geq \mathcal{N}'(E(y, y, \dots, y), u, K\rho^{\beta-1}|\rho - v|t); \end{aligned} \quad (7.11)$$

$$\begin{aligned} & \mathcal{N}((A_h(z) - h(z), u), t) \\ & \geq \mathcal{N}'(O(z, z, \dots, z), u, K\tau^{\beta-1}|\tau - v|t); \end{aligned} \quad (7.12)$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ , respectively. The mappings  $A_f, A_g, A_h$  are respectively defined as

$$\lim_{\alpha \rightarrow \infty} \mathcal{N} \left( \left( A_f(x) - \frac{f(\kappa^{\alpha\gamma}x)}{\kappa^{\alpha\gamma}}, u \right), t \right) = 1; \quad (7.13)$$

$$\lim_{\alpha \rightarrow \infty} \mathcal{N} \left( \left( A_g(y) - \frac{g(\rho^{\alpha\gamma}y)}{\rho^{\alpha\gamma}}, u \right), t \right) = 1; \quad (7.14)$$

$$\lim_{\alpha \rightarrow \infty} \mathcal{N} \left( \left( A_h(z) - \frac{h(\tau^{\alpha\gamma}z)}{\tau^{\alpha\gamma}}, u \right), t \right) = 1; \quad (7.15)$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ .

*Proof.* Changing

$$(x_1, x_2, \dots, x_n) = (x, x, \dots, x) \quad \text{in (7.1);}$$

$$(y_2, y_4, \dots, y_{2n}) = (y, y, \dots, y) \quad \text{in (7.2);}$$

$$(z_1, z_3, \dots, z_{2n-1}) = (z, z, \dots, z) \quad \text{in (7.3);}$$

we arrive the following inequalities

$$\begin{aligned} & \mathcal{N} \left( \left( f \left( \sum_{a=1}^n a x \right) - \left( \sum_{a=1}^n a \right) f(x), u \right), t \right) \\ & \geq \mathcal{N}'(N(x, x, \dots, x), u, t); \end{aligned} \quad (7.16)$$

$$\begin{aligned} & \mathcal{N} \left( \left( g \left( \sum_{a=1}^n 2a y \right) - \left( \sum_{a=1}^n 2a \right) g(y), u \right), t \right) \\ & \geq \mathcal{N}'(E(y, y, \dots, y), u, t); \end{aligned} \quad (7.17)$$

$$\begin{aligned} & \mathcal{N} \left( \left( h \left( \sum_{a=1}^n (2a-1) z \right) - \left( \sum_{a=1}^n (2a-1) \right) h(z), u \right), t \right) \\ & \geq \mathcal{N}'(O(z, z, \dots, z), u, t); \end{aligned} \quad (7.18)$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and all  $t > 0$ . Define

$$\sum_{a=1}^n a = \frac{n(n+1)}{2} = \kappa \quad \text{in (7.16);}$$

$$\sum_{a=1}^n 2a = n(n+1) = \rho \quad \text{in (7.17);}$$

$$\sum_{a=1}^n (2a-1) = n^2 = \tau \quad \text{in (7.18);}$$

we obtain the succeeding inequalities

$$\mathcal{N}((f(\kappa x) - \kappa f(x), u), t) \geq \mathcal{N}'(N(x, x, \dots, x), u, t); \quad (7.19)$$

$$\mathcal{N}((g(\rho y) - \rho g(y), u), t) \geq \mathcal{N}'(E(y, y, \dots, y), u, t); \quad (7.20)$$

$$\mathcal{N}((h(\tau z) - \tau h(z), u), t) \geq \mathcal{N}'(O(z, z, \dots, z), u, t); \quad (7.21)$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ . Using (2QBFN3) it follows from above inequalities

$$\mathcal{N} \left( \left( \frac{f(\kappa x)}{\kappa} - f(x), u \right), t \right) \geq \mathcal{N}'(N(x, x, \dots, x), u, \kappa^\beta t); \quad (7.22)$$

$$\mathcal{N} \left( \left( \frac{g(\rho y)}{\rho} - g(y), u \right), t \right) \geq \mathcal{N}'(E(y, y, \dots, y), u, \rho^\beta t); \quad (7.23)$$

$$\mathcal{N} \left( \left( \frac{h(\tau z)}{\tau} - h(z), u \right), t \right) \geq \mathcal{N}'(O(z, z, \dots, z), u, \tau^\beta t); \quad (7.24)$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ . Replacing, using and substituting

$$x = \kappa^\alpha x, \quad (2QBFN3), (7.7) \text{ in (7.22) and } t = v^{\alpha\beta} t \text{ in (7.22);}$$

$$y = \rho^\alpha y, \quad (2QBFN3), (7.8) \text{ in (7.23) and } t = v^{\alpha\beta} t \text{ in (7.23);}$$

$$z = \tau^\alpha z, \quad (2QBFN3), (7.9) \text{ in (7.24) and } t = v^{\alpha\beta} t \text{ in (7.24);}$$

we arrive the following inequalities

$$\begin{aligned} & \mathcal{N} \left( \left( \frac{f(\kappa^{\alpha+1} x)}{\kappa^{\alpha+1}} - \frac{f(\kappa^\alpha x)}{\kappa^\alpha}, u \right), \frac{v^{\alpha\beta} t}{\kappa^{\alpha\beta}} \right) \\ & \geq \mathcal{N}'(N(x, x, \dots, x), u, \kappa^\beta t); \end{aligned} \quad (7.25)$$

$$\begin{aligned} & \mathcal{N} \left( \left( \frac{g(\rho^{\alpha+1} y)}{\rho^{\alpha+1}} - \frac{g(\rho^\alpha y)}{\rho^\alpha}, u \right), \frac{v^{\alpha\beta} t}{\rho^{\alpha\beta}} \right) \\ & \geq \mathcal{N}'(E(y, y, \dots, y), u, \rho^\beta t); \end{aligned} \quad (7.26)$$

$$\begin{aligned} & \mathcal{N} \left( \left( \frac{h(\tau^{\alpha+1} z)}{\tau^{\alpha+1}} - \frac{h(\tau^\alpha z)}{\tau^\alpha}, u \right), \frac{v^{\alpha\beta} t}{\tau^{\alpha\beta}} \right) \\ & \geq \mathcal{N}'(O(z, z, \dots, z), u, \tau^\beta t); \end{aligned} \quad (7.27)$$





for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ . It is easy to see that we have

$$\frac{f(\kappa^\alpha x)}{\kappa^\alpha} - f(x) = \sum_{\sigma=0}^{\alpha-1} \frac{f(\kappa^{\sigma+1} x)}{\kappa^{\sigma+1}} - \frac{f(\kappa^\sigma x)}{\kappa^\sigma}; \quad (7.28)$$

$$\frac{g(\rho^\alpha y)}{\rho^\alpha} - g(y) = \sum_{\sigma=0}^{\alpha-1} \frac{g(\rho^{\sigma+1} y)}{\rho^{\sigma+1}} - \frac{g(\rho^\sigma y)}{\rho^\sigma}; \quad (7.29)$$

$$\frac{h(\tau^\alpha z)}{\tau^\alpha} - h(z) = \sum_{\sigma=0}^{\alpha-1} \frac{h(\tau^{\sigma+1} z)}{\tau^{\sigma+1}} - \frac{h(\tau^\sigma z)}{\tau^\sigma}; \quad (7.30)$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ . From (7.28) and (7.25); (7.29) and (7.26); (7.30) and (7.27);

we achieve the subsequent inequalities

$$\begin{aligned} & \mathcal{N} \left( \left( \frac{f(\kappa^\alpha x)}{\kappa^\alpha} - f(x), u \right), \sum_{\sigma=0}^{\alpha-1} \frac{v^{\sigma\beta} t}{\kappa^{\sigma\beta}} \right) \\ & \geq \min \bigcup_{\sigma=0}^{\alpha-1} \left\{ \mathcal{N} \left( \left( \frac{f(\kappa^{\sigma+1} x)}{\kappa^{\sigma+1}} - \frac{f(\kappa^\sigma x)}{\kappa^\sigma}, u \right), \frac{K v^{\sigma\beta} t}{\kappa^{\sigma\beta}} \right) \right\} \geq \mathcal{N}' \left( N(x, x, \dots, x), u, K \kappa^\beta t \right); \end{aligned} \quad (7.31)$$

$$\begin{aligned} & \mathcal{N} \left( \left( \frac{g(\rho^\alpha y)}{\rho^\alpha} - g(y), u \right), \sum_{\sigma=0}^{\alpha-1} \frac{v^{\sigma\beta} t}{\rho^{\sigma\beta}} \right) \\ & \geq \min \bigcup_{\sigma=0}^{\alpha-1} \left\{ \mathcal{N} \left( \left( \frac{g(\rho^{\sigma+1} y)}{\rho^{\sigma+1}} - \frac{g(\rho^\sigma y)}{\rho^\sigma}, u \right), \frac{K v^{\sigma\beta} t}{\rho^{\sigma\beta}} \right) \right\} \geq \mathcal{N}' \left( N(y, y, \dots, y), u, K \rho^\beta t \right); \end{aligned} \quad (7.32)$$

$$\begin{aligned} & \mathcal{N} \left( \left( \frac{h(\tau^\alpha z)}{\tau^\alpha} - h(z), u \right), \sum_{\sigma=0}^{\alpha-1} \frac{v^{\sigma\beta} t}{\tau^{\sigma\beta}} \right) \\ & \geq \min \bigcup_{\sigma=0}^{\alpha-1} \left\{ \mathcal{N} \left( \left( \frac{h(\tau^{\sigma+1} z)}{\tau^{\sigma+1}} - \frac{h(\tau^\sigma z)}{\tau^\sigma}, u \right), \frac{K v^{\sigma\beta} t}{\tau^{\sigma\beta}} \right) \right\} \geq \mathcal{N}' \left( N(z, z, \dots, z), u, K \tau^\beta t \right); \end{aligned} \quad (7.33)$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ . It is easy to verify that the sequences

$$\left\{ \frac{f(\kappa^\alpha x)}{\kappa^\alpha} \right\}; \left\{ \frac{g(\rho^\alpha y)}{\rho^\alpha} \right\}; \left\{ \frac{h(\tau^\alpha z)}{\tau^\alpha} \right\}$$

are Cauchy sequences. Indeed, changing

$$x = \kappa^\delta x \text{ and using (2QBFN3), (7.7) in (7.31);}$$

$$y = \rho^\delta y \text{ and using (2QBFN3), (7.8) in (7.32);}$$

$$z = \tau^\delta z \text{ and using (2QBFN3), (7.9) in (7.33);}$$

$$\begin{aligned} & \mathcal{N} \left( \left( \frac{f(\kappa^{\alpha+\delta} x)}{\kappa^{\alpha+\delta}} - \frac{f(\kappa^\delta x)}{\kappa^\delta}, u \right), t \right) \\ & \geq \mathcal{N}' \left( N(x, x, \dots, x), u, \frac{K \kappa^\beta t}{\sum_{\sigma=0}^{\alpha-1} \frac{v^{(\sigma+\delta)\beta}}{\kappa^{(\sigma+\delta)\beta}}} \right); \end{aligned} \quad (7.34)$$

$$\begin{aligned} & \mathcal{N} \left( \left( \frac{g(\rho^{\alpha+\delta} y)}{\rho^{\alpha+\delta}} - \frac{g(\rho^\delta y)}{\rho^\delta}, u \right), t \right) \\ & \geq \mathcal{N}' \left( N(y, y, \dots, y), u, \frac{K \rho^\beta t}{\sum_{\sigma=0}^{\alpha-1} \frac{v^{(\sigma+\delta)\beta}}{\rho^{(\sigma+\delta)\beta}}} \right); \end{aligned} \quad (7.35)$$

$$\begin{aligned} & \mathcal{N} \left( \left( \frac{h(\tau^{\alpha+\delta} z)}{\tau^{\alpha+\delta}} - \frac{h(\tau^\delta z)}{\tau^\delta}, u \right), t \right) \\ & \geq \mathcal{N}' \left( N(z, z, \dots, z), u, \frac{K \tau^\beta t}{\sum_{\sigma=0}^{\alpha-1} \frac{v^{(\sigma+\delta)\beta}}{\tau^{(\sigma+\delta)\beta}}} \right); \end{aligned} \quad (7.36)$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$  and all  $\alpha, \delta > 0$ . Since

$$\begin{aligned} 0 < v < \kappa \quad & \text{and} \quad \sum_{\sigma=0}^{\alpha} \left( \frac{v}{\kappa} \right)^\sigma < \infty; \\ 0 < v < \rho \quad & \text{and} \quad \sum_{\sigma=0}^{\alpha} \left( \frac{v}{\rho} \right)^\sigma < \infty; \\ 0 < v < \tau \quad & \text{and} \quad \sum_{\sigma=0}^{\alpha} \left( \frac{v}{\tau} \right)^\sigma < \infty; \end{aligned}$$

the Cauchy criterion for convergence and (2QBFN5) implies that our sequences are Cauchy sequences. Due to the completeness of  $\mathcal{B}$ , this sequences converges to some points  $A_f; A_g; A_h$  respectively and its is defined by

$$\lim_{\alpha \rightarrow \infty} \mathcal{N} \left( \left( A_f(x) - \frac{f(\kappa^\alpha x)}{\kappa^\alpha}, u \right), t \right) = 1; \quad (7.37)$$

$$\lim_{\alpha \rightarrow \infty} \mathcal{N} \left( \left( A_g(y) - \frac{g(\rho^\alpha y)}{\rho^\alpha}, u \right), t \right) = 1; \quad (7.38)$$

$$\lim_{\alpha \rightarrow \infty} \mathcal{N} \left( \left( A_h(z) - \frac{h(\tau^\alpha z)}{\tau^\alpha}, u \right), t \right) = 1; \quad (7.39)$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ . Letting  $\delta = 0$  and letting  $\alpha \rightarrow \infty$  in (7.34), (7.35) and (7.36) respectively, we



arrive

$$\begin{aligned} & \mathcal{N}((A_f(x) - f(x), u), t) \\ & \geq \mathcal{N}'(N(x, x, \dots, x), u, K\kappa^{\beta-1}(\kappa - \nu)t); \\ & \mathcal{N}((A_g(y) - g(y), u), t) \\ & \geq \mathcal{N}'(E(y, y, \dots, y), u, K\rho^{\beta-1}(\rho - \nu)t); \\ & \mathcal{N}((A_h(z) - h(z), u), t) \\ & \geq \mathcal{N}'(O(z, z, \dots, z), u, K\tau^{\beta-1}(\tau - \nu)t); \end{aligned}$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ , respectively with  $\gamma = 1$ .

To show the mappings  $A_f(x); A_g(y); A_h(z)$  satisfies the functional equations (1.27), (1.28) and (1.29) by replacing

$$\begin{aligned} (x_1, x_2, \dots, x_n) &= (\kappa^\alpha x_1, \kappa^\alpha x_2, \dots, \kappa^\alpha x_n) \quad \text{in (7.1)} \\ (y_2, y_4, \dots, y_{2n}) &= (\rho^\alpha y_2, \rho^\alpha y_4, \dots, \rho^\alpha y_{2n}) \quad \text{in (7.2)} \\ (z_1, z_3, \dots, z_{2n-1}) &= (\tau^\alpha z_1, \tau^\alpha z_3, \dots, \tau^\alpha z_{2n-1}) \quad \text{in (7.3)} \end{aligned}$$

respectively, we arrive

$$\begin{aligned} & \mathcal{N}\left(\frac{1}{\kappa^\alpha} \left(f\left(\sum_{a=1}^n a \kappa^\alpha x_a\right) - \sum_{a=1}^n (a f(\kappa^\alpha x_a))\right), u, t\right) \\ & \geq \mathcal{N}'(N(\kappa^\alpha x_1, \kappa^\alpha x_2, \dots, \kappa^\alpha x_n), u, \kappa^{\alpha\beta} t); \quad (7.40) \\ & \mathcal{N}\left(\frac{1}{\rho^\alpha} \left(g\left(\sum_{a=1}^n 2a \rho^\alpha y_{2a}\right) - \sum_{a=1}^n (2a g(\rho^\alpha y_{2a}))\right), u, t\right) \\ & \geq \mathcal{N}'(E(\rho^\alpha y_2, \rho^\alpha y_4, \dots, \rho^\alpha y_{2n}), u, \rho^{\alpha\beta} t); \quad (7.41) \\ & \mathcal{N}\left(\frac{1}{\tau^\alpha} \left(h\left(\sum_{a=1}^n (2a-1) \tau^\alpha z_{2a-1}\right) - \sum_{a=1}^n ((2a-1) h(\tau^\alpha z_{2a-1}))\right), u, t\right) \\ & \geq \mathcal{N}'(O(\tau^\alpha z_1, \tau^\alpha z_3, \dots, \tau^\alpha z_{2n-1}), u, \tau^{\alpha\beta} t); \quad (7.42) \end{aligned}$$

for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ . Now

$$\begin{aligned} & \mathcal{N}\left(\left(A_f\left(\sum_{a=1}^n a x_a\right) - \sum_{a=1}^n (a A_f(x_a))\right), u, t\right) \\ & = \min\left\{\mathcal{N}\left(\left(A_f\left(\sum_{a=1}^n a x_a\right) - \frac{1}{\kappa^\alpha} f\left(\sum_{a=1}^n a \kappa^\alpha x_a\right)\right), u, \frac{Kt}{3}\right)\right\} \end{aligned}$$

$$\begin{aligned} & \mathcal{N}\left(\left(-\sum_{a=1}^n (a A_f(x_a))\right. \right. \\ & \quad \left. \left. + \frac{1}{\kappa^\alpha} \sum_{a=1}^n (a f(\kappa^\alpha x_a))\right), u, \frac{Kt}{3}\right), \\ & \mathcal{N}\left(\frac{1}{\kappa^\alpha} \left(f\left(\sum_{a=1}^n a \kappa^\alpha x_a\right) - \sum_{a=1}^n (a f(\kappa^\alpha x_a))\right), u, \frac{Kt}{3}\right)\right\}; \quad (7.43) \end{aligned}$$

$$\begin{aligned} & \mathcal{N}\left(\left(A_g\left(\sum_{a=1}^n 2a y_{2a}\right) - \sum_{a=1}^n (2a A_g(y_{2a}))\right), u, t\right) \\ & = \min\left\{\mathcal{N}\left(\left(A_g\left(\sum_{a=1}^n 2a y_{2a}\right) - \frac{1}{\rho^\alpha} g\left(\sum_{a=1}^n 2a \rho^\alpha y_{2a}\right)\right), u, \frac{Kt}{3}\right), \right. \end{aligned}$$

$$\begin{aligned} & \left. \mathcal{N}\left(\left(-\sum_{a=1}^n (2a A_g(y_{2a}))\right. \right. \right. \\ & \quad \left. \left. + \frac{1}{\rho^\alpha} \sum_{a=1}^n (2a g(\rho^\alpha y_{2a}))\right), u, \frac{Kt}{3}\right), \\ & \mathcal{N}\left(\frac{1}{\rho^\alpha} \left(g\left(\sum_{a=1}^n 2a \rho^\alpha y_{2a}\right) - \sum_{a=1}^n (2a g(\rho^\alpha y_{2a}))\right), u, \frac{Kt}{3}\right)\right\}; \quad (7.44) \end{aligned}$$

$$\begin{aligned} & \mathcal{N}\left(\left(A_h\left(\sum_{a=1}^n (2a-1) z_{2a-1}\right) - \sum_{a=1}^n ((2a-1) A_h(z_{2a-1}))\right), u, t\right) \\ & = \min\left\{\mathcal{N}\left(\left(A_h\left(\sum_{a=1}^n (2a-1) z_{2a-1}\right) - \frac{1}{\tau^\alpha} h\left(\sum_{a=1}^n (2a-1) \tau^\alpha z_{2a-1}\right)\right), u, \frac{Kt}{3}\right), \right. \end{aligned}$$

$$\begin{aligned} & \left. \mathcal{N}\left(\left(-\sum_{a=1}^n ((2a-1) A_h(z_{2a-1}))\right. \right. \right. \\ & \quad \left. \left. + \frac{1}{\tau^\alpha} \sum_{a=1}^n ((2a-1) h(\tau^\alpha z_{2a-1}))\right), u, \frac{Kt}{3}\right), \\ & \mathcal{N}\left(\frac{1}{\tau^\alpha} \left(h\left(\sum_{a=1}^n (2a-1) \tau^\alpha z_{2a-1}\right) - \sum_{a=1}^n ((2a-1) h(\tau^\alpha z_{2a-1}))\right), u, \frac{Kt}{3}\right)\right\}; \quad (7.45) \end{aligned}$$



for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ . Letting  $\alpha \rightarrow \infty$  in

(7.43) and using (7.37), (7.55), (7.4), (2QBFN2)

(7.44) and using (7.38), (7.56), (7.5), (2QBFN2)

(7.45) and using (7.39), (7.57), (7.6), (2QBFN2)

we reach the following equations

$$\begin{aligned} & \mathcal{N} \left( \left( A_f \left( \sum_{a=1}^n a x_a \right) - \sum_{a=1}^n (a A_f(x_a)), u \right), t \right) \\ &= \min \left\{ 1, 1, \mathcal{N}' \left( N(\kappa^\alpha x_1, \kappa^\alpha x_2, \dots, \kappa^\alpha x_n), u, \kappa^{\alpha\beta} t \right) \right\}; \end{aligned} \tag{7.46}$$

$$\begin{aligned} & \mathcal{N} \left( \left( A_g \left( \sum_{a=1}^n 2a y_{2a} \right) - \sum_{a=1}^n (2a A_g(y_{2a})), u \right), t \right) \\ &= \min \left\{ 1, 1, \mathcal{N}' \left( E(\rho^\alpha y_2, \rho^\alpha y_4, \dots, \rho^\alpha y_{2n}), u, \rho^{\alpha\beta} t \right) \right\}; \end{aligned} \tag{7.47}$$

$$\begin{aligned} & \mathcal{N} \left( \left( A_h \left( \sum_{a=1}^n (2a-1) z_{2a-1} \right) - \sum_{a=1}^n ((2a-1) A - h(z_{2a-1})), u \right), t \right) \\ &= \min \left\{ 1, 1, \mathcal{N}' \left( O(\tau^\alpha z_1, \tau^\alpha z_3, \dots, \tau^\alpha z_{2n-1}), u, \tau^{\alpha\beta} t \right) \right\}; \end{aligned} \tag{7.48}$$

$$A_f \left( \sum_{a=1}^n a x_a \right) = \sum_{a=1}^n (a A_f(x_a)); \tag{7.49}$$

$$A_g \left( \sum_{a=1}^n 2a y_{2a} \right) = \sum_{a=1}^n (2a A_g(y_{2a})); \tag{7.50}$$

$$A_h \left( \sum_{a=1}^n (2a-1) z_{2a-1} \right) = \sum_{a=1}^n ((2a-1) A - h(z_{2a-1})); \tag{7.51}$$

for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$ . Thus  $A_f(x); A_g(y); A_h(z)$  satisfies the functional equations (1.27), (1.28) and (1.29) for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$ .

In order to prove the being  $A_f(x); A_g(y); A_h(z)$  are unique. Let  $A'_f(x); A'_g(y); A'_h(z)$  be another additive mappings satisfying (7.10), (7.11), (7.12) and (7.13), (7.14), (7.15) respectively.

$$\begin{aligned} & \mathcal{N} \left( (A_f(x) - A'_f(x), u), t \right) \\ &= \mathcal{N} \left( \left( \frac{1}{\kappa^\alpha} (A_f(x) - A'_f(x)), u \right), t \right) \\ &\geq \min \left\{ \mathcal{N} \left( (A_f(\kappa^\alpha x) - f(\kappa^\alpha x), u), \frac{K \kappa^{\alpha\beta} t}{2} \right), \right. \\ &\quad \left. \mathcal{N} \left( (A'_f(\kappa^\alpha x) - f(\kappa^\alpha x), u), \frac{K \kappa^{\alpha\beta} t}{2} \right) \right\} \\ &\geq \mathcal{N}' \left( N(\kappa^\alpha x, \kappa^\alpha x, \dots, \kappa^\alpha x), u, \frac{K \kappa^{\beta-1} (\kappa - \nu) t}{2} \right) \\ &= \mathcal{N}' \left( N(x, x, \dots, x), u, \frac{K \kappa^{\beta-1} (\kappa - \nu) t}{2 \nu^\alpha} \right) \\ &\rightarrow 1 \text{ as } \alpha \rightarrow \infty; \end{aligned} \tag{7.52}$$

$$\begin{aligned} & \mathcal{N} \left( (A_g(y) - A'_g(y), u), t \right) \\ &= \mathcal{N} \left( \left( \frac{1}{\rho^\alpha} (A_g(y) - A'_g(y)), u \right), t \right) \\ &\geq \min \left\{ \mathcal{N} \left( (A_g(\rho^\alpha y) - g(\rho^\alpha y), u), \frac{K \rho^{\alpha\beta} t}{2} \right), \right. \\ &\quad \left. \mathcal{N} \left( (A'_g(\rho^\alpha y) - g(\rho^\alpha y), u), \frac{K \rho^{\alpha\beta} t}{2} \right) \right\} \\ &\geq \mathcal{N}' \left( E(\rho^\alpha y, \rho^\alpha y, \dots, \rho^\alpha y), u, \frac{K \rho^{\beta-1} (\rho - \nu) t}{2} \right) \\ &= \mathcal{N}' \left( E(y, y, \dots, y), u, \frac{K \rho^{\beta-1} (\rho - \nu) t}{2 \nu^\alpha} \right) \\ &\rightarrow 1 \text{ as } \alpha \rightarrow \infty; \end{aligned} \tag{7.53}$$

$$\begin{aligned} & \mathcal{N} \left( (A_h(z) - A'_h(z), u), t \right) \\ &= \mathcal{N} \left( \left( \frac{1}{\tau^\alpha} (A_h(z) - A'_h(z)), u \right), t \right) \\ &\geq \min \left\{ \mathcal{N} \left( (A_h(\tau^\alpha z) - h(\tau^\alpha z), u), \frac{K \tau^{\alpha\beta} t}{2} \right), \right. \\ &\quad \left. \mathcal{N} \left( (A'_h(\tau^\alpha z) - h(\tau^\alpha z), u), \frac{K \tau^{\alpha\beta} t}{2} \right) \right\} \\ &\geq \mathcal{N}' \left( E(\tau^\alpha z, \tau^\alpha z, \dots, \tau^\alpha z), u, \frac{K \tau^{\beta-1} (\tau - \nu) t}{2} \right) \\ &= \mathcal{N}' \left( E(z, z, \dots, z), u, \frac{K \tau^{\beta-1} (\tau - \nu) t}{2 \nu^\alpha} \right) \\ &\rightarrow 1 \text{ as } \alpha \rightarrow \infty; \end{aligned} \tag{7.54}$$

for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ . Hence the mappings  $A_f(x); A_g(y); A_h(z)$  are unique. Thus the theorem holds for  $\gamma = 1$ .



Also, if we replace

$$x = \frac{x}{\kappa} \text{ in (7.19);}$$

$$y = \frac{y}{\rho} \text{ in (7.20);}$$

$$z = \frac{z}{\tau} \text{ in (7.21);}$$

we arrive

$$\begin{aligned} & \mathcal{N} \left( \left( f(x) - \kappa f \left( \frac{x}{\kappa} \right), u \right), t \right) \\ & \geq \mathcal{N}' \left( N \left( \frac{x}{\kappa}, \frac{x}{\kappa}, \dots, \frac{x}{\kappa} \right), u, t \right); \end{aligned} \tag{7.55}$$

$$\begin{aligned} & \mathcal{N} \left( \left( g(y) - \rho g \left( \frac{y}{\rho} \right), u \right), t \right) \\ & \geq \mathcal{N}' \left( E \left( \frac{y}{\rho}, \frac{y}{\rho}, \dots, \frac{y}{\rho} \right), u, t \right); \end{aligned} \tag{7.56}$$

$$\begin{aligned} & \mathcal{N} \left( \left( h(z) - \tau h \left( \frac{z}{\tau} \right), u \right), t \right) \\ & \geq \mathcal{N}' \left( O \left( \frac{z}{\tau}, \frac{z}{\tau}, \dots, \frac{z}{\tau} \right), u, t \right); \end{aligned} \tag{7.57}$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ . The rest of proof is similar to that of the case  $\gamma = 1$ . Thus the theorem holds for  $\gamma = -1$  also. This completes the proof of the Theorem.  $\square$

The following corollary is an immediate consequence of Theorem 7.7 concerning the Hyers - Ulam, Hyers - Ulam - Rassias and J.M.Rassias stabilities of the functional equations (1.27), (1.28) and (1.29).

**Corollary 7.8.** *Let  $f, g, h : \mathcal{A} \rightarrow \mathcal{B}$  be a mapping satisfying the following inequalities*

$$\begin{aligned} & \mathcal{N} \left( \left( f \left( \sum_{a=1}^n a x_a \right) - \sum_{a=1}^n (a f(x_a)), u \right), t \right) \\ & \geq \begin{cases} \mathcal{N}'(\lambda, u, t), & \mu \neq 1; \\ \mathcal{N}' \left( \sum_{a=1}^n \lambda |x_a, u|^\mu, u, t \right), & \mu \neq 1; \\ \mathcal{N}' \left( \prod_{a=1}^n \lambda |x_a, u|^\mu + \sum_{a=1}^n \lambda |x_a, u|^{n\mu}, u, t \right), & n\mu \neq 1; \end{cases} \end{aligned} \tag{7.58}$$

$$\begin{aligned} & \mathcal{N} \left( \left( g \left( \sum_{a=1}^n 2a y_{2a} \right) - \sum_{a=1}^n (2a g(y_{2a})), u \right), t \right) \\ & \geq \begin{cases} \mathcal{N}'(\lambda, u, t), & \mu \neq 1; \\ \mathcal{N}' \left( \sum_{a=1}^n \lambda |y_{2a}, u|^\mu, u, t \right), & \mu \neq 1; \\ \mathcal{N}' \left( \prod_{a=1}^n \lambda |y_{2a}, u|^\mu + \sum_{a=1}^n \lambda |y_{2a}, u|^{n\mu}, u, t \right), & n\mu \neq 1; \end{cases} \end{aligned} \tag{7.59}$$

$$\begin{aligned} & \mathcal{N} \left( \left( h \left( \sum_{a=1}^n (2a-1) z_{2a-1} \right) \right. \right. \\ & \quad \left. \left. - \sum_{a=1}^n ((2a-1) h(z_{2a-1})), u \right), t \right) \end{aligned}$$

$$\geq \begin{cases} \mathcal{N}'(\lambda, u, t) \\ \mathcal{N}' \left( \sum_{a=1}^n \lambda |z_{2a-1}, u|^\mu, u, t \right), & \mu \neq 1; \\ \mathcal{N}' \left( \prod_{a=1}^n \lambda |z_{2a-1}, u|^\mu \right. \\ \quad \left. + \sum_{a=1}^n \lambda |z_{2a-1}, u|^{n\mu}, u, t \right), & n\mu \neq 1; \end{cases} \tag{7.60}$$

for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ , where  $\lambda$  and  $\mu$  are positive constants. Then there exists one and only additive mapping  $A_f, A_g, A_h : \mathcal{A} \rightarrow \mathcal{B}$  satisfying (1.27), (1.28), (1.29), and

$$\begin{aligned} & \mathcal{N} \left( (A_f(x) - f(x), u), t \right) \\ & \geq \begin{cases} \mathcal{N}'(\lambda, u, K\kappa^{\beta-1}|\kappa-1|t); \\ \mathcal{N}' \left( n\lambda|x, u|^\mu, u, K\kappa^{\beta-1}|\kappa-v|t \right); \\ \mathcal{N}' \left( (n+1)\lambda|x, u|^{n\mu}, u, K\kappa^{\beta-1}|\kappa-v|t \right); \end{cases} \end{aligned} \tag{7.61}$$

$$\begin{aligned} & \mathcal{N} \left( (A_g(y) - g(y), u), t \right) \\ & \geq \begin{cases} \mathcal{N}'(\lambda, u, K\rho^{\beta-1}|\rho-1|t); \\ \mathcal{N}' \left( n\lambda|x, u|^\mu, u, K\rho^{\beta-1}|\rho-v|t \right); \\ \mathcal{N}' \left( (n+1)\lambda|x, u|^{n\mu}, u, K\rho^{\beta-1}|\rho-v|t \right); \end{cases} \end{aligned} \tag{7.62}$$

$$\begin{aligned} & \mathcal{N} \left( (A_h(z) - h(z), u), t \right) \\ & \geq \begin{cases} \mathcal{N}'(\lambda, u, K\tau^{\beta-1}|\tau-1|t); \\ \mathcal{N}' \left( n\lambda|x, u|^\mu, u, K\tau^{\beta-1}|\tau-v|t \right); \\ \mathcal{N}' \left( (n+1)\lambda|x, u|^{n\mu}, u, K\tau^{\beta-1}|\tau-v|t \right); \end{cases} \end{aligned} \tag{7.63}$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ , respectively.

### 8. Stability Results In Random Quasi-Beta-2-Banach Space

In this section, we investigate the generalized Ulam - Hyers stability of the functional equations (1.27), (1.28) and (1.29) in random quasi- $\beta$ -2-Banach spaces using Hyers Method.

Now, we give basic definitions and notations in random quasi- $\beta$ -2-Banach space.

From now on,  $\Delta^+$  is the space of distribution functions, that is, the space of all mappings

$$F : R \cup \{-\infty, \infty\} \longrightarrow [0, 1],$$

such that  $F$  is leftcontinuous and nondecreasing on  $R$ ,  $F(0) = 0$  and  $F(+\infty) = 1$ .  $D^+$  is a subset of  $\Delta^+$  consisting of all functions  $F \in \Delta^+$  for which  $l^-F(+\infty) = 1$ , where  $l^-f(x)$  denotes the left limit of the function  $f$  at the point  $x$ , that is,

$$l^-f(x) = \lim_{t \rightarrow x^-} f(t).$$

The space  $\Delta^+$  is partially ordered by the usual pointwise ordering of functions, that is,  $F \leq G$  if and only if  $F(t) \leq G(t)$



for all  $t \in \mathbb{R}$ . The maximal element for  $\Delta^+$  in this order is the distribution function  $\varepsilon_0$  given by

$$\varepsilon_0(t) = \begin{cases} 0, & \text{if } t \leq 0, \\ 1, & \text{if } t \geq 0. \end{cases} \quad (8.1)$$

**Definition 8.1.** A mapping  $T : [0, 1] \times [0, 1] \rightarrow [0, 1]$  is called a continuous triangular norm (briefly, a continuous  $t$ -norm) if  $T$  satisfies the following conditions:

- (a)  $T$  is commutative and associative;
- (b)  $T$  is continuous;
- (c)  $T(a, 1) = a$  for all  $a \in [0, 1]$ ;
- (d)  $T(a, b) \leq T(c, d)$  whenever  $a \leq c$  and  $b \leq d$  for all  $a, b, c, d \in [0, 1]$ .

Typical examples of continuous  $t$ -norms are  $T_P(a, b) = ab$ ,  $T_M(a, b) = \min(a, b)$  and  $T_L(a, b) = \max(a + b - 1, 0)$  (the Lukasiewicz  $t$ -norm). Recall (see [21, 22]) that if  $T$  is a  $t$ -norm and  $x_n$  is a given sequence of numbers in  $[0, 1]$ , then  $T_{i=1}^n x_{n+i}$  is defined recurrently by

$$T_{i=1}^1 x_i = x_1 \quad \text{and} \quad T_{i=1}^n x_i = T(T_{i=1}^{n-1} x_i, x_n) \quad \text{for } n \geq 2.$$

$T_{i=n}^\infty x_i$  is defined as  $T_{i=1}^\infty x_{n+i}$ . It is known [22] that, for the Lukasiewicz  $t$ -norm, the following implication holds:

$$\lim_{n \rightarrow \infty} (T_L)_{i=1}^\infty x_{n+i} = 1 \iff \sum_{n=1}^\infty (1 - x_n) < \infty \quad (8.2)$$

**Definition 8.2.** A random quasi- $\beta$ -2-normed space is a quartile  $(X, \mathcal{R}, T, t)$ , where  $X$  is a vector space,  $T$  is a continuous  $t$ -norm,  $\mathcal{R}$  is a mapping from  $X$  into  $D^+$  and  $u \in D^+$  satisfying the following conditions:

- (2QBRN1)  $\mathcal{R}_x(u, t) = \varepsilon_0(u, t)$  for all  $t > 0$  if and only if  $x = 0$ ;
- (2QBRN2)  $\mathcal{R}_{\alpha x}(u, t) = \mathcal{R}_x(u, t/|\alpha|^\beta)$  for all  $x \in X$ , and  $\alpha \in \mathbb{R}$  with  $\alpha \neq 0$ ;
- (2QBRN3)  $\mathcal{R}_{x+y}(u, (t+s)) \geq T(\mathcal{R}_x(u, Kt), \mathcal{R}_y(u, Ks))$  for all  $x, y \in X$ ,  $t, s \geq 0$  and a constant  $K \geq 1$ .

**Example 8.3.** Every normed spaces  $(X, \|\cdot\|)$  defines a random quasi- $\beta$ -2-normed space  $(X, \mathcal{R}, T_M, u)$ , where

$$\mathcal{R}_x(u, t) = \frac{t}{t + \|x\|}$$

and  $T_M$  is the minimum  $t$ -norm.

**Definition 8.4.** Let  $(X, \mathcal{R}, T, u)$  be a random quasi- $\beta$ -2-normed space.

- (1) A sequence  $\{x_n\}$  in  $X$  is said to be convergent to a point  $x \in X$  if, for any  $\varepsilon > 0$  and  $\lambda > 0$ , there exists a positive integer  $N$  such that  $\mathcal{R}_{x_n-x}(\varepsilon) > 1 - \lambda$  for all  $n \geq N$ .

(2) A sequence  $\{x_n\}$  in  $X$  is called a Cauchy sequence if, for any  $\varepsilon > 0$  and  $\lambda > 0$ , there exists a positive integer  $N$  such that  $\mathcal{R}_{x_n-x_m}(\varepsilon) > 1 - \lambda$  for all  $n \geq m \geq N$  and all  $t > 0$ .

(3) A random quasi- $\beta$ -2-normed space  $(X, \mathcal{R}, T, u)$  is said to be complete if every Cauchy sequence in  $X$  is convergent to a point in  $X$ .

(4) A complete random quasi- $\beta$ -2-normed space  $(X, \mathcal{R}, T, u)$  is called random quasi- $\beta$ -2-Banach space.

To prove stability results, let us consider  $\mathcal{A}$  be a random quasi- $\beta$ -2-normed space and  $\mathcal{B}$  be a random quasi- $\beta$ -2-Banach space.

**Theorem 8.5.** Let  $f, g, h : \mathcal{A} \rightarrow \mathcal{B}$  be a mapping satisfying the following inequalities

$$\mathcal{R}'_f(\sum_{a=1}^n a x_a) - \sum_{a=1}^n (a f(x_a)) (u, t) \geq \mathcal{R}'_{\mathcal{N} x_1, x_2, \dots, x_n} (u, t); \quad (8.3)$$

$$\mathcal{R}'_g(\sum_{a=1}^n 2a y_{2a}) - \sum_{a=1}^n (2a g(y_{2a})) (u, t) \geq \mathcal{R}'_{\mathcal{E} y_2, y_4, \dots, y_{2n}} (u, t); \quad (8.4)$$

$$\mathcal{R}'_h(\sum_{a=1}^n (2a-1) z_{2a-1}) - \sum_{a=1}^n ((2a-1) h(z_{2a-1})) (u, t) \geq \mathcal{R}'_{\mathcal{O} z_1, z_3, \dots, z_{2n-1}} (u, t); \quad (8.5)$$

for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ , for which there exist a function  $\mathcal{R}'_{\mathcal{N}} : \mathcal{A}^n \rightarrow D^+$ ,  $\mathcal{R}'_{\mathcal{E}} : \mathcal{A}^{2n} \rightarrow D^+$  and  $\mathcal{R}'_{\mathcal{O}} : \mathcal{A}^{2n-1} \rightarrow D^+$  with the conditions

$$\begin{aligned} T_{\alpha=0}^\infty \mathcal{R}'_{\mathcal{N} \kappa^{\alpha\gamma x}, \kappa^{\alpha\gamma x}, \dots, \kappa^{\alpha\gamma x}} (u, \kappa^{\alpha\gamma t}) &= 1 \\ &= \lim_{\alpha \rightarrow \infty} \mathcal{R}'_{\mathcal{N} \kappa^{\alpha\gamma x_1}, \kappa^{\alpha\gamma x_2}, \dots, \kappa^{\alpha\gamma x_n}} (u, \kappa^{\alpha\gamma t}); \end{aligned} \quad (8.6)$$

$$\begin{aligned} T_{\alpha=0}^\infty \mathcal{R}'_{\mathcal{E} \rho^{\alpha\gamma y}, \rho^{\alpha\gamma y}, \dots, \rho^{\alpha\gamma y}} (u, \rho^{\alpha\gamma t}) &= 1 \\ &= \lim_{\alpha \rightarrow \infty} \mathcal{R}'_{\mathcal{E} \rho^{\alpha\gamma y_2}, \rho^{\alpha\gamma y_4}, \dots, \rho^{\alpha\gamma y_{2n}}} (u, \rho^{\alpha\gamma t}); \end{aligned} \quad (8.7)$$

$$\begin{aligned} T_{\alpha=0}^\infty \mathcal{R}'_{\mathcal{O} \tau^{\alpha\gamma z}, \tau^{\alpha\gamma z}, \dots, \tau^{\alpha\gamma z}} (u, \tau^{\alpha\gamma t}) &= 1 \\ &= \lim_{\alpha \rightarrow \infty} \mathcal{R}'_{\mathcal{O} \tau^{\alpha\gamma z_1}, \tau^{\alpha\gamma z_3}, \dots, \tau^{\alpha\gamma z_{2n-1}}} (u, \tau^{\alpha\gamma t}); \end{aligned} \quad (8.8)$$

for all  $x, y, z, x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$  all  $u \in \mathcal{B}$  and  $t > 0$  with  $\gamma = \pm 1$ . Then there exists one and only additive mapping  $A_f, A_g, A_h : \mathcal{A} \rightarrow \mathcal{B}$  satisfying (1.27), (1.28), (1.29), and

$$\mathcal{R}_{A_f(x)-f(x)}(u, t) \geq T_{\alpha=0}^\infty \mathcal{R}'_{\mathcal{N} \kappa^{\alpha\gamma x}, \kappa^{\alpha\gamma x}, \dots, \kappa^{\alpha\gamma x}} (u, K \kappa^{\beta\alpha\gamma t}); \quad (8.9)$$

$$\mathcal{R}_{A_g(y)-g(y)}(u, t) \geq T_{\alpha=0}^\infty \mathcal{R}'_{\mathcal{E} \rho^{\alpha\gamma y}, \rho^{\alpha\gamma y}, \dots, \rho^{\alpha\gamma y}} (u, \rho^{\beta\alpha\gamma t}); \quad (8.10)$$

$$\mathcal{R}_{A_h(z)-h(z)}(u, t) \geq T_{\alpha=0}^\infty \mathcal{R}'_{\mathcal{O} \tau^{\alpha\gamma z}, \tau^{\alpha\gamma z}, \dots, \tau^{\alpha\gamma z}} (u, \tau^{\beta\alpha\gamma t}); \quad (8.11)$$



for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ , respectively. The mappings  $A_f, A_g, A_h$  are respectively defined as

$$\mathcal{R}_{A_f(x)}(u, t) = \lim_{\alpha \rightarrow \infty} \mathcal{R}_{\frac{f(\kappa^\alpha \gamma_x)}{\kappa^\alpha \gamma}}(u, t); \tag{8.12}$$

$$\mathcal{R}_{A_g(y)}(u, t) = \lim_{\alpha \rightarrow \infty} \mathcal{R}_{\frac{g(\rho^\alpha \gamma_y)}{\rho^\alpha \gamma}}(u, t); \tag{8.13}$$

$$\mathcal{R}_{A_h(z)}(u, t) = \lim_{\alpha \rightarrow \infty} \mathcal{R}_{\frac{h(\tau^\alpha \gamma_z)}{\tau^\alpha \gamma}}(u, t); \tag{8.14}$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ .

*Proof.* Changing

$$(x_1, x_2, \dots, x_n) = (x, x, \dots, x) \text{ in (8.3);}$$

$$(y_2, y_4, \dots, y_{2n}) = (y, y, \dots, y) \text{ in (8.4);}$$

$$(z_1, z_3, \dots, z_{2n-1}) = (z, z, \dots, z) \text{ in (8.5);}$$

we arrive the following inequalities

$$\mathcal{R}_{f(\sum_{a=1}^n a x) - (\sum_{a=1}^n a) f(x)}(u, t) \geq \mathcal{R}'_{\mathcal{N}_{x,x,\dots,x}}(u, t); \tag{8.15}$$

$$\mathcal{R}_{g(\sum_{a=1}^n 2a y) - (\sum_{a=1}^n 2a) g(y)}(u, t) \geq \mathcal{R}'_{\mathcal{E}_{y,y,\dots,y}}(u, t); \tag{8.16}$$

$$\mathcal{R}_{h(\sum_{a=1}^n (2a-1) z) - (\sum_{a=1}^n (2a-1)) h(z)}(u, t) \geq \mathcal{R}'_{\mathcal{O}_{z,z,\dots,z}}(u, t); \tag{8.17}$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and all  $t > 0$ . Define

$$\sum_{a=1}^n a = \frac{n(n+1)}{2} = \kappa \text{ in (8.15);}$$

$$\sum_{a=1}^n 2a = n(n+1) = \rho \text{ in (8.16);}$$

$$\sum_{a=1}^n (2a-1) = n^2 = \tau \text{ in (8.17);}$$

we obtain the succeeding inequalities

$$\mathcal{R}_{f(\kappa x) - \kappa f(x)}(u, t) \geq \mathcal{R}'_{\mathcal{N}_{x,x,\dots,x}}(u, t); \tag{8.18}$$

$$\mathcal{R}_{g(\rho y) - \rho g(y)}(u, t) \geq \mathcal{R}'_{\mathcal{E}_{y,y,\dots,y}}(u, t); \tag{8.19}$$

$$\mathcal{R}_{h(\tau z) - \tau h(z)}(u, t) \geq \mathcal{R}'_{\mathcal{O}_{z,z,\dots,z}}(u, t); \tag{8.20}$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ . Using (2QBRN2) it follows from above inequalities

$$\mathcal{R}_{\frac{f(\kappa x)}{\kappa} - f(x)}(u, t) \geq \mathcal{R}'_{\mathcal{N}_{x,x,\dots,x}}(u, \kappa^\beta t); \tag{8.21}$$

$$\mathcal{R}_{\frac{g(\rho y)}{\rho} - g(y)}(u, t) \geq \mathcal{R}'_{\mathcal{E}_{y,y,\dots,y}}(u, \rho^\beta t); \tag{8.22}$$

$$\mathcal{R}_{\frac{h(\tau z)}{\tau} - h(z)}(u, t) \geq \mathcal{R}'_{\mathcal{O}_{z,z,\dots,z}}(u, \tau^\beta t); \tag{8.23}$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ . The rest of proof is similar tracing to that of Theorem 7.7.  $\square$

The following corollary is an immediate consequence of Theorem 8.5 concerning the Hyers - Ulam, Hyers - Ulam - Rassias and J.M.Rassias stabilities of the functional equations (1.27), (1.28) and (1.29).

**Corollary 8.6.** Let  $f, g, h : \mathcal{A} \rightarrow \mathcal{B}$  be a mapping satisfying the following inequalities

$$\begin{aligned} &\mathcal{R} \left( \left( f \left( \sum_{a=1}^n a x_a \right) - \sum_{a=1}^n (a f(x_a)), u \right), t \right) \\ &\geq \begin{cases} \mathcal{R}'(\lambda, u, t) & \mu \neq 1; \\ \mathcal{R}'(\sum_{a=1}^n \lambda |x_a, u|^\mu, u, t), & \\ \mathcal{R}'(\prod_{a=1}^n \lambda |x_a, u|^\mu + \sum_{a=1}^n \lambda |x_a, u|^{n\mu}, u, t), & n\mu \neq 1; \end{cases} \end{aligned} \tag{8.24}$$

$$\begin{aligned} &\mathcal{R} \left( \left( g \left( \sum_{a=1}^n 2a y_{2a} \right) - \sum_{a=1}^n (2a g(y_{2a})), u \right), t \right) \\ &\geq \begin{cases} \mathcal{R}'(\lambda, u, t) & \mu \neq 1; \\ \mathcal{R}'(\sum_{a=1}^n \lambda |y_{2a}, u|^\mu, u, t) & \\ \mathcal{R}'(\prod_{a=1}^n \lambda |y_{2a}, u|^\mu + \sum_{a=1}^n \lambda |y_{2a}, u|^{n\mu}, u, t), & n\mu \neq 1; \end{cases} \end{aligned} \tag{8.25}$$

$$\begin{aligned} &\mathcal{R} \left( \left( h \left( \sum_{a=1}^n (2a-1) z_{2a-1} \right) - \sum_{a=1}^n ((2a-1) h(z_{2a-1})), u \right), t \right) \\ &\geq \begin{cases} \mathcal{R}'(\lambda, u, t) & \mu \neq 1; \\ \mathcal{R}'(\sum_{a=1}^n \lambda |z_{2a-1}, u|^\mu, u, t), & \\ \mathcal{R}'(\prod_{a=1}^n \lambda |z_{2a-1}, u|^\mu + \sum_{a=1}^n \lambda |z_{2a-1}, u|^{n\mu}, u, t), & n\mu \neq 1; \end{cases} \end{aligned} \tag{8.26}$$

for all  $x_1, x_2, \dots, x_n, y_2, y_4, \dots, y_{2n}, z_1, z_3, \dots, z_{2n-1} \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ , where  $\lambda$  and  $\mu$  are positive constants. Then there exists one and only additive mapping  $A_f, A_g, A_h : \mathcal{A} \rightarrow \mathcal{B}$  satisfying (1.27), (1.28), (1.29), and

$$\begin{aligned} &\mathcal{R}((A_f(x) - f(x), u), t) \\ &\geq \begin{cases} \mathcal{R}'(\lambda, u, K\kappa^{\beta-1}|\kappa - 1|t); \\ \mathcal{R}'(n\lambda|x, u|^\mu, u, K\kappa^{\beta-1}|\kappa - \nu|t); \\ \mathcal{R}'((n+1)\lambda|x, u|^{n\mu}, u, K\kappa^{\beta-1}|\kappa - \nu|t); \end{cases} \end{aligned} \tag{8.27}$$

$$\begin{aligned} &\mathcal{R}((A_g(y) - g(y), u), t) \\ &\geq \begin{cases} \mathcal{R}'(\lambda, u, K\rho^{\beta-1}|\rho - 1|t); \\ \mathcal{R}'(n\lambda|x, u|^\mu, u, K\rho^{\beta-1}|\rho - \nu|t); \\ \mathcal{R}'((n+1)\lambda|x, u|^{n\mu}, u, K\rho^{\beta-1}|\rho - \nu|t); \end{cases} \end{aligned} \tag{8.28}$$

$$\begin{aligned} &\mathcal{R}((A_h(z) - h(z), u), t) \\ &\geq \begin{cases} \mathcal{R}'(\lambda, u, K\tau^{\beta-1}|\tau - 1|t); \\ \mathcal{R}'(n\lambda|x, u|^\mu, u, K\tau^{\beta-1}|\tau - \nu|t); \\ \mathcal{R}'((n+1)\lambda|x, u|^{n\mu}, u, K\tau^{\beta-1}|\tau - \nu|t); \end{cases} \end{aligned} \tag{8.29}$$

for all  $x, y, z \in \mathcal{A}$  and all  $u \in \mathcal{B}$  and  $t > 0$ , respectively.





## Acknowledgment

This work is dedicated to all the mathematicians and research scholars working in this field.

## References

- [1] J. Aczel *Lectures on Functional Equations and Their Applications*, Academic Press, New York (1966). MR:348020 (1967).
- [2] J. Aczel and J. Dhombres, *Functional Equations in Several Variables*, Cambridge Univ, Press, 1989.
- [3] T. Aoki, *On the stability of the linear transformation in Banach spaces*, J. Math. Soc. Japan. 2 (1950), 64-66.
- [4] M. Arunkumar, *Solution and stability of Arun-additive functional equations*, International Journal Mathematical Sciences and Engineering Applications, 4, No.3, August 2010.
- [5] M. Arunkumar, C. Leela Sabari, *Solution and stability of a functional equation originating from a chemical equation*, International Journal Mathematical Sciences and Engineering Applications, f 5 (2) (2011), 1-8..
- [6] M. Arunkumar, S. Karthikeyan, *Solution and Stability of n-dimensional Additive functional equation*, International Journal of Applied Mathematics, 25 (2), (2012), 163-174.
- [7] M. Arunkumar, S. Hema latha, C. Devi Shaymala Mary, *Functional equation originating from arithmetic Mean of consecutive terms of an arithmetic Progression are stable in banach space: Direct and fixed point method*, JP Journal of Mathematical Sciences, 3, Issue 1, 2012, Pages 27-43.
- [8] M. Arunkumar, G. Vijayanandhraj, S. Karthikeyan, *Solution and Stability of a Functional Equation Originating From n Consecutive Terms of an Arithmetic Progression*, Universal Journal of Mathematics and Mathematical Sciences, 2, No. 2, (2012), 161-171.
- [9] M. Arunkumar, *Solution and stability of modified additive and quadratic functional equation in generalized 2-normed spaces*, International Journal Mathematical Sciences and Engineering Applications, 7 No. I (2013), 383-391.
- [10] M. Arunkumar, *Functional equation originating from sum of First natural numbers is stable in Cone Banach Spaces: Direct and Fixed Point Methods*, International Journal of Information Science and Intelligent System (IJISIS ISSN:2307-9142), 2 No. 4 (2013), 89 - 104.
- [11] M. Arunkumar, *Stability of n- dimensional Additive functional equation in Generalized 2 - normed space*, Demonstrato Mathematica 49 (3), (2016), 319 - 33.
- [12] M. Arunkumar, P. Agilan, *Additive functional equation and inequality are Stable in Banach space and its applications*, Malaya Journal of Matematik (MJM), Vol 1, Issue 1, (2013), 10-17.
- [13] A. Malceski, R. Malceski, K. Anevska and S. Malceski, *Remark about Quasi 2-Normed Space*, Applied Mathematical Sciences, 9, 2015, no. 55, 2717-2727.
- [14] S. Czerwik, *Functional Equations and Inequalities in Several Variables*, World Scientific, River Edge, NJ, 2002.
- [15] S. Gähler *2 Metrische Raume und ihre Topologische Struktur*, Math. Nachr., 26 (1963), 115-148.
- [16] S. Gähler, *Lineare 2-Normierte Raume*, Math. Nachr., 28 (1964) 1-43.
- [17] S. Gähler, *Über 2-Banach-Raume*, Math. Nachr. 42 (1969), 335-347.
- [18] P. Gävruta, *A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings*, J. Math. Anal. Appl., 184 (1994), 431-436.
- [19] D. H. Hyers, *On the stability of the linear functional equation*, Proc. Nat. Acad. Sci. USA. 27 (1941), 222-224.
- [20] D. H. Hyers, G. Isac and Th.M. Rassias, *Stability of functional equations in several variables*, Birkhauser, Basel, 1998.
- [21] O. Hadzic and E. Pap, *Fixed Point Theory in Probabilistic Metric Spaces*, vol. 536 of Mathematics and Its Applications, Kluwer Academic, Dordrecht, The Netherlands, 2001.
- [22] O. Hadzic, E. Pap and M. Budincevic, *Countable extension of triangular norms and their applications to the fixed point theory in probabilistic metric spaces*, Kybernetika. 38, no. 3 (2002), 363-382.
- [23] S.M. Jung, *Hyers-Ulam-Rassias Stability of Functional Equations in Mathematical Analysis*, Hadronic Press, Palm Harbor, 2001.
- [24] Pl. Kannappan, *Functional Equations and Inequalities with Applications*, Springer Monographs in Mathematics, 2009.
- [25] A. K. Katsaras, *Fuzzy topological vector spaces II*, Fuzzy Sets and Systems, 12 (1984), 143-154.
- [26] M. Kir and M. Acikgoz, *A study involving the completion of quasi 2-normed space*, Int. J. Anal. (2013). <http://dx.doi.org/10.1155/2013/512372>.
- [27] D.O. Lee, *Hyers-Ulam stability of an addtiive type functional equation*, J. Appl. Math. and Computing, 13 (2003) no.1-2, 471-477.
- [28] Z. Lewandowska, *Generalized 2-normed spaces*, Stuspske Prace Matematyczno-Fizyczne, 1 (2001), no.4, 33-40.
- [29] L. Maligranda, *A result of Tosio Aoki about a generalization of Hyers-Ulam stability of additive functions- a question of priority*, Aequationes Math., 75 (2008), 289-296.
- [30] M.J.Rassias, M. Arunkumar, E.Sathya, *ai Type n- Variable Multi n- Dimensional Additive Functional Equation*, Malaya Journal of Matematik, 5(2) (2017), 278-292.
- [31] J. M. Rassias, *On approximately of approximately linear mappings by linear mappings*, J. Funct. Anal. USA, 46, (1982) 126-130.
- [32] Th. M. Rassias, *On the stability of the linear mapping in*



- Banach spaces*, Proc. Amer. Math. Soc., 72 (2) (1978) 297-300.
- [33] K. Ravi, M. Arunkumar, *On a  $n$ -dimensional additive Functional Equation with fixed point Alternative*, Proceedings of ICMS 2007, 314-330.
- [34] K. Ravi, M. Arunkumar and J.M. Rassias, *On the Ulam stability for the orthogonally general Euler-Lagrange type functional equation*, International Journal of Mathematical Sciences, Autumn 2008 Vol.3, No. 08, 36-47.
- [35] S. Rolewicz, *Metric Linear Spaces*, Reidel, Dordrecht, 1984.
- [36] P. K. Sahoo, Pl. Kannappan, *Introduction to Functional Equations*, Chapman and Hall/CRC Taylor and Francis Group, 2011.
- [37] B. Schweizer and A. Sklar, *Probabilistic Metric Spaces*, North-Holland Series in Probability and Applied Mathematics, North-Holland Publishing, New York, NY, USA, 1983.
- [38] A. N. Sherstnev, *On the notion of a random normed space*, Doklady Akademii Nauk SSSR. 149 (1963), 280-283 (Russian).
- [39] J. Tabor, *Stability of the Cauchy functional equation in quasi-Banach spaces*, Ann. Polon. Math. **83** (2004) 243-255.
- [40] S. M. Ulam, *Problems in Modern Mathematics*, Chapter VI, Science Ed., Wiley, New York, 1940.
- [41] A. White, *2-Banach spaces*, Doctorial Diss., St. Louis Univ., 1968.
- [42] A. White, *2-Banach spaces*, Math. Nachr. 42 (1969), 43-60.

\*\*\*\*\*  
ISSN(P):2319 – 3786  
Malaya Journal of Matematik  
ISSN(O):2321 – 5666  
\*\*\*\*\*

